

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Memoranda of the Field Experiments at Rothamsted: May 1881



[Full Table of Content](#)

Default Title

Rothamsted Research

Rothamsted Research (1882) *Default Title* ; Memoranda Of The Field Experiments At Rothamsted: May 1881, pp 1 - 27 - DOI: <https://doi.org/10.23637/ERADOC-1-245>

MEMORANDA
OF THE
ORIGIN, PLAN, AND RESULTS
OF THE
FIELD AND OTHER EXPERIMENTS,
CONDUCTED
On the Farm and in the Laboratory of
JOHN BENNET LAWES, LL.D., F.R.S.,
AT ROTHAMSTED, HERTS;
ALSO A STATEMENT OF THE
PRESENT AND PREVIOUS CROPPING, ETC.,
OF THE
ARABLE LAND NOT UNDER EXPERIMENT.

MAY, 1881.

CONTENTS.

	PAGE
Origin, Scope, and Plan of the Rothamsted Experiments	3-5
Lists of Papers published	6-8
Experiments on Permanent Meadow Land; The Park	9
Experiments on Barley; Hoos Field	10
Experiments on Wheat; Broadbalk Field	11
Experiments on Oats; Geescroft Field	12
Experiments on Beans; Geescroft Field	13
Experiments on Clover; Hoos Field	13-15
Experiments with various Leguminous Plants	16
Experiments on Turnips; Barn Field	17
Experiments on Sugar-beet; Barn Field	18-19
Experiments on Mangold-Wurzel; Barn Field	20-21
Experiments on Potatos; Hoos Field	22-23
Experiments on Rotation; Agdell Field	24
Experiments with different descriptions of Wheat; Rickyard Field	25
Cropping, &c., of the Arable Land not under Experiment	26-27

B

MEMORANDA
 OF THE
 ORIGINAL PLAN AND RESULTS
 OF THE
 FIELD AND OTHER EXPERIMENTS
 ON THE PLANT AND IN THE LABORATORY OF
 JOHN BRETHERTON, M.D., F.R.S.
 AT BOTHAMPTON, HANTS.
 WITH A STATEMENT OF THE
 PROGRESS AND PREVIOUS CHRONIC, ETC.
 OF THE
 DISEASE AND NOT UNDER EXPERIMENT.

MAY 1841

CONTENTS

1-2	Introductory Remarks
3-4	On the Nature of the Disease
5-6	On the Progress of the Disease
7-8	On the Symptoms of the Disease
9-10	On the Treatment of the Disease
11-12	On the Prevention of the Disease
13-14	On the Origin of the Disease
15-16	On the Duration of the Disease
17-18	On the Mortality of the Disease
19-20	On the Recovery of the Disease
21-22	On the Cause of the Disease
23-24	On the Effect of the Disease
25-26	On the Nature of the Disease
27-28	On the Progress of the Disease
29-30	On the Symptoms of the Disease
31-32	On the Treatment of the Disease
33-34	On the Prevention of the Disease
35-36	On the Origin of the Disease
37-38	On the Duration of the Disease
39-40	On the Mortality of the Disease
41-42	On the Recovery of the Disease
43-44	On the Cause of the Disease
45-46	On the Effect of the Disease
47-48	On the Nature of the Disease
49-50	On the Progress of the Disease
51-52	On the Symptoms of the Disease
53-54	On the Treatment of the Disease
55-56	On the Prevention of the Disease
57-58	On the Origin of the Disease
59-60	On the Duration of the Disease
61-62	On the Mortality of the Disease
63-64	On the Recovery of the Disease
65-66	On the Cause of the Disease
67-68	On the Effect of the Disease
69-70	On the Nature of the Disease
71-72	On the Progress of the Disease
73-74	On the Symptoms of the Disease
75-76	On the Treatment of the Disease
77-78	On the Prevention of the Disease
79-80	On the Origin of the Disease
81-82	On the Duration of the Disease
83-84	On the Mortality of the Disease
85-86	On the Recovery of the Disease
87-88	On the Cause of the Disease
89-90	On the Effect of the Disease
91-92	On the Nature of the Disease
93-94	On the Progress of the Disease
95-96	On the Symptoms of the Disease
97-98	On the Treatment of the Disease
99-100	On the Prevention of the Disease
101-102	On the Origin of the Disease
103-104	On the Duration of the Disease
105-106	On the Mortality of the Disease
107-108	On the Recovery of the Disease
109-110	On the Cause of the Disease
111-112	On the Effect of the Disease
113-114	On the Nature of the Disease
115-116	On the Progress of the Disease
117-118	On the Symptoms of the Disease
119-120	On the Treatment of the Disease
121-122	On the Prevention of the Disease
123-124	On the Origin of the Disease
125-126	On the Duration of the Disease
127-128	On the Mortality of the Disease
129-130	On the Recovery of the Disease
131-132	On the Cause of the Disease
133-134	On the Effect of the Disease
135-136	On the Nature of the Disease
137-138	On the Progress of the Disease
139-140	On the Symptoms of the Disease
141-142	On the Treatment of the Disease
143-144	On the Prevention of the Disease
145-146	On the Origin of the Disease
147-148	On the Duration of the Disease
149-150	On the Mortality of the Disease
151-152	On the Recovery of the Disease
153-154	On the Cause of the Disease
155-156	On the Effect of the Disease
157-158	On the Nature of the Disease
159-160	On the Progress of the Disease
161-162	On the Symptoms of the Disease
163-164	On the Treatment of the Disease
165-166	On the Prevention of the Disease
167-168	On the Origin of the Disease
169-170	On the Duration of the Disease
171-172	On the Mortality of the Disease
173-174	On the Recovery of the Disease
175-176	On the Cause of the Disease
177-178	On the Effect of the Disease
179-180	On the Nature of the Disease
181-182	On the Progress of the Disease
183-184	On the Symptoms of the Disease
185-186	On the Treatment of the Disease
187-188	On the Prevention of the Disease
189-190	On the Origin of the Disease
191-192	On the Duration of the Disease
193-194	On the Mortality of the Disease
195-196	On the Recovery of the Disease
197-198	On the Cause of the Disease
199-200	On the Effect of the Disease
201-202	On the Nature of the Disease
203-204	On the Progress of the Disease
205-206	On the Symptoms of the Disease
207-208	On the Treatment of the Disease
209-210	On the Prevention of the Disease
211-212	On the Origin of the Disease
213-214	On the Duration of the Disease
215-216	On the Mortality of the Disease
217-218	On the Recovery of the Disease
219-220	On the Cause of the Disease
221-222	On the Effect of the Disease
223-224	On the Nature of the Disease
225-226	On the Progress of the Disease
227-228	On the Symptoms of the Disease
229-230	On the Treatment of the Disease
231-232	On the Prevention of the Disease
233-234	On the Origin of the Disease
235-236	On the Duration of the Disease
237-238	On the Mortality of the Disease
239-240	On the Recovery of the Disease
241-242	On the Cause of the Disease
243-244	On the Effect of the Disease
245-246	On the Nature of the Disease
247-248	On the Progress of the Disease
249-250	On the Symptoms of the Disease
251-252	On the Treatment of the Disease
253-254	On the Prevention of the Disease
255-256	On the Origin of the Disease
257-258	On the Duration of the Disease
259-260	On the Mortality of the Disease
261-262	On the Recovery of the Disease
263-264	On the Cause of the Disease
265-266	On the Effect of the Disease
267-268	On the Nature of the Disease
269-270	On the Progress of the Disease
271-272	On the Symptoms of the Disease
273-274	On the Treatment of the Disease
275-276	On the Prevention of the Disease
277-278	On the Origin of the Disease
279-280	On the Duration of the Disease
281-282	On the Mortality of the Disease
283-284	On the Recovery of the Disease
285-286	On the Cause of the Disease
287-288	On the Effect of the Disease
289-290	On the Nature of the Disease
291-292	On the Progress of the Disease
293-294	On the Symptoms of the Disease
295-296	On the Treatment of the Disease
297-298	On the Prevention of the Disease
299-300	On the Origin of the Disease
301-302	On the Duration of the Disease
303-304	On the Mortality of the Disease
305-306	On the Recovery of the Disease
307-308	On the Cause of the Disease
309-310	On the Effect of the Disease
311-312	On the Nature of the Disease
313-314	On the Progress of the Disease
315-316	On the Symptoms of the Disease
317-318	On the Treatment of the Disease
319-320	On the Prevention of the Disease
321-322	On the Origin of the Disease
323-324	On the Duration of the Disease
325-326	On the Mortality of the Disease
327-328	On the Recovery of the Disease
329-330	On the Cause of the Disease
331-332	On the Effect of the Disease
333-334	On the Nature of the Disease
335-336	On the Progress of the Disease
337-338	On the Symptoms of the Disease
339-340	On the Treatment of the Disease
341-342	On the Prevention of the Disease
343-344	On the Origin of the Disease
345-346	On the Duration of the Disease
347-348	On the Mortality of the Disease
349-350	On the Recovery of the Disease
351-352	On the Cause of the Disease
353-354	On the Effect of the Disease
355-356	On the Nature of the Disease
357-358	On the Progress of the Disease
359-360	On the Symptoms of the Disease
361-362	On the Treatment of the Disease
363-364	On the Prevention of the Disease
365-366	On the Origin of the Disease
367-368	On the Duration of the Disease
369-370	On the Mortality of the Disease
371-372	On the Recovery of the Disease
373-374	On the Cause of the Disease
375-376	On the Effect of the Disease
377-378	On the Nature of the Disease
379-380	On the Progress of the Disease
381-382	On the Symptoms of the Disease
383-384	On the Treatment of the Disease
385-386	On the Prevention of the Disease
387-388	On the Origin of the Disease
389-390	On the Duration of the Disease
391-392	On the Mortality of the Disease
393-394	On the Recovery of the Disease
395-396	On the Cause of the Disease
397-398	On the Effect of the Disease
399-400	On the Nature of the Disease

ORIGIN, SCOPE, AND PLAN,
OF THE
ROTHAMSTED EXPERIMENTS.

THE following statement of the origin, scope, and plan, of the Rothamsted Investigations, was drawn up in answer to a circular letter issued by a Committee appointed to arrange for the commemoration of the twenty-fifth anniversary of the establishment of the First Experimental Station in Germany (Möckern), which was held in Leipzig in September 1877. The precise form of the statement depended on the order and form of the questions to which it is an answer. It has already been published in German, almost in full, with the series of reports of other Experimental Stations, which was issued at the time of the Jubilee Meeting.¹ It has been thought that it would be of some interest as an introduction to the *Memoranda of the Plan and Results of the Field Experiments, &c.*, annually issued at Rothamsted, and which here follow it. To the general statement, which, with a few slight alterations correcting it up to date, is given in the form in which it was originally drawn up, are appended lists of the titles of all the papers already published, with full reference to the Journals in which they appeared.

Mr. Lawes was the founder of the Rothamsted Experimental Station. He commenced experiments with different manuring substances, first with plants in pots, and afterwards in the field, soon after entering into possession of his hereditary property at Rothamsted² in 1834. The researches of De Saussure on vegetation were the chief subjects of his study to this end. Of all the experiments so made, those in which the neutral phosphate of lime, in bones, bone-ash, and apatite, was rendered soluble by means of sulphuric acid, and the mixture applied for root-crops, gave the most striking results. The results obtained on a small scale in 1837, 1838, and 1839, were such as to lead to more extensive trials in the field in 1840 and 1841, and subsequently.

In 1843 more systematic field experiments were commenced; and a barn, which had previously been partially applied to laboratory purposes, became almost exclusively devoted to agricultural investigations. The foundation of the Rothamsted Experimental Station may be said to date from that time (1843).

The Rothamsted station has up to the present time been entirely disconnected from any external organization, and has been maintained entirely by Mr. Lawes. He has further set apart a sum of £100,000, and certain areas of land, for the continuance of the investigations after his death.

In 1854-5 a new laboratory was built, by public subscription of agriculturists, and presented to Mr. Lawes, in July 1855, from which date the old barn-laboratory was abandoned, and the new one has been occupied.

From June 1843, up to the present time, Dr. J. H. Gilbert has been associated with Mr. Lawes, and has had the direction of the laboratory.

The number of assistants and other helps has increased from time to time. At first only one laboratory man was employed; but very soon a chemical assistant was necessary, and next a computer and record-keeper.

During the last twenty-five years the staff has consisted of—
One or two, and sometimes three, chemists.

Two or three general assistants. One of these is usually employed in routine chemical work, but sometimes in more general work. The chief occupation of the general assistants is to superintend the field experiments—that is, the making of the manures, the measurement of the plots, the application of the

manures, and the harvesting of the crops; also the taking of samples, the preparation of them for preservation or analysis, and the determinations of dry matter, ash, &c. These assistants also superintend any experiments made with animals. There are now about 30,000 bottles of samples of experimentally-grown vegetable produce, of animal products, of ashes, or of soils, stored in the laboratory.

A botanical assistant is also occasionally employed, with from three to six boys under him, and with him is generally associated one of the permanent general assistants, who at other times undertakes the botanical work.

Two or three (for some time past three) computers and record-keepers have been occupied in calculating and tabulating field, feeding, and laboratory results, copying, &c.

One, and sometimes two, laboratory men are employed.

Besides the permanent laboratory staff, chemical assistance is frequently engaged in London, or elsewhere; and, in this way, for some years past, Mr. R. Richter, of Berlin, has been almost constantly occupied with analytical work sent from Rothamsted.

The field experiments, and occasionally feeding experiments, also employ a considerable but a very variable number of agricultural labourers.

Nothing has been done at Rothamsted in the way of manure-, feeding-stuff-, or seed-control.

The investigations may be classed under two heads:—

I.—FIELD EXPERIMENTS, EXPERIMENTS ON
VEGETATION, Etc.

The general scope and plan of the field experiments has been:—

To grow some of the most important crops of rotation, each separately, year after year, for many years in succession on the same land, without manure, with farmyard-manure, and with a great variety of chemical manures; the same description of manure being, as a rule, applied year after year on the same plot. Experiments on an actual course of rotation, without manure, and with different manures, have also been made. In this way field experiments have been conducted as follows:—

(1) Die landwirthschaftlichen Versuchs-Stationen. Band xxii. 1877.

(2) Rothamsted is in Hertfordshire, twenty-five miles from London, on the Midland Railway; station, Harpenden.

On Wheat, thirty-eight years in succession; 13 acres, 37 plots, many of which are duplicates of others.

On Barley, thirty years in succession; $4\frac{1}{2}$ acres, 29 plots.

On Oats, ten years (including one year fallow); $\frac{3}{4}$ acre, 6 plots.

On Wheat, alternated with fallow, thirty years; 1 acre, 2 plots.

On different descriptions of Wheat, fourteen years; 4-8 acres (each year in a different field), now more than 20 plots.

On Beans, thirty-two years (including one year Wheat and five years fallow); $1\frac{1}{4}$ acre, 10 plots. Also twenty-seven years; 1 acre, 5 plots.

On Beans, alternated with Wheat, twenty-eight years; 1 acre 10 plots.

On Clover, with fallow or a corn-crop intervening, twenty-six years; 3 acres, 18 plots. The land is now devoted to experiments with various Leguminous plants, commenced in 1878.

On Turnips, twenty-eight years (including three years' barley); about 8 acres, 40 plots.

On Sugar Beet, five years; about 8 acres, 41 plots.

On Mangold-Wurzel, six years (in progress); about 8 acres, 41 plots.

On Potatos, six years (in progress); 2 acres, 10 plots.

On Rotation, thirty-four years; about $2\frac{1}{2}$ acres, 12 plots.

On permanent Grass-land, twenty-six years; about 7 acres, 22 plots.

Comparative experiments with different manures have also been made on other descriptions of soil, in other localities.

Samples of all the experimental crops are taken, and brought to the laboratory. Weighed portions of each are partially dried, and preserved for future reference or analysis. Duplicate weighed portions of each are dried at 100° C., the dry matter determined, and then burnt to ash on platinum sheets in cast-iron muffles. The quantities of ash are determined and recorded, and the ashes themselves are preserved for reference, or analysis.

In a large proportion of the samples the nitrogen is determined; and in some the amount existing as albuminoids, amides, and nitric acid.

In selected cases, illustrating the influence of season, manures, exhaustion, &c., complete ash-analyses have been made, numbering in all more than 600.

Also in selected cases, illustrating the influence of season and manuring, quantities of the experimentally-grown Wheat grain have been sent to the mill, and the proportion and composition of the different mill-products determined.

In the Sugar Beet, Mangold-Wurzel, and Potatos, the sugar in the juice has in most cases been determined by polariscope, and frequently by copper also.

In the case of the experiments on the mixed herbage of permanent grass land, besides the samples taken for the determination of the chemical composition (dry matter, ash, nitrogen, woody fibre, fatty matter, and composition of ash), carefully averaged samples have frequently been taken for the determination of the botanical composition. In this way, on four occasions, at intervals of five years—viz., in 1862, 1867, 1872, and 1877—a sample of the produce of each plot was taken, and submitted to careful botanical separation, and the percentage, by weight, of each species in the mixed herbage determined. Partial separations, in the case of samples from selected plots (frequently of both first and second crops), have also been made in other years.

INVESTIGATION OF SOILS.

Samples of the soils of most of the experimental plots have been taken from time to time, generally to the depth of 9, 18, and 27 inches, but sometimes to twice this depth. In this way about 600 samples have been taken, submitted to partial mechanical separation, and portions of the mould have been care-

fully prepared and preserved for analysis. In a large proportion of the samples the loss on drying at different temperatures, and at ignition, has been determined. In most the nitrogen determinable by burning with soda-lime has been estimated. In some the carbon, and in some the nitrogen as nitric acid, have been determined. Some experiments have also been made on the comparative absorptive capacity (for water and ammonia) of different soils and subsoils. The systematic investigation of the amount, and condition, of the nitrogen, and of some of the more important mineral constituents, of the soils of the different plots, and from different depths, is now in progress or contemplated.

RAINFALL AND DRAINAGE.

Almost from the commencement of the experiments the rainfall has been measured—for twenty-eight years in a gauge of one-thousandth of an acre area, as well as in an ordinary small funnel-gauge of 5 inches diameter. An 8-inch "Board of Trade" copper gauge is also now in use, commencing January 1, 1881. From time to time the nitrogen, as ammonia and as nitric acid, has been determined in the rain waters. The chlorine has, also, in some cases been determined.

Three "drain gauges," also of one-thousandth of an acre each, for the determination of the quantity and composition of the water percolating respectively through 20 inches, 40 inches, and 60 inches depth of soil (with its subsoil in natural state of consolidation) have also been constructed. A more numerous series of smaller "drain gauges," arranged for the investigation of the influence of different crops, and of different manures, has been constructed; but they have been found to be not sufficiently water-tight. Each of the differently manured plots of the permanent experimental Wheat-field having a separate pipe-drain, the drainage-waters have been and are frequently collected and analysed.

Professor Frankland has determined the nitrogen, as ammonia, as nitric acid, and as organic nitrogen, and also some other constituents, in many samples both of the rain and of the various drainage waters collected at Rothamsted. Dr. Voelcker also has determined the combined nitrogen, and likewise the incombustible constituents, in many of the drainage waters.

The nitrogen existing as nitric acid, sometimes that in other forms, and also some other constituents, are, and for some time past have been, determined periodically, in both the rain and the various drainage waters.

AMOUNT OF WATER TRANSPIRED BY PLANTS.

For several years in succession, experiments were made to determine the amount of water given off by plants during their growth. In this way various plants, including representatives of the gramineous, the leguminous, and other families, have been experimented upon. Similar experiments have also been made with various evergreen and deciduous trees.

BOTANICAL CHARACTERISTICS, &c.

Having regard to the difference in the character and amount of the constituents assimilated by plants of different botanical relationships, under equal external conditions, or by the same description of plants, under varying conditions, observations have been made on the character and range of the roots of different plants, and on their relative development of stem, leaf, &c. In the case of various crops, but more especially with Wheat and Beans, samples have been taken at different stages of growth, and the composition determined, in more or less detail, sometimes of the entire plant, and sometimes of the separated parts. In a few cases the amounts of dry matter, ash, nitrogen, &c., in the above-ground growth of a given area, at different stages of development, have been determined. The amounts of stubble of different crops have also occasionally been estimated.

ASSIMILATION OF FREE NITROGEN.

Experiments were made for several years in succession to determine whether plants assimilate free or uncombined nitrogen, and also various collateral points. Plants of the gramineous, the leguminous, and of other families were operated upon. The late Dr. Pugh took a prominent part in this inquiry.

II.—EXPERIMENTS ON ANIMALS, ETC.

Experiments with the animals of the farm were commenced early in 1847, and have been continued, at intervals, up to the present time.

The following points have been investigated:—

1. The amount of food, and of its several constituents, consumed in relation to a given live weight of animal within a given time.
2. The amount of food, and of its several constituents, consumed to produce a given amount of increase in live weight.
3. The proportion, and relative development, of the different organs or parts of different animals.
4. The proximate and ultimate composition of the animals in different conditions as to age and fatness, and the probable composition of their increase in live weight during the fattening process.
5. The composition of the solid and liquid excreta (the manure) in relation to that of the food consumed.
6. The loss or expenditure of constituents by respiration and the cutaneous exhalations—that is, in the mere sustenance of living meat-and-manure-making machine.

The general plan of experimenting was as follows:—

To provide data as to the amount of food, or its several constituents, consumed in relation to a given live weight of animal within a given time, and to produce a given amount of increase in live weight, several hundred animals—oxen, sheep, and pigs—have been experimented upon. Selected lots of animals were supplied, for many weeks, or for months consecutively, with weighed quantities of foods, selected and allotted according to the special point under inquiry. The composition of the foods was determined by analysis. The weights of the animals were taken at the commencement, at intervals during the progress, and at the conclusion of the experiment.

The amount, and relative development, of the different organs and parts were determined in two calves, two heifers, fourteen bullocks, one lamb, 249 sheep, and fifty-nine pigs.

The percentage of water, mineral matter, fat, and nitrogenous substance, were determined in certain separated parts, and in the entire bodies, of ten animals—namely, one calf, two oxen, one lamb, four sheep, and two pigs. Complete analyses of the ashes, respectively, of the entire carcasses, of the mixed internal and other “offal” parts, and of the entire bodies, of each of these ten animals have also been made.

From the data provided, as just described, as to the chemical composition of the different descriptions of animal, in different conditions as to age and fatness, the composition of the increase whilst fattening, and the relation of the constituents stored up in increase to those consumed in food, have been estimated.

To ascertain the composition of the manure in relation to that of the food consumed, oxen, sheep, and pigs have been experimented upon.

In the case of oxen, the food and litter (sometimes with an acid absorbent), were weighed, sampled, and analysed; the

animals were fed in boxes, for periods of from five to nine weeks, and the total dung produced was well mixed, weighed, sampled, and analysed. The constituents determined in the food and litter on the one hand, and in the dung on the other, were dry matter, ash, and nitrogen.

In the case of sheep no litter was used; the animals were kept in lots of five, on rafters, through which (but with some little loss) the solid and liquid excreta passed on to a sheet-zinc flooring at such an incline that the liquid drained off at once into carboys containing acid, and the solid matter was removed two or three times daily, and also mixed with acid. The constituents determined in the food and manure were dry matter, mineral matter, sometimes woody fibre, and nitrogen.

In the case of pigs, individual male animals were experimented upon, each for periods of three, five, or ten days only. Each animal was kept in a frame, preventing it from turning round, and having a zinc bottom, with an outlet for the liquid to run into a bottle, and it was watched night and day, and the voidings carefully collected as soon as passed, which could easily be done, as the animal never passed either fæces or urine without getting up, and in getting up he rang a bell, and so attracted the notice of the attendant. The constituents determined were, in the food and fæces, dry matter, ash and nitrogen, and in the urine, dry matter, ash, nitrogen, and urea.

The loss or expenditure of constituents, by respiration and the cutaneous exhalations, has not been determined directly, that is by means of a respiration-apparatus, but only by difference, that is, by calculation, founded on the amounts of dry matter, ash, and nitrogen, in the food, and in the (increase) fæces, and urine.

Independently of the points of inquiry above enumerated, the results obtained have supplied data for the consideration of the following questions:—

1. The characteristic demands of the animal body (for nitrogenous or non-nitrogenous constituents of food) in the exercise of muscular power.
2. The sources in the food of the fat produced in the animal body.
3. The comparative characters of animal and vegetable food in human dietaries.

SUPPLEMENTARY INVESTIGATIONS.

In conjunction with Professor Way, an extensive investigation was undertaken on the application of town sewage to different crops, but especially to grass. The amount, and the composition, of both the sewage and the produce grown were determined; and, in selected cases, the composition of the land drainage-water was also determined. Comparative experiments were also made on the feeding qualities of the differently grown produce; the amount of increase yielded by oxen, and the amount and composition of the milk yielded by cows, being determined. In this inquiry part of the analytical work was performed at Rothamsted, but most of it by Professor Way in London.

The chemistry of the malting process, the loss of food constituents during its progress, and the comparative feeding value of barley and malt, have been investigated.

Although many of the results of the investigations above enumerated have already been published, a large proportion as yet remains unpublished.

The following lists give the titles of the papers already published, arranged in two series, and within each series arranged in chronological order; and they show in what Journal each paper appeared.

SERIES I.—REPORTS OF FIELD EXPERIMENTS, EXPERIMENTS ON VEGETATION, &c.
PUBLISHED 1847—1881, INCLUSIVE.

- | | |
|--|---|
| <p>1. Agricultural Chemistry (Jour. Roy. Ag. Soc. Eng., vol. viii., p. 226) 1847</p> <p>2. Agricultural Chemistry, Turnip Culture (Jour. Roy. Ag. Soc. Eng., vol. viii., p. 494) 1847</p> <p>3. Experimental Investigation into the Amount of Water Given Off by Plants during their Growth, especially in relation to the Fixation and Source of their various Constituents (Jour. Hort. Soc. Lond., vol. v., p. 38) 1850</p> <p>4. Report of some Experiments undertaken at the suggestion of Professor Lindley, to ascertain the Comparative Evaporating Properties of Evergreen and Deciduous Trees (Jour. Hort. Soc. Lond., vol. vi., p. 227) 1851</p> <p>5. Agricultural Chemistry, especially in relation to the Mineral Theory of Baron Liebig (Jour. Roy. Ag. Soc. Eng., vol. xii., p. 1) 1851</p> <p>6. On the Amounts of, and Methods of Estimating, Ammonia and Nitric Acid in Rain-water (Report of the British Association for the Advancement of Science for 1854) 1854</p> <p>7. Report of the Right Hon. the Earl of Leicester, on the Experiments, conducted by Mr. Keary, on the Growth of Wheat upon the same land for four successive years, at Holkham Park Farm (Jour. Roy. Ag. Soc. Eng., vol. xvi., p. 207) 1855</p> <p>8. On some points connected with Agricultural Chemistry; being a reply to Baron Liebig's "Principles of Agricultural Chemistry" (Jour. Roy. Ag. Soc. Eng., vol. xvi., p. 411) 1855</p> <p>9. On the Growth of Wheat by the Lois Weedon System, on the Rothamsted Soil; and on the Combined Nitrogen in Soils (Jour. Roy. Ag. Soc. Eng., vol. xvii., p. 582) 1856</p> <p>10. On some points in the Composition of Wheat Grain, its Products in the Mill, and Bread (Journal of the Chemical Society of London, vol. x., p. 1) .. 1857</p> <p>11. On the Growth of Barley by Different Manures continuously on the Same Land; and on the Position of the Crop in Rotation (Jour. Roy. Ag. Soc. Eng., vol. xviii., p. 454) 1857</p> <p>12. Report of Experiments with different Manures on Permanent Meadow Land, with Tabular Appendix (Jour. Roy. Ag. Soc. Eng., vols. xix., p. 552, and xx., pp. 228 and 398) 1858-9</p> <p>13. Report of Experiments on the Growth of Red Clover by different Manures (Jour. Roy. Ag. Soc. Eng., vol. xxi., p. 178) 1860</p> <p>14. On the Sources of the Nitrogen of Vegetation; with special reference to the question whether Plants Assimilate Free or Uncombined Nitrogen.—Abstract (Proceedings of the Royal Society of London, vol. x., p. 544) 1860</p> <p>15. On the Application of Different Manures to Different Crops, and on their Proper Distribution on the Farm 1861</p> | <p>16. On some Points in connection with the Exhaustion of Soils.—Abstract (Report of the British Association for the Advancement of Science for 1861) .. 1861</p> <p>17. On the Sources of the Nitrogen of Vegetation, with special reference to the question whether Plants Assimilate Free or Uncombined Nitrogen (Philosophical Transactions, part 2, 1861) 1861</p> <p>18. Report of Experiments made at Rodmersham, Kent, on the Growth of Wheat by different Descriptions of Manure for several years in succession on the same land (Jour. Roy. Ag. Soc. Eng., vol. xxiii., p. 31) .. 1862</p> <p>19. The Effects of Different Manures on the Mixed Herbage of Grass Land (Jour. Roy. Ag. Soc. Eng., vol. xxiv., p. 131) 1863</p> <p>20. On the Sources of the Nitrogen of Vegetation, with special reference to the question whether Plants assimilate Free or Uncombined Nitrogen (Jour. Chem. Soc., new series, vol. i.; entire series, vol. xvi.) 1863</p> <p>21. Liebig and the "Mineral Theory" (note, extracted from a paper by Messrs. Lawes and Gilbert, Jour. Roy. Ag. Soc. Eng., vol. xxiv., part 2) 1863</p> <p>22. Further Report of Experiments with Different Manures on Permanent Meadow Land (Jour. Roy. Ag. Soc. Eng., vol. xxiv., part 2) 1863</p> <p>23. Report of Experiments on the Growth of Wheat for Twenty Years in Succession on the same land (Jour. Roy. Ag. Soc. Eng., vol. xxv., parts 1 and 2) 1864</p> <p>24. On the Selection of Artificial Manures for the Sugar-cane 1864</p> <p>25. On the Accumulation of the Nitrogen of Manure in the Soil (Report of the British Association for the Advancement of Science for 1866) 1866</p> <p>26. Preliminary Notice of Results on the Composition of Wheat grown for twenty years in succession on the same land (Report of the British Association for the Advancement of Science for 1867) 1867</p> <p>27. On the Home Produce, Imports, and Consumption of Wheat (Jour. Roy. Ag. Soc. Eng., vol. vi., s.s., part 2) 1868</p> <p>28. Exhaustion of the Soil in relation to Landlords' Covenants, and the Valuation of Unexhausted Improvements (read before the London Farmers' Club, April 4, 1870) 1870</p> <p>29. Scientific Agriculture with a view to Profit (read before the Maidstone Farmers' Club, Dec. 15, 1870) 1870</p> <p>30. Reports of Experiments on the Influence of various Manures on different Species of Plants (Proceedings of the Royal Horticultural Society) 1870</p> <p>31. Effects of the Drought of 1870 on some of the Experimental Crops at Rothamsted (Jour. Roy. Ag. Soc. Eng., vol. vii., s.s., part 1) 1871</p> <p>32. Notes on Clover Sickness (Jour. Roy. Hort. Soc., vol. iii.) 1871</p> <p>33. Report of Experiments on the Growth of Barley for Twenty Years in Succession on the same land (Jour. Roy. Ag. Soc. Eng., vol. ix., s.s., parts 1 and 2) 1873</p> |
|--|---|

34. Unexhausted Tillages and Manures, with reference to the Landlord and Tenant (Ireland) Act, 1870 .. 1874
35. On the more frequent Growth of Barley on Heavy Land (read before the London Farmers' Club, February 1, 1875) 1875
36. On the Valuation of Unexhausted Manures (Jour. Roy. Ag. Soc. Eng., vol. xi., s.s., part 1) 1875
37. Note on the Occurrence of "Fairy Rings" (Jour. Linn. Soc., Botany, vol. xv., p. 17) 1875
38. On some points in connection with Vegetation (Address delivered at South Kensington in the Chemical Section of the Science Conferences) 1876
39. On Rainfall, Evaporation, and Percolation (Proceedings of the Inst. of Civil Engineers, vol. xiv., part 3) 1876
40. Freedom in the Growth and Sale of the Crops of the Farm, considered in relation to the interests of the Landowner and the Tenant Farmer (Jour. Soc. Arts, December 14, 1877) 1877
41. On Nitrification; Part I., a Report of Experiments made in the Rothamsted Laboratory (Jour. Chem. Soc., January, 1878) 1878
42. Composition of Potatoes (Note—Jour. Roy. Hort. Soc., vol. v., part 5; Proceedings, p. xxxvii.) 1878
43. Is Higher Farming a remedy for Lower Prices? (Lecture delivered before the East Berwickshire Agricultural Association, May 3, 1879. Published by G. Macaskie, 'Warder' Office, Berwick) .. 1879
44. On Nitrification; Part II., a Report of Experiments made in the Rothamsted Laboratory (Jour. Chem. Soc., July, 1879) 1879
45. On the Determination of Nitric Acid as Nitric Oxide, by means of its action on Mercury; a Report of Experiments made in the Rothamsted Laboratory (Jour. Chem. Soc., July, 1879) 1879
46. On the Determination of Nitric Acid by means of Indigo, with special reference to Water Analysis; a Report of Experiments made in the Rothamsted Laboratory (Jour. Chem. Soc., September, 1879) .. 1879
47. Agricultural, Botanical, and Chemical Results of Experiments on the Mixed Herbage of Permanent Meadow, conducted for more than twenty years in succession on the same Land.—Abstract (Proceedings of the Royal Society, No. 197, 1879) 1879
48. On some points in connection with Agricultural Chemistry.—Abstract (Report of the British Association for the Advancement of Science for 1879) .. 1879
49. Our Climate and our Wheat-Crops (Jour. Roy. Ag. Soc. Eng., vol. xvi., s.s., part 1) 1880
50. On the Home Produce, Imports, Consumption, and Price of Wheat, over twenty-eight (or twenty-seven) harvest-years, 1852-53 to 1879-80 inclusive (Jour. of the Statistical Society, June, 1880) 1880
51. Agricultural, Botanical, and Chemical Results of Experiments on the Mixed Herbage of Permanent Meadow, conducted for more than twenty years in succession on the same Land.—Part I. The Agricultural Results (Philosophical Transactions, part 1, 1880) 1880
52. Sketch of the Progress of Agricultural Chemistry: Address to the Chemical Section of the British Association (Report of the British Association for the Advancement of Science for 1880) 1880
53. On the determination of Nitric Acid as Nitric Oxide by means of its reaction with Ferrous Salts.—Part I. A Report of Experiments made in the Rothamsted Laboratory (Jour. Chem. Soc., July, 1880) 1880
54. On the determination of Carbon in Soils; a Report of Experiments made in the Rothamsted Laboratory (Jour. Chem. Soc., September, 1880) 1880
55. On the Home Produce, Imports, Consumption, and Price of Wheat, over twenty-seven (or twenty-eight) harvest-years, 1852-3 to 1879-80 (Jour. Roy. Ag. Soc., vol. xvi., s.s., part 2, 1880) 1880
56. Letter on "Bread Reform" (Journal of the Society of Arts, January 21, 1881) 1881
57. On the Amount and Composition of the Rain and Drainage-Waters collected at Rothamsted; Parts I. and II. (Jour. Roy. Ag. Soc., Eng., vol. xvii., s.s., part 1, 1881) 1881
58. Letters on "Fertility" (Agricultural Gazette, Feb. 21 and 28; March 7, 14, and 21; April 4, 11, 18, and 25; May 2 and 9, 1881) 1881

SERIES II.—REPORTS OF EXPERIMENTS ON THE FEEDING OF ANIMALS, SEWAGE UTILISATION, &c.
PUBLISHED 1849—1877, INCLUSIVE.

- | | | | |
|---|------|---|------|
| 1. Agricultural Chemistry : Sheep Feeding and Manure, Part I. (With Tabular Appendix in 1856.) (Jour. Roy. Ag. Soc. Eng., vol. x., p. 276) | 1849 | Food, and on its relations to Bread—Abstract (Jour. Chem. Soc., vol. xii., p. 54) | 1860 |
| 2. Report of Experiments on the Comparative Fattening Qualities of Different Breeds of Sheep; Hampshire and Sussex Downs (Jour. Roy. Ag. Soc. Eng., vol. xii., p. 414) | 1851 | 15. Fifth Report of Experiments on the Feeding of Sheep (Jour. Roy. Ag. Soc. Eng., vol. xxii., p. 189) | 1861 |
| 3. Report of Experiments on the Comparative Fattening Qualities of Different Breeds of Sheep—Cotswolds (Jour. Roy. Ag. Soc. Eng., vol. xiii., p. 179) | 1852 | 16. Report of Experiments on the Fattening of Oxen at Woburn Park Farm (Jour. Roy. Ag. Soc. Eng., vol. xxii., p. 200) | 1861 |
| 4. On the Composition of Foods in relation to Respiration and the Feeding of Animals (Report of the British Association for the Advancement of Science for 1852) | 1852 | 17. Experiments on the Question whether the Use of Condiments increases the Assimilation of Food by Fattening Animals, or adds to the Profits of the Feeder (Edinburgh Veterinary Review and Annals of Comparative Pathology, July, 1862) | 1862 |
| 5. Agricultural Chemistry : Pig Feeding (Jour. Roy. Ag. Soc. Eng., vol. xiv., p. 459) | 1853 | 18. Supplementary Report of Experiments on the Feeding of Sheep (Jour. Roy. Ag. Soc. Eng., vol. xxiii., p. 191) | 1862 |
| 6. On the Equivalency of Starch and Sugar in Food (Report of the British Association for the Advancement of Science for 1854) | 1854 | 19. The Utilisation of Town Sewage (Jour. Roy. Ag. Soc. Eng., vol. xxiv., p. 65) | 1863 |
| 7. Experiments on the Comparative Fattening Qualities of Different Breeds of Sheep—Leicesters and Cross-breeds (Jour. Roy. Ag. Soc. Eng., vol. xvi., p. 45) | 1855 | 20. On the Chemistry of the Feeding of Animals for the Production of Meat and Manure (read before the Royal Dublin Society, March 31, 1864) | 1864 |
| 8. On the Sewage of London (Journal of the Society of Arts, March 7, 1855) | 1855 | 21. On the Sewage of Towns (Third Report and Appendices 1, 2, and 3, of the Royal Commission. Presented to Parliament) | 1865 |
| 9. Letter on the Utilisation of Town Sewage (from the Report ordered by the House of Commons to be printed, Aug. 3, 1857. Appendix xii., p. 477) | 1857 | 22. Report (presented to Parliament) of Experiments undertaken by Order of the Board of Trade to Determine the Relative Values of Unmalted and Malted Barley as Food for Stock | 1866 |
| 10. Experimental Inquiry into the Composition of some of the Animals Fed and Slaughtered as Human Food. Abstract (Proceedings of the Royal Society of London, vol. ix., p. 348) | 1858 | 23. On the Composition, Value, and Utilisation of Town Sewage (Jour. Chem. Soc., New Series, vol. iv.; Entire Series, vol. xix.) | 1866 |
| 11. Observations on the recently-introduced Manufactured Foods for Agricultural Stock (Jour. Roy. Ag. Soc. Eng., vol. xix., p. 199) | 1858 | 24. Food, in its Relations to the various Exigencies of the Animal Body (Phil. Mag., July, 1866) | 1866 |
| 12. Experimental Inquiry into the Composition of some of the Animals Fed and Slaughtered as Human Food (Philosophical Transactions, Part 2, 1859) | 1859 | 25. On the Sources of the Fat of the Animal Body (Phil. Mag., December, 1866) | 1866 |
| 13. On the Composition of Oxen, Sheep, and Pigs, and of their Increase while Fattening (Jour. Roy. Ag. Soc. Eng., vol. xxi., p. 433) | 1860 | 26. Note—On Sewage Utilisation (Proceedings of the Institution of Civil Engineers, vol. xiv., Part 3) | 1876 |
| 14. On the Composition of the Animal Portion of our | | 27. On some Points in connection with Animal Nutrition (Address delivered at South Kensington in the Biological Section of the Science Conferences) | 1876 |
| | | 28. On the Formation of Fat in the Animal Body (Journal of Anatomy and Physiology, vol. xi., Part 4) | 1877 |

THE PARK.
EXPERIMENTS WITH DIFFERENT MANURES ON PERMANENT MEADOW LAND.

The Land has probably been laid down with Grass for some centuries. No fresh seed has been artificially sown within the last 40 years certainly; nor is there record of any having been sown since the Grass was first laid down. The experiments commenced in 1856, at which time the character of the herbage appeared uniform over all the Plots. Excepting as explained in the Table, and in the foot-notes, the same description of Manure has been applied year after year to the same Plot.

During the first 19 years of the experiments, 1856-1874, the first crop only, each year, was mown, made into hay, removed from the land, and weighed. As a rule, the second crop was fed-off by sheep having no other food, the object being not to disturb the condition of the manuring. A given number was allotted to each Plot, according to the amount of produce, penned upon a portion of it, and the area extended, day by day, until the whole was eaten down. Frequently, however, the animals suffered considerably; and in 1866, 1870, 1873, and 1874, the second crops (and third, if any) were cut, and spread on the respective Plots. In the twentieth season, 1875, the second crops being unusually heavy, and the weather favourable, they were, for the first time, cut, weighed as hay, and removed. In 1877 and 1878 the second crops were again made into hay, weighed, and removed. In 1879 the second crops were cut, sampled, carted, and weighed, green; the dry matter in the weighed samples was determined, and the produce reckoned into hay by adding one-fourth to the calculated dry matter per acre. In 1880 the second crops were again made into hay, weighed and removed. It is intended in future to adopt this plan whenever the weather will permit. Owing to this change in the treatment of the crops, the average produce per annum is given, separately, for the first 20 years, 1856-1875, first crops only, and for the succeeding 5 years, 1876-1880, first and second crops (18).

(Area under experiments, about 7 acres.)

PLOTS.	Manures, per acre, per Annum.	PRODUCE PER ACRE, WEIGHED AS HAY.										PLOTS.	
		Average per Annum, 20 Years, 1856-75. (First Crops only.)		Average per Annum, 5 Years, 1876-80. (First and Second Crops.)		Average per Annum, 20 Years, 1856-75. (First Crops only.)		Average per Annum, 5 Years, 1876-80. (First and Second Crops.)		Twenty-fifth Season, 1880.		Total.	
		Cwts.	Cwts.	Cwts.	Cwts.	Cwts.	Cwts.	Cwts.	Cwts.	Cwts.	Cwts.	Cwts.	Cwts.
1	(1856-63, 8 years, 14 tons Farmyard Manure, and 200 lbs. Ammonia-salts (1); average produce 49½ cwts. (1))	48½	37½	32½	367	23½	167	12½	29½	15	7½	18½	1
2	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	41½	32	23½	367	23½	167	12½	29½	15	7½	18½	2
3	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	22½	20	21½	22½	21½	22½	21½	22½	9	9	18	3
4	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	33½	30½	32½	30½	32½	30½	32½	30½	24½	24½	48½	4
5	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	30½	22	19½	14½	14½	14½	14½	14½	8½	8½	17	5
6	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	31½	30½	26½	26½	26½	26½	26½	26½	22½	22½	45	6
7	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	33½	36½	35½	35½	35½	35½	35½	35½	22½	22½	45	7
8	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	33½	26½	26½	26½	26½	26½	26½	26½	16½	16½	33	8
9	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	53½	46½	51	51	51	51	51	51	40½	40½	92	9
10	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	52½	39½	46½	46½	46½	46½	46½	46½	25½	25½	71	10
11	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	61½	57½	57½	57½	57½	57½	57½	57½	30	30	91	11
12	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	63½	61½	61½	61½	61½	61½	61½	61½	44½	44½	108	12
13	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	55½	59½	57½	57½	57½	57½	57½	57½	51½	51½	103	13
14	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	53½	60½	57½	57½	57½	57½	57½	57½	51½	51½	103	14
15	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	36½	35	33½	33½	33½	33½	33½	33½	19	19	54	15
16	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	45½	47½	46½	46½	46½	46½	46½	46½	37½	37½	84	16
17	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	34½	33½	33½	33½	33½	33½	33½	33½	21½	21½	53	17
18	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	21	33½	33½	33½	33½	33½	33½	33½	15	15	30	18
19	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	16½	16½	33	19
20	(1856-63, 8 years, 14 tons Farmyard Manure; average produce 42½ cwts. (1))	15	15	30	20

(1) "Ammonia-salts"—in all cases equal parts Sulphate and Muriate of Ammonia of Commerce.
 (2) The "Superphosphate of Lime" is, in all cases, made from 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid Sp. Gr. 1.7 (and water).
 (3) Plots 6, 8, and 10, had, besides the Manures specified, 2000 lbs. Sawdust per acre per annum for the first 7 years, 1856-1862, but without effect.
 (4) 200 lbs. 1856-63 inclusive.
 (5) 500 lbs. in 1862 and 1863. (6) Only 400 lbs. in 1856-60-61.
 (7) The application of Silicates did not commence until 1862; 8 years (1862-1870), 200 lbs. Silicate Lime, and 200 lbs. Silicate Soda; 1871, and since, 400 lbs. Silicate Soda.
 (8) The manures specified were first applied in 1859 (previously, 1856-7 and 8, Sawdust only).
 (9) Averages of 8 years, 10 years, and 18 years, as these experiments did not commence until 1868.
 (10) Averages of 4 years, 10 years, and 11 years, as the experiment only commenced in 1866.
 (11) Averages of 4 years only, 1872-75.
 (12) As the second crops were not removed in 1876, those of 1875, which were, are brought in to give the 5 years, average of second crops.

HOOS FIELD.
EXPERIMENTS ON THE GROWTH OF BARLEY YEAR AFTER YEAR ON THE SAME LAND, WITHOUT MANURE, AND WITH DIFFERENT KINDS OF MANURE.

Previous Cropping—1847, Swedish Turnips, with Dung and Superphosphate of Lime, the Roots carted off; 1848, Clover; 1850, Wheat; 1851, Barley manured with Ammonia-salts. First Experimental Barley Crop in 1852. Barley every year since. The crop of the present year, 1881, is, therefore, the 30th Barley crop in succession. Unless stated to the contrary in the Table, or in the foot-notes, the same Manure has been applied year after year to the same Plot.

(Area under experiment, about 4½ acres.)

PLOTS.	Description of Manures applied	PRODUCTS PER ACRE.																		PLOTS.				
		Average per Annum.									Total Straw.										Twenty-ninth Season, 1880.			
		Quantity.			Weight per Bushel.			Dressed Corn.			14 Years, 1862-79.			14 Years, 1867-79.			28 Years, 1862-79.				Dressed Corn.	Total Straw.		
		Bushels.		Cwts.	Dressed Corn.		Cwts.	Dressed Corn.		Cwts.	Dressed Corn.		Cwts.	Dressed Corn.	Cwts.	Bushels.	Cwts.							
		14 Years, 1860-79.	23 Years, 1862-79.		14 Years, 1862-79.	14 Years, 1867-79.		28 Years, 1862-79.	14 Years, 1867-79.		28 Years, 1862-79.	14 Years, 1867-79.						28 Years, 1862-79.						
1 O.	Unmanured continuously	1 O.
2 O.	3½ cwts. Superphosphate of Lime (1)	2 O.
3 O.	275 lbs. Sulphate of Potass, 100 lbs. (2) Sulphate Soda, 100 lbs. (3) Sulphate Magnesia	3 O.
4 O.	200 lbs. (4) Sulphate Potass, 100 lbs. (5) Sulphate Soda, 100 lbs. (6) Superphosphate	4 O.
1 A.	200 lbs. Ammonia-salts (7)	1 A.
2 A.	200 lbs. Ammonia-salts and 3½ cwts. Superphosphate	2 A.
3 A.	200 lbs. Ammonia-salts, 200 lbs. (8) Sulph. Potass, 100 lbs. (9) Sulph. Soda, 100 lbs. Magnesia	3 A.
4 A.	200 lbs. Ammonia-salts, 200 lbs. (10) Sulph. Potass, 100 lbs. (11) Sulph. Soda, 100 lbs. Mag., 3½ cwts. Superphos.	4 A.
1 A.A.	275 lbs. Nitrate Soda	1 A.A.
2 A.A.	275 lbs. Nitrate Soda and 3½ cwts. Superphosphate	2 A.A.
3 A.A.	275 lbs. Nitrate Soda, 200 lbs. (12) Sulph. Potass, 100 lbs. (13) Sulph. Soda, 100 lbs. Magnesia	3 A.A.
4 A.A.	275 lbs. Nitrate Soda, 200 lbs. (14) Sulph. Potass, 100 lbs. (15) Sulph. Soda, 100 lbs. Mag., 3½ cwts. Superphos.	4 A.A.
1 A.A.S.	275 lbs. Nitrate Soda, 400 lbs. Silicate Soda (16)	1 A.A.S.
2 A.A.S.	275 lbs. Nitrate Soda, 400 lbs. Silicate Soda, and 3½ cwts. Superphosphate (17)	2 A.A.S.
3 A.A.S.	275 lbs. Nitrate Soda, 400 lbs. Silicate Soda, 200 lbs. (18) Sulph. Potass, 100 lbs. (19) Sulph. Soda, 100 lbs. Sulph. Mag.	3 A.A.S.
4 A.A.S.	275 lbs. Nitrate Soda, 400 lbs. Silicate Soda, 200 lbs. (20) Sulph. Potass, 100 lbs. (21) Sulph. Soda, 100 lbs. Sulph. Mag., and 3½ cwts. Superphosphate	4 A.A.S.
1 C.	1000 lbs. Rape-cake and 3½ cwts. Superphosphate	1 C.
2 C.	1000 lbs. Rape-cake	2 C.
3 C.	1000 lbs. Rape-cake, 200 lbs. (22) Sulph. Potass, 100 lbs. (23) Sulph. Soda, 100 lbs. Sulph. Mag.	3 C.
4 C.	1000 lbs. Rape-cake, 200 lbs. (24) Sulph. Potass, 100 lbs. (25) Sulph. Soda, 100 lbs. Sulph. Mag., 3½ cwts. Superphos.	4 C.
1 N.	275 lbs. Nitrate of Soda	1 N.
2 N.	275 lbs. (26) Nitrate of Soda	2 N.
5 O.	200 lbs. (27) Sulphate of Potass, 3½ cwts. Superphosphate (28)	5 O.
5 A.	200 lbs. (29) Sulphate of Potass, 3½ cwts. Superphosphate, and 200 lbs. (30) Ammonia-salts	5 A.
M.	100 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, and 3½ cwts. Superphosphate	M.
6 (1)	Unmanured continuously	6 (1)
7 (2)	Ashes (burnt soil and turf)	7 (2)

excepting the addition of the Silicates, have been and are, in other respects, manured in the same way as the "A.A." plots.

(1) 2000 lbs. Rape-cake per annum for the first six years, and 1000 lbs. only, each year since.

(2) 500 lbs. Sulphate of Potass, and 3½ cwts. Superphosphate of Lime, without Nitrate of Soda, the first year (1852). Nitrate of Soda applied each year from 1852-4-5-6, and 7; and 275 lbs. only, each year since.

(3) 450 lbs. Nitrate of Soda, and 100 lbs. Sulphate of Potass, in 1852-4-5-6, and 7; and 275 lbs. only, each year since.

(4) Ammonia-salts also the first year, but not since. (1) By mistake 400 lbs. in 1880.

(5) Averages of 2 years, 14 years, and 16 years. (2) Averages of 13 years, 14 years, and 27 years.

(6) Averages of 8 years, 13 years, and 21 years. (3) Averages of 13 years, 14 years, and 27 years.

(7) Averages of 20 years (with dung), 8 years (unmanured), and 28 years.

(8) Weighed, owing to the foison of the plot, from the wet season.

(9) A.A.S. = Ammonia-salts, Superphosphate, and Silicate of Soda.

BROADBALK FIELD.

EXPERIMENTS ON THE GROWTH OF WHEAT YEAR AFTER YEAR ON THE SAME LAND, WITHOUT MANURE, AND WITH DIFFERENT KINDS OF MANURE. Previous Cropping—1839, Turnips, with Farmyard Manure; 1840, Barley; 1841, Peas; 1842, Wheat; 1843, Oats; the last four Crops Unmanured. First Experimental Wheat Crop in 1844. Wheat every year since; and, with some exceptions, nearly the same description of Manure on the same Plots each year—especially during the last 30 years (1852 and since). The Crop of the present year, 1881, is, therefore, the 38th Wheat Crop in succession. From the commencement of the experiments in 1843-4 up to 1876-7 inclusive, the mineral manures, the ammonia-salts, and rape-cake, &c., if any, were sown in the autumn, before the seed; in 1845, when, owing to the wet autumn and winter, all the manures were spring-sown; and for the crops of 1873, 4, 5, 6, and 7, the ammonia-salts applied to Plot 15 were top-dressed in the spring. Nitrate of soda has, however, always been sown in the autumn. But, in consequence of the ascertained great loss of the nitrogen of the manures by drainage, especially in wet winters, it has been decided to apply only the mineral manures (and Farmyard-manure) in the autumn, and the ammonia-salts, as well as the nitrate, in the spring; excepting on Plot 19, where, for comparison, the ammonia-salts are sown in the autumn. This plan was adopted for the first time for the crop of 1878.

(Area under experiment, about 13 acres.)

PLOTS.	PRODUCE PER ACRE.												Plots.
	Average per Annum.						Thirty-Seventh Season, 1880.						
	Dressed Corn.			Total Straw.			Dressed Corn.			Total Straw.			
	Quantity.	Weight per Bushel.	Quantity.	Weight per Bushel.	Quantity.	Weight per Bushel.	Quantity.	Weight per Bushel.	Quantity.	Weight per Bushel.	Quantity.	Weight per Bushel.	
0	18	58	14	58	15	58	11	58	16	58	11	58	0
1	16	58	10	58	15	58	9	58	11	58	12	58	1
2	35	58	31	58	30	58	30	58	34	58	32	58	2
3	15	58	11	58	14	58	9	58	11	58	11	58	3
4	16	58	11	58	14	58	9	58	12	58	11	58	4
5 (a and b)	28	58	19	58	16	58	10	58	17	58	15	58	5 (a and b)
6 (a and b)	37	58	28	58	26	58	18	58	23	58	26	58	6 (a and b)
7 (a and b)	39	58	33	58	31	58	29	58	34	58	33	58	7 (a and b)
8 (a and b)	36	58	31	58	29	58	28	58	31	58	30	58	8 (a and b)
9 { a } { b }	26	58	21	58	21	58	17	58	23	58	21	58	9 { a } { b }
10 { a } { b }	23	58	18	58	19	58	17	58	19	58	18	58	10 { a } { b }
11 (a and b)	30	58	22	58	26	58	21	58	25	58	25	58	11 (a and b)
12 (a and b)	35	58	26	58	31	58	25	58	30	58	29	58	12 (a and b)
13 (a and b)	35	58	27	58	31	58	28	58	32	58	30	58	13 (a and b)
14 (a and b)	33	58	27	58	30	58	27	58	31	58	29	58	14 (a and b)
15 { a } { b }	35	58	28	58	31	58	27	58	31	58	30	58	15 { a } { b }
16 (a and b)	39	58	13	58	45	58	12	58	14	58	13	58	16 (a and b)
17 (a and b)	18	58	13	58	15	58	11	58	14	58	12	58	17 (a and b)
18 (a and b)	32	58	24	58	28	58	22	58	26	58	23	58	18 (a and b)
19	32	58	24	58	28	58	22	58	26	58	23	58	19
20	15	58	10	58	13	58	10	58	12	58	11	58	20
21	22	58	17	58	19	58	14	58	17	58	16	58	21
22	21	58	18	58	19	58	15	58	17	58	16	58	22

(1) 300 lbs. per annum for Crop of 1858, and previously.
 (2) 200 lbs. per annum for Crop of 1858, and previously.
 (3) "Superphosphate of Lime"—in all cases, excepting for Plot 19, made from 200 lbs. Bone-ash, 150 lbs. Sulphuric acid sp. gr. 1.7 (and water).
 (4) "Ammonia-salts," in all cases, equal parts Sulphate and Murate of Ammonia of Common Salt, 550 lbs. each, and 1852, 275 lbs. of each, 1853, 1854, 90, 475 lbs. Nitrate in 1852, 550 lbs. of Potash, Soda, or Magn. Sulphate, 1854, 90, 475 lbs. Nitrate in 1853, 550 lbs. of each year since, 550 lbs. Nitrate is reckoned to contain the same amount of Nitrogen as 400 lbs. "Ammonia-salts."
 (5) For 1858, and previously—1/2 time as much.
 (6) For 1872 and previously, made with Murate instead of Sulphuric Acid.
 (7) For 1872 and previously, 400 lbs. Sulphate Ammonia, sown in the Autumn; for 1873, 4, 5, 6, and 7, 400 lbs. Ammonia-salts, sown in the Spring; for 1878 and since, 400 lbs. Ammonia-salts, sown in the Autumn.
 (8) For 1872 and previously, 300 lbs. Sulphate Ammonia and 500 lbs. Rape-cake, sown in the Autumn; for 1873, 4, 5, 6, and 7, 400 lbs. Ammonia-salts, sown in the Spring; for 1878 and since, 400 lbs. Ammonia-salts, sown in the Autumn.
 (9) For 1872 and previously, 300 lbs. Sulphate Ammonia and 500 lbs. Rape-cake, sown in the Autumn; for 1873, 4, 5, 6, and 7, 400 lbs. Ammonia-salts, sown in the Spring; for 1878 and since, 400 lbs. Ammonia-salts, sown in the Autumn.
 (10) "a" portion of Plot 15. For the crop of 1880, and since, the return of the straw was been discontinued.
 (11) The Plots marked "a" and "b" are divided into duplicate portions, "a" and "b," respectively, which are manured alike; excepting that, for the crops of 1864-5-6 and 7, the "a" portions of plots 5, 6, 7, 8, 9, 16, and 17 (or 18), received a mixture of soluble Silicates in addition to the other Manures, but, hitherto, without any material effect; and for the crops of 1868 to 1879 inclusive, cut straw (that produced in the previous season) was applied (instead of Silicates) on the "a" portions of plots 5, 6, 7, 8, 11, 12, 13, 14, and 17 (or 18); also for the crop of 1874, and each succeeding crop to 1879 inclusive, the straw of the previous season was cut up and applied to the "a" portion of Plot 15. For the crop of 1880, and since, the return of the straw was been discontinued.
 (12) The Manures of Plots 17 and 18 are, year by year, transposed.
 (13) Made with Murate instead of Sulphuric Acid.
 (14) Averages of Mineral Manures, alternated with Ammonia-salts.
 (15) Averages of Ammonia-salts, alternated with Mineral Manures.
 (16) Averages of Ammonia-salts, alternated with Mineral Manures.
 (17) Averages of 14 years, 13 years, and 27 years only; as, in 1868, owing to a mistake in carting, the produce could not be ascertained.
 (18) The Plots marked "a" and "b" are divided into duplicate portions, "a" and "b," respectively, which are manured alike; excepting that, for the crops of 1864-5-6 and 7, the "a" portions of plots 5, 6, 7, 8, 9, 16, and 17 (or 18), received a mixture of soluble Silicates in addition to the other Manures, but, hitherto, without any material effect; and for the crops of 1868 to 1879 inclusive, cut straw (that produced in the previous season) was applied (instead of Silicates) on the "a" portions of plots 5, 6, 7, 8, 11, 12, 13, 14, and 17 (or 18); also for the crop of 1874, and each succeeding crop to 1879 inclusive, the straw of the previous season was cut up and applied to the "a" portion of Plot 15. For the crop of 1880, and since, the return of the straw was been discontinued.

GEESCHOPT FIELD.

EXPERIMENTS ON THE GROWTH OF OATS YEAR AFTER YEAR ON THE SAME LAND; WITHOUT MANURE, AND WITH DIFFERENT KINDS OF MANURE.

Previous Cropping—1847 and 1848, Clover, Experimental Manures; 1849—1859, Beans, Experimental Manures; 1860, Fallow; 1861 and 1862, Wheat, Unmanured; 1863, Fallow; 1864, Beans, Dunged; 1865, Wheat, Unmanured; 1866, Beans, Unmanured; 1867 and 1868, Wheat, Unmanured.

The first Experimental Oat Crop was in 1869; the last in 1878, since which time, owing to the wetness and the foulness of the land, it has been left fallow.

(Area under Experiment, $\frac{1}{2}$ acre.)

PLOTS.	MANURES, PER ACRE, PER ANNUM.	PRODUCE PER ACRE.											
		1ST SEASON, 1869.		2ND SEASON, 1870.		3RD SEASON, 1871.		4TH SEASON, 1872.		5TH SEASON, 1873.		AVERAGE PER ANNUM 5 YEARS, 1869-1873.	
		Dressed Corn.	Total Straw.	Dressed Corn.	Total Straw.	Dressed Corn.	Total Straw.	Dressed Corn.	Total Straw.	Dressed Corn.	Total Straw.	Dressed Corn.	Total Straw.
Bushels.	lbs. per Bushel.	Bushels.	lbs. per Bushel.	Bushels.	lbs. per Bushel.	Bushels.	lbs. per Bushel.	Bushels.	lbs. per Bushel.	Bushels.	lbs. per Bushel.	Bushels.	lbs. per Bushel.
1	Unmanured	36 $\frac{1}{2}$	86 $\frac{1}{2}$	16 $\frac{1}{2}$	35	20 $\frac{1}{2}$	33 $\frac{1}{2}$	15	36 $\frac{1}{2}$	10 $\frac{1}{2}$	27 $\frac{1}{2}$	19 $\frac{1}{2}$	53 $\frac{1}{2}$
2	{ 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 400 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwts. Superphosphate of Lime (1)	45	83 $\frac{1}{2}$	19 $\frac{1}{2}$	35 $\frac{1}{2}$	22	35 $\frac{1}{2}$	19 $\frac{1}{2}$	37 $\frac{1}{2}$	17	28 $\frac{1}{2}$	17	35
3	400 lbs. Ammonia-salts (2)	56 $\frac{1}{2}$	87 $\frac{1}{2}$	30	34 $\frac{1}{2}$	57 $\frac{1}{2}$	36 $\frac{1}{2}$	55 $\frac{1}{2}$	37 $\frac{1}{2}$	36 $\frac{1}{2}$	32 $\frac{1}{2}$	47	35 $\frac{1}{2}$
4	{ 400 lbs. Ammonia-salts, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwts. Superphosphate	75 $\frac{1}{2}$	89 $\frac{1}{2}$	50 $\frac{1}{2}$	36	58 $\frac{1}{2}$	35 $\frac{1}{2}$	62 $\frac{1}{2}$	39 $\frac{1}{2}$	48 $\frac{1}{2}$	34 $\frac{1}{2}$	59	37
5	550 lbs. Nitrate of Soda (3)	62 $\frac{1}{2}$	88 $\frac{1}{2}$	36 $\frac{1}{2}$	35 $\frac{1}{2}$	55	36 $\frac{1}{2}$	42 $\frac{1}{2}$	36 $\frac{1}{2}$	39 $\frac{1}{2}$	30 $\frac{1}{2}$	47 $\frac{1}{2}$	35 $\frac{1}{2}$
6	{ 550 lbs. Nitrate of Soda, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwts. Superphosphate	69 $\frac{1}{2}$	88 $\frac{1}{2}$	50	35 $\frac{1}{2}$	60 $\frac{1}{2}$	33 $\frac{1}{2}$	44 $\frac{1}{2}$	37 $\frac{1}{2}$	63 $\frac{1}{2}$	33 $\frac{1}{2}$	57 $\frac{1}{2}$	35 $\frac{1}{2}$
SECOND 5 YEARS; MINERAL MANURES AS BEFORE, AMMONIA-SALTS AND NITRATE OF SODA ONLY HALF AS MUCH AS PREVIOUSLY.													
PLOTS.	MANURES, PER ACRE, PER ANNUM.	6TH SEASON, 1874.		7TH SEASON, 1875.		8TH SEASON, 1876 (1).		9TH SEASON, 1877 (1) FALLOW.		10TH SEASON, 1878.		AVERAGE PER ANNUM 4 YEARS, 1874, 5, 6, and 8.	
		Bushels.	lbs. per Bushel.	Bushels.	lbs. per Bushel.	Bushels.	lbs. per Bushel.	Bushels.	lbs. per Bushel.	Bushels.	lbs. per Bushel.	Bushels.	lbs. per Bushel.
		Dressed Corn.	Total Straw.	Dressed Corn.	Total Straw.	Dressed Corn.	Total Straw.	Dressed Corn.	Total Straw.	Dressed Corn.	Total Straw.	Dressed Corn.	Total Straw.
1	Unmanured	12	31 $\frac{1}{2}$	12 $\frac{1}{2}$	29 $\frac{1}{2}$	8 $\frac{1}{2}$	32	22 $\frac{1}{2}$	32	13 $\frac{1}{2}$	31 $\frac{1}{2}$
2	{ 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwts. Superphosphate of Lime (1)	13 $\frac{1}{2}$	31 $\frac{1}{2}$	13 $\frac{1}{2}$	29 $\frac{1}{2}$	7 $\frac{1}{2}$	30	17 $\frac{1}{2}$	36 $\frac{1}{2}$	13 $\frac{1}{2}$	31 $\frac{1}{2}$
3	200 lbs. Ammonia-salts (2)	37 $\frac{1}{2}$	33 $\frac{1}{2}$	30 $\frac{1}{2}$	32 $\frac{1}{2}$	17 $\frac{1}{2}$	34 $\frac{1}{2}$	30	32 $\frac{1}{2}$	25 $\frac{1}{2}$	33 $\frac{1}{2}$
4	{ 200 lbs. Ammonia-salts, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwts. Superphosphate	46 $\frac{1}{2}$	34 $\frac{1}{2}$	30 $\frac{1}{2}$	34 $\frac{1}{2}$	29 $\frac{1}{2}$	35 $\frac{1}{2}$	45 $\frac{1}{2}$	37	38	35 $\frac{1}{2}$
5	275 lbs. Nitrate of Soda (3)	35 $\frac{1}{2}$	30 (4)	23 $\frac{1}{2}$	31 $\frac{1}{2}$	12 $\frac{1}{2}$	30 $\frac{1}{2}$	34 $\frac{1}{2}$	34 $\frac{1}{2}$	26 $\frac{1}{2}$	31 $\frac{1}{2}$
6	{ 275 lbs. Nitrate of Soda, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3 $\frac{1}{2}$ cwts. Superphosphate	23 $\frac{1}{2}$	33 $\frac{1}{2}$	28 $\frac{1}{2}$	33 $\frac{1}{2}$	19 $\frac{1}{2}$	33 $\frac{1}{2}$	37	36 $\frac{1}{2}$	25 $\frac{1}{2}$	34 $\frac{1}{2}$

(1) "Superphosphate of Lime"—in all cases, made from 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid sp. gr. 1.7 (and water).
 (2) "Ammonia-salts"—in each case, equal parts Sulphate and Muriate of Ammonia of Commerce.
 (3) 550 lbs. Nitrate of Soda is reckoned to contain the same amount of Nitrogen as 400 lbs. "Ammonia-salts."
 (4) On these plots, where large quantities of Nitrate of Soda had been applied year after year, the land, though more worked, was so wet that it could not be got into favourable condition for sowing, and the plant was very irregular. Owing to the extremely wet condition of the land, especially on the Nitrate plots, it was not sown until April 6, and then with a very unfavourable seed bed; and there being a heavy fall of snow a week later, the plant came up very irregularly, and much of it perished from standing surface-water.
 (5) Owing to the very wet winter, 1876-7, the land could not be worked in time for sowing, and was therefore left fallow in 1877; no manures being applied.

EXPERIMENTS ON THE GROWTH OF LEGUMINOUS CROPS.

I.—BEANS, PEAS, AND TARES—GEESCROFT FIELD.

EXPERIMENTS on the growth of Leguminous corn-crops (beans, peas, and tares), with different descriptions of manure, were commenced in 1847, about nine acres being devoted to the purpose.

Experiments with BEANS were continued without a break, for thirteen consecutive seasons, to 1859 inclusive; but, during the later years, the crop fell off very much, and the land became very foul.

In 1860 the land was fallowed.

In 1861 a crop of wheat, without manure, was taken.

In 1862 beans were again sown, but with some variation in the manuring.

In 1863 the land was fallowed.

In 1864, 5, 6, 7, 8, and 9, beans were grown, with much the same manures on the same plots, each year, as in 1862.

In the winter of 1869-70, 5000 lbs. of fresh burnt lime were applied per acre, over all the plots.

In 1870 beans were grown with the same manures on the respective plots as in 1864-69.

In October 1870 winter beans were sown (without manure), but the plants were to so great an extent destroyed by the severe weather which followed, that, in April 1871, the crop was ploughed up, and the land left fallow.

During the winter and early spring of 1871-2, the land was so wet that it could not be prepared in time for sowing. It was therefore left fallow for 1872; at the end of May it was subsoiled to a depth of about 12 inches, and re-ploughed in July.

The winter and early spring of 1872-3 were also so extremely wet, that it was again impossible to prepare the land in time for sowing; it was, however, ploughed up towards the end of March, again left fallow, and re-ploughed in July and October (1873).

On February 2, 1874, the land was again set with Beans, but without manure.

In 1875 Beans were re-sown, with the same manures on the respective plots as in 1864-1870; but owing to the wetness of the land in the first instance, and the subsequent hindrance by other spring sowing, they were not put in until April 1 and 2.

The wetness of the winter 1875-6, again prevented the preparation of the land in due time; and, though the manures were sown, and the land ploughed, it was left fallow during the summer of 1876.

Early in October, 1876, winter Beans were put in (drilled), without further manuring.

In 1878 the usual manures were sown, and beans were drilled on February 26.

Owing to the wetness of the winter, and the foul condition of the land, it was left fallow in 1879.

Owing to the continued wetness in the autumn, the severe winter, and foulness of the land, it could not be got into order for sowing, and remained fallow in 1880.

During 1880 the land was ploughed, scarified, and partially cleaned, but owing to the wetness of the autumn and the wetness and severity of the winter, it was again impossible to work the land in time for sowing; and it still remains fallow (1881).

The general result of the experiments with BEANS has been that mineral constituents used as manure (more particularly potass), increased the produce very much during the early years; and, to a certain extent, afterwards, whenever the season was favourable for the crop. Ammonia-salts, on the other hand, produced very little effect; notwithstanding that a Leguminous crop contains two, three, or more times as much nitrogen as a Gramineous one grown under similar conditions as to soil, &c. Nitrate of soda has, however, produced more marked effects. But Leguminous crops grown too frequently on the same land seem to be peculiarly subject to disease, which no conditions of manuring that we have hitherto tried seem to obviate.

Experiments with PEAS were soon abandoned, owing to the difficulty of keeping the land free from weeds, and an alternation

of BEANS and WHEAT was substituted; the Beans being manured much as in the experiments with the same crop grown continuously as above described. But the wetness of the winter of 1871-72 prevented the sowing of the Beans for the season of 1872; and again the wetness of the autumn and winter of 1872-3 prevented the sowing of the wheat until April 4, 1873, when Nursery wheat was put in, which, however, did not come to maturity, but was cut in the middle of September, yielding about 27 cwts. of gross produce per acre, containing too little corn to be worth thrashing. The land was ploughed in October 1873, and sown with beans February 3, 1874. On October 23, 1874, wheat was sown without manure. Beans should have been sown in 1876; indeed, the manures were sown, but, for the reason stated above, the land was left fallow; and wheat was put in October 24 (1876). In 1878 Beans were drilled, on February 26, with the usual manures. Owing to the wetness of the winter, and the condition of the land, it was left fallow in 1879; and it continues so up to the present time (May 1881).

In alternating WHEAT with BEANS, the remarkable result had been obtained, that nearly as much wheat, and nearly as much nitrogen, were yielded in eight crops of wheat in alternation with the highly nitrogenous beans, as in sixteen crops of wheat grown consecutively without manure in another field, and also nearly as much as were obtained in a third field in eight crops alternated with bare fallow.

Experiments with TARES, like those with Peas, were soon abandoned, and for the same reasons. Beans were at first substituted, with some variation in the description of the manures employed; but this experiment has likewise been abandoned for some years.

II.—RED CLOVER (*Trifolium pratense*).

1. Experiments on ordinary arable land.—HOOS FIELD.

EXPERIMENTS on the growth of Clover, on ordinary arable land, with many different descriptions of manure, were commenced in 1849, and, with the occasional interposition of a corn-crop, or fallow, were continued up to 1877, inclusive.

As with other Leguminous crops, the result was, that mineral constituents applied as manure (particularly potass) considerably increased the early crops. Ammonia-salts had little or no beneficial effect, and were sometimes injurious. It may be added, that the beneficial effects of long previous applications of potass have been apparent whenever there was any growth at all. To go a little more into detail:—

In the first year, 1849, the crops were throughout very heavy; especially with mineral, and without nitrogenous manure.

In autumn 1849 wheat was sown, and in spring 1850 Red Clover. In 1851 small cuttings were taken; and in 1852, though the crops were not heavy, there was by no means a failure.

Since that time, however, all attempts to grow clover year after year on this ordinary arable land have failed to give anything like a full crop, or even a plant which would stand the usual time on the ground.

Small cuttings were obtained in the autumns of 1855 and 1859, from seed sown in the spring of those years; and small but rather heavier cuttings in June and August 1865, from seed sown in 1864.

In April 1868 a portion only of the land was sown with Clover, and the plant for the most part died off in the winter.

In April 1869 the same portion was resown, and gave a small cutting in September of that year; but the plant again died off in the winter.

In April 1870 Clover was sown over the whole of the experimental land, this time in conjunction with Barley; but on those portions which had also been sown in 1868 and 1869 the plant again died off during the winter and early spring; whilst from those which had not been sown in 1868 and 1869 two small cuttings were taken in 1871.

EXPERIMENTS ON THE GROWTH OF LEGUMINOUS CROPS—*continued*.

In the spring of 1872, the plant being then almost entirely gone, the land was ploughed up. It was again ploughed in July 1872, and in March 1873; the intention being to sow some other Leguminous crop; but owing to the wetness and lateness of the season this was not done; the land was again left fallow, and re-ploughed in the beginning of June and the end of July (1873).

On May 4, 1874, the land was again ploughed, and sown with Red Clover seed, May 5, without manure. The plant came up well, and was very forward in September, when the flowering stems were cut down, but left on the land. During the winter and early spring the plant on those portions from which cuttings had been taken in 1871 almost entirely failed, and the land was ploughed up in May, and again in August (1875); whilst on those from which none had been taken since 1869 a fair plant remained, and two small cuttings were obtained, namely on June 23, and on August 9 and 12 (1875). On September 22, this portion of the land was ploughed up.

In May (1876) the whole was re-ploughed, again in July and September, and left fallow.

In May 1877, Barley and Clover were sown over the whole of the experimental Land, without further manuring, but the clover plant completely died off during the winter.

On two occasions (1851 and 1854) heavy dressings of Farnyard dung were applied to some of the plots; and in 1854 some received a dressing of 20 tons of dung, and 5000 lbs. of lime, per acre.

On some portions of the land Clover was sown 12 times during the 30 years, 1848–1877, and more frequently alone than with a corn-crop; but in 9 out of the last 10 trials the plant died off in the winter and spring succeeding the sowing the seed.

In 1878 the land was devoted to experiments with various Leguminous plants, differently manured, and these experiments are still in progress (1881); for further particulars see p. 16.

In reference to these field experiments on clover, it may be added that, in 1864, a portion of the land was trenched 2 feet deep, and one-third of the manure was mixed with the layer from 24 to 16 inches, one-third from 16 to 8 inches, and the remainder from 8 inches upwards. Owing to the characters of the season, the mechanical condition of the land was at first very unfavourable after this treatment; but, although many years have now elapsed, and the excess of constituents supplied was in some cases considerable, the plant has died off as completely on these plots as elsewhere.

Again, in the winter of 1867–8 a number of small beds, each 3 yards \times 2, were arranged on the previously unmanured plot of the experimental land. These were dug, some to the depth of 9 inches, some to the depth of 18, some to the depth of 27, and some to the depth of 36 inches, and sown to the respective depths with different mixtures; supplying in some cases very large amounts of potass, soda, lime, magnesia, phosphoric acid, sulphuric acid, nitrate of soda, &c.

From three similar sized beds, the soil was removed to the depths of 9, 18, and 27 inches respectively, and replaced by soil taken at the same depths from a garden border, on a portion of which Clover had been grown successfully since 1854, as described on the next page.

In April 1868 clover was sown on the whole of these small beds (as well as on some other portions of the experimental land); but the plant for the most part died off during the following winter.

In April 1869 the small beds (and the other portions as in 1868) were re-sown, small quantities of clover were cut in September of that year, but the plant again died off in the winter.

In April 1870 Clover was again sown on the small beds in conjunction with barley (as on all the rest of the experimental land), but the plant again died in the winter.

In the spring of 1871 the small beds were again re-sown, and the three with garden-soil were entirely enclosed, both around and above, by galvanised wire netting. Small cuttings were taken from these small beds in July 1872; and (excepting from the beds of garden-soil, which had yielded considerably more than the others in 1872), larger cuttings were taken in July 1873. The produce was the largest where potass and nitrate of soda were employed, and where they were applied in the largest quantity, and to the greatest depths.

In April 1874 there was still some healthy plant on all the beds, but it was considered to be too irregular to preserve. It was, therefore, dug in. The artificially-manured beds were re-manured as before, but only to the depth of 9 inches, and seed was sown on May 4th, July 6th, and October 22nd; each time the plant coming up well, but subsequently dying off. On the three beds of garden-soil, the plant from the first sowing (May 4), for the most part stood; requiring only to be made good here and there on July 6; and in September small cuttings were taken.

More small beds were arranged in the spring of 1874; on which the manures were dug in, at the various depths, on May 11th to 14th, and the seed sown on May 16th. At this time, the wire netting was removed from above the three beds of garden-soil, but the whole series of small beds was now surrounded with netting, to keep out ground game. One series of the new plots received sulphate of potass only, another nitrate of soda only, and a third the two together. The plants came up fairly well, but there were some blanks in the rows, which were re-sown on October 22 (1874). A cutting was taken on June 22 and 23 (1875) from these new beds; the blanks in the rows were re-sown on July 24; a second cutting was taken on August 17; and the blanks were again re-sown on September 22 (1875). The plant was the most even on the beds with sulphate of potass, less so on those with nitrate of soda, and less still on those with both together. The amount of produce was also greater with each of the manures used separately, than with the mixture of the two.

In May, 1875, the plant was entirely gone on the old artificially-manured beds, which were then dug up, and prepared for re-sowing. On the three beds of garden-soil, though the rows were imperfect, some healthy plants still remained, and gave a small cutting on June 22. On July 24 these beds were dug up; and they, as well as the artificially manured ones just referred to, were re-sown with seed. All came up well, but in May (1876), the plants on the beds of garden-soil were entirely gone, and those on the artificially manured ones nearly so, but they yielded small cuttings on July 17 (1876).

The plants on the new artificially manured beds, like those on the older ones, showed failure in the spring of 1876; but also, like them, gave small cuttings in July. All the small beds were dug up in August (1876); the artificially manured ones re-manured as in 1874, the manures dug in to a depth of 9 inches, and seed was sown on September 1, which came up, but the plants died off on all the plots in the winter of 1876–7.

In May 1877, all the small beds were dug up, and sown with Barley and Clover. To try the effects of shelter, the Barley stubble was left unusually high, but the young clover plants completely died off during the winter (1877–8).

In the spring of 1878 the beds were dug up, and cleaned; and they were re-sown with Clover, without further manuring, on June 12 and 13. All came up well, but the plant was almost entirely destroyed by "Fly."

In May 1879, there remained about a quarter of a plant on the plot with the largest amount of mineral manure, including potass, and sown to the greatest depth, and perhaps a third of a plant where the same mineral manures, with nitrate of soda in addition, had been applied; but there was scarcely a single plant on any of the other plots. On June 9 and 10, 1879, all the beds were cleaned, and re-sown with seed, which came up well; but a very wet and cold season following, most of the plants died off during the summer and autumn.

Early in June 1880, all the small beds were cleaned, and forked up; and on June 10, they were re-sown with seed without further manure. All came up well, but the plants were for the most part destroyed by the severe winter which followed.

In May, 1881, there was perhaps half a plant on two or three only out of the forty small beds; namely, where the mixed mineral manure, including potass, was used without nitrogenous manure; and the greatest vigour was where the manure was applied in the largest quantity, and to the greatest depths. On no other beds (not even on the three made up of garden-soil) was there nearly as much plant; and now (May 1881) all the small beds have been cleaned, the clover plants forked in, manures also forked in, as in 1876, to a depth of 8 or 9 inches, and clover seed sown.

It will be observed that, although in the earlier years, the three small beds in the field which had been artificially made up of surface-soil and subsoil brought from a highly manured

EXPERIMENTS WITH VARIOUS LEGUMINOUS PLANTS.—HOOS FIELD.

The land upon which attempts had been made to grow Red Clover in frequent succession since 1849, was devoted to experiments with various Leguminous Plants in 1878; so that the present season, 1881, is the fourth year of these experiments. The object was to ascertain whether, among a selection of plants all belonging to the Leguminous family, but of different habits of growth, and especially of different character and range of roots, some could be grown successfully for a longer time, and would yield more produce, containing more nitrogen as well as other constituents, than others; all being supplied with the same descriptions and quantities of manuring substances, applied to the surface soil. Further, whether the success in some cases, and the failure in others, would afford additional evidence as to the source of the nitrogen of the Leguminosae generally, and as to the causes of the failure of Red Clover in particular, when it is grown too frequently on the same land. Below, is given a list of the plants now growing (1881), all of which were sown on April 9; and, excepting in the few cases stated, the same description has been sown in each of the four years. The same mineral manure (if any) is applied to the same plot of each *Series*. *Series 1*, has mineral manure only; *Series 2*, the same mineral manure, and nitrates of soda; *Series 3*, the same mineral manure, and on one portion ammonia-salts, on another a dressing of cow's urine, and on another rape-cake, in addition. The mineral manures were applied in the quantities per acre stated in the Table, in 1878, and in 1880. The nitrates of soda, and the ammonia-salts, have been applied, in each of the four years, 1878, 1879, 1880, and 1881. The cows'-urine was applied in the second year only, 1879. The rape-cake was applied in the first and third years, 1878 and 1880; in the former on three lands, but in the latter on only two. It is estimated that already several times as much nitrogen has been removed in some of the other plants as in the Red Clover; but it is too soon to form any general conclusions from the results.

(Area under Experiment, about 3 acres, each plot, about $\frac{1}{4}$ th acre.)

Nos.		PLANTS GROWN ON EACH PLOT.		Notes.	
		Botanical Names.	Common Names.		
1		<i>Trifolium pratense</i>	Common Red, or Broad Clover.		
2		<i>Trifolium pratense</i> pervenne ..	Pervennial Clover, or Cow-grass.		
3		<i>Trifolium pratense</i> hybridum ..	Stations hybrid (Cow Clover).		
4		<i>Trifolium repens</i> ..	Common white, or Dutch Clover.		
5		<i>Trifolium repens</i> pervenne ..	Giant pervennial, White Clover.		
6		<i>Trifolium hybridum</i> ..	Alsike Clover.		
7		<i>Trifolium incarnatum</i> ..	Early Red, or Crimson Clover.		
8		<i>Trifolium hardiflorum album</i> ..	Late White Clover		
9		<i>Medicago lupulina</i> ..	Non-such, or Black Medick.		
10		<i>Melilotus alba</i> ..	White Clover, or Purple Medick.		
11		<i>Melilotus officinalis</i> ..	Balsam Clover.		
12		<i>Lotus corniculatus</i> ..	Common Bird's-foot Trefoil		
13		<i>Vicia sativa</i> ..	Summer Tare, or Vetch.		
14		<i>Onobrychis sativa</i> ..	Common Saintoin ..		
MANURES; QUANTITIES PER ACRE.					
Plots.		Series 1 (6 Lands). ¹		Series 2, ¹	Series 3 (5 Lands).
		Without Mineral Manure, or with Mineral Manure only.		Each Plot as Series 1, and—	Each Plot as Series 1, and—
1		Without Mineral Manure.		(5 Lands);	(2 Lands);
2		5 cwts. Superphosphate of Lime (°)		Each Plot as Series 1, and—	Each Plot as Series 1, and—
3		1000 lbs. Sulphate Potash			
4		1000 lbs. Sulphate Potash, 5 cwts. Superphosphate			
5		1000 lbs. Sulphate Potash, 250 lbs. Chloride Sodium, 250 lbs. Sulphate Lime, 250 lbs. Sulphate Magnesia			
6		1000 lbs. Sulphate Potash, 250 lbs. Chloride Sodium, 250 lbs. Sulphate Lime, 250 lbs. Sulphate Magnesia, 5 cwts. Superphosphate			
				Nitrate Soda, 520 lbs. in 1878; 275 lbs. in 1879, 1880, and 1881.	Ammonia-salts, 400 lbs. in 1878; 200 lbs. in 1879, 1880, and 1881.
					Cows' Urine, 6120 lbs. in 1879 only. (°)
					Rape Cake, 2000 lbs. in 1878 and 1880.

(1) In November 1879, Lime was applied to one land of Series 1, and to the adjoining land of Series 2, in addition to the other manures.
 (2) One of the two lands had received rape-cake in 1878.
 (3) "Superphosphate of Lime"—in all cases, made from 300 lbs. Bone-ash, 225 lbs. Sulphuric acid sp. gr. 1.7 (and water).

EXPERIMENTS ON ROOT-CROPS.—BARN FIELD.

Experiments with Turnips were commenced in 1843. Eight acres, divided into numerous Plots, were set apart for the purpose, and the crop was grown for ten consecutive years on the same land; "Norfolk Whites" 1843-1848, and "Swedes" 1849-1852; on some Plots without manure, and on others with different descriptions of manure. Barley was then grown for three consecutive seasons, 1853-1855, without manure, in order to test the comparative corn-growing condition of the different Plots, and also to equalise their condition, as far as possible, by the exhaustion of some of the most active and immediately available constituents supplied by the previous manuring. A new series of experiments with Swedes was arranged in 1856, having regard to the character of the manures previously applied on the different Plots, and to the results previously obtained. This second series was continued for fifteen years, namely, from 1856 to 1870 inclusive. The results obtained with Norfolk Whites in the first three years, 1843, 1844, and 1845, were published in the 'Journal of the Royal Agricultural Society of England,' vol. viii. Part II, 1847; and an abstract of the results obtained from 1845 to 1870 inclusive, is given in the Table below. During the five years, 1871-1875, the land was devoted to experiments with Sugar-Beet, for particulars of which see pp. 18 and 19. In 1876 experiments with Mangold-wurzel were substituted, and are still in progress (see pages 20 and 21).

(Area under experiment, about 8 acres; quantities, average, per acre, per annum.)

NORFOLK WHITE TURNIPS; FOUR SEASONS, 1845-1848; Roots and Leaves carted off the Land.

SERIES 1. Manures as under; no Cross-dressing.		Each Plot as Series 1, and Cross-dressed as under—											
		SERIES 2. No Cross-dressing.		SERIES 3. 160 lbs. Sulphate Ammonia. 75 lbs. Muriate Ammonia.		SERIES 4. 160 lbs. Sulphate Ammonia. 75 lbs. Muriate Ammonia. 1840 lbs. Rape-cake.		SERIES 5. 1840 lbs. Rape-cake.					
		Average Produce, per Acre, per Annum.											
PLOTS.		Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.	
		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.
3	Gypsum 1845; without Manure 1846 and since (average 1846, 7, 8)	1	4	0	17	1	7	1	0	5	10	3	19
4	Superphosphate, each year; Potass, Soda, and Magnesia, 1847-8 ..	8	1	2	15	9	15	4	3	10	5	6	1
5	Superphosphate, each year;	8	16	2	19	9	18	4	8	10	1	6	3
6	Superphosphate, each year; and Potass 1847-8	8	0	2	19	9	16	5	4	10	7	6	17
7													

SWEDISH TURNIPS; FOUR SEASONS, 1849-1852; Roots and Leaves carted off the Land (excepting 1849, when the Leaves were too small to weigh or remove).

SERIES 1. Manures as under; no Cross-dressing.		Each Plot as Series 1, and Cross-dressed, as under, in 1849 and 1850. No Cross-dressing in 1851 and 1852.												
		SERIES 2. No Cross-dressing.		SERIES 3. 200 lbs. Ammonia-salts.		SERIES 4. 200 lbs. Ammonia-salts. 2000 lbs. Rape-cake.		SERIES 5. 2000 lbs. Rape-cake.						
		Average Produce, per Acre, per Annum.												
PLOTS.		Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.		
		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	
3	Without Manure, 1846 and since	2	6	0	6	3	17	0	6	7	0	17	7	14
4	Superphosphate, Sulphates Potass and Magnesia, and Soda-ash ..	7	17	0	10	9	9	0	11	13	1	0	18	12
5	Superphosphate	7	9	0	11	8	14	0	13	11	4	1	1	10
6	Superphosphate, and Sulphate Potass	6	16	0	9	8	14	0	10	12	8	0	17	11
7														

BARLEY, without Manure (after Roots manured as above); THREE SEASONS, 1853-1855.

PLOTS.		SERIES 1.		SERIES 2.		SERIES 3.		SERIES 4.		SERIES 5.	
		Dressed Corn.	Straw.	Dressed Corn.	Straw.	Dressed Corn.	Straw.	Dressed Corn.	Straw.	Dressed Corn.	Straw.
		Bushels.	Cwts.	Bushels.	Cwts.	Bushels.	Cwts.	Bushels.	Cwts.	Bushels.	Cwts.
3	18 $\frac{3}{4}$	12 $\frac{1}{2}$	20 $\frac{1}{2}$	12 $\frac{3}{4}$	20 $\frac{1}{2}$	12 $\frac{3}{4}$	24 $\frac{1}{2}$	15 $\frac{3}{4}$	25 $\frac{1}{2}$	16
4	20 $\frac{1}{2}$	12 $\frac{1}{2}$	22 $\frac{1}{2}$	13 $\frac{3}{4}$	22 $\frac{1}{2}$	13 $\frac{3}{4}$	25	14 $\frac{3}{4}$	25 $\frac{1}{2}$	14 $\frac{5}{8}$
5	21	11 $\frac{3}{4}$	23	12 $\frac{3}{4}$	23	12 $\frac{3}{4}$	26 $\frac{1}{2}$	15	27	15 $\frac{3}{8}$
6	18 $\frac{3}{4}$	10 $\frac{3}{4}$	20 $\frac{1}{2}$	11 $\frac{3}{4}$	25	14 $\frac{3}{4}$	25	14 $\frac{3}{4}$	25	14 $\frac{3}{4}$
7											

SWEDISH TURNIPS; FIFTEEN SEASONS, 1856-1870.(?) Roots and Leaves carted off the Land.

SERIES 1. Manures as under; no Cross-dressing.		Each Plot as Series 1, and Cross-dressed as under—											
		SERIES 2. 5 years, 1856-1860. 3000 lbs. Saw-dust. 328 lbs. Nitric Acid.		SERIES 3. 5 years, 1856-1860. 200 lbs. Ammonia-salts.		SERIES 4. 5 years, 1856-1860. 200 lbs. Ammonia-salts. 3000 lbs. Sawdust.		SERIES 5. 5 years, 1856-1860. 3000 lbs. Sawdust.					
		Average Produce, per Acre, per Annum.											
PLOTS.		Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.	
		Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.
1	Farmyard Manure, 14 tons	6	4	0	17	7	9	1	2	8	8	1	4
2	Farmyard Manure, 14 tons, and Superphosphate	6	7	0	16	7	13	1	3	8	5	1	5
3	Without Manure, 1846, and since	0	11	0	3	0	19	0	4	0	13	0	3
4	Superphosph., each year; Sulph. Potass, Soda, and Magnesia, 1856-60	2	16	0	8	5	2	0	16	4	12	0	14
5	Superphosphate, each year	2	12	0	9	4	13	0	18	3	16	0	15
6	Superphosphate, each year; Sulphate Potass, 1856-1860	2	7	0	7	4	11	0	14	4	5	0	13
7	Superphosph., each year; Sulph. Potass, and 36 $\frac{1}{2}$ Amm.-salts, 1856-60	2	12	0	7	4	13	0	14	4	12	0	14
8	Unman. 1853, and since; previously part Unman.; part Superphosph.	1	3	0	4	1	13	0	5	1	2	0	5

NOTE.—"Sulphate of Ammonia" is estimated to contain 23 per cent. Ammonia and "Muriate of Ammonia" 21 per cent. "Ammonia-salts" in each case, equal parts Sulphate and Muriate of Ammonia of commerce; and the mixture is estimated to contain 25 per cent. Ammonia. The 328 lbs. Nitric Acid (Sp. gr. 1.35), mixed with sawdust, and used as a cross-dressing on the Plots of Series 2, from 1856-1860, were estimated to contain Nitrogen = 50 lbs. Ammonia.

(?) The crops of 1859 and 1860 failed, and were ploughed in; but, as the manures were applied, and there would be accumulation within the soil for the succeeding crops, the average produce is calculated as for 15 years, that is the produce of the 13 years in, in each case, divided by 15.

EXPERIMENTS ON SUGAR BEET—BARN FIELD—continued.

SUMMARY OF THE COMPOSITION OF THE SUGAR-BEET ROOTS.

As it will be some time before we shall be able to report fully the results obtained illustrating the influence of different manures, and different seasons, on the composition of Sugar-beet, an abstract of the analytical results obtained is given below. In interpreting the figures it must be borne in mind that with forty different experiments each year, and in each year 4 or 5 or more times as much produce on some plots as on others, it would be impossible to sample each at its best, and all in the same condition of ripeness. Each year the seed was sown on all the plots at the same time; and the samples (each consisting of the vertical fourths of 10 or 15 roots) were taken from all within a period of about a week, beginning with the ripest. It is obvious, however, that the smaller crops would be much riper than the larger ones. The dry matter, ash, and nitrogen, as given in the table, are determined in the roots themselves; but they have generally been determined in the expressed juice also. The sugar is determined in the juice; and calculated into its percentage in the roots, on the assumption that they contain uniformly 95 per cent. of juice. But, with roots varying so much in character of growth, size, and ripeness, this will not be the case. Nevertheless, the results so calculated, approximately, and usefully, represent both the actual and relative amounts of sugar in the various roots. According to recent experiments of Schiebler, and others, however, the percentage of the juice in the roots, reckoned from the determined percentage of dry matter in the juice, and in the roots, respectively, has been over-estimated. According to these new results, the amount of true juice will average more nearly 90, than 95 per cent. If this be established, the percentage of sugar in the roots will be less (perhaps $\frac{1}{5}$ to $\frac{1}{10}$ less) than given in the Table below. It need only further be observed that although, in comparable cases, the larger crops generally give a juice containing a lower percentage of sugar, and higher percentages of mineral matter and of nitrogen, yet the larger crops yielded very much more sugar over a given area of land.

For Manures and Produce, see facing page.	CROSS-DRESSED MANURES, PER ACRE, PER ANNUM.																			
	SERIES 1. No Cross-dressing.				SERIES 2. As Series 1, and Cross-dressed with 550 lbs. Nitrate Soda.				SERIES 3. As Series 1, and Cross-dressed with 400 lbs. "Ammonia-salts."				SERIES 4. As Series 1, and Cross-dressed with 2000 lbs. Rape-cake, and 400 lbs. "Ammonia-salts."				SERIES 5. As Series 1, and Cross-dressed with 2000 lbs. Rape-cake.			

FIRST SEASON, 1871. (Results in all cases the means of determinations made on two samples, collected at the end of October, and the end of November, respectively).

PLOTS.	Mean Per Cent. Total Dry Matter, Sugar, Mineral Matter (Crude Ash), and Nitrogen in the Roots.																			
	SERIES 1.				SERIES 2.				SERIES 3.				SERIES 4.				SERIES 5.			
	Dry Matter.	Sugar.	Ash.	Nitrogen.	Dry Matter.	Sugar.	Ash.	Nitrogen.	Dry Matter.	Sugar.	Ash.	Nitrogen.	Dry Matter.	Sugar.	Ash.	Nitrogen.	Dry Matter.	Sugar.	Ash.	Nitrogen.
1	17.04	11.77	0.821	0.142	14.83	9.76	0.945	0.184	16.07	11.05	0.934	0.246	14.73	9.36	1.021	0.244	13.44	10.25	0.892	0.192
2	17.24	11.91	0.826	0.146	15.03	9.80	0.970	0.200	15.12	9.95	0.977	0.213	14.80	9.23	0.988	0.249	16.11	10.80	0.909	
3	17.47	12.51	0.711		15.36	10.37	0.861		17.75	10.98	0.901		16.71	9.66	0.915		16.95	11.72	0.758	
4	18.07	12.99	0.738		15.72	10.81	0.828		18.68	11.87	0.907		16.87	9.90	1.002		16.61	11.69	0.767	
5	17.89	13.23	0.746		15.93	11.07	0.787		16.36	11.44	0.754		14.63	9.28	0.843		16.84	11.85	0.722	
6	18.09	13.00	0.778		15.29	10.47	0.856		16.33	11.51	0.843		15.28	9.71	0.856		17.05	12.08	0.812	
7	17.97	13.17	0.762		15.86	10.49	0.901		16.71	11.50	0.826		15.99	10.23	0.904		17.57	12.30	0.782	
8	18.32	13.02	0.791		15.98	11.07	0.856		16.08	10.88	0.764		14.90	9.33	0.806		16.73	11.93	0.747	

SECOND SEASON, 1872. (Samples collected early in November.)

1	18.23	12.97	0.874		17.07	12.04	0.973		17.07	11.95	0.962		17.17	12.07	0.930		17.75	12.35	0.925	
2	18.07	13.04	0.822		15.97	11.12	1.000		16.04	10.43	0.982		17.07	11.81	0.955		17.95	12.82	0.875	
3	19.22	13.99	0.767		17.83	12.78	0.823		19.62	14.38	0.691		17.87	12.60	0.720		19.12	13.95	0.683	
4	19.08	14.16	0.778	0.110	16.97	12.19	0.860	0.148	18.55	13.32	0.800	0.128	18.49	12.66	0.965	0.184	18.67	13.38	0.795	0.130
5	18.67	13.92	0.712	0.101	16.37	11.16	0.866	0.167	18.40	13.02	0.734	0.167	15.82	10.40	0.918	0.250	18.07	13.22	0.705	0.139
6	18.83	13.81	0.773	0.098	17.08	11.88	0.891	0.167	18.70	13.46	0.837	0.166	17.38	12.15	0.879	0.173	18.41	13.17	0.780	0.162
7	19.03	13.94	0.742		16.66	11.22	0.937		18.71	13.35	0.787		17.98	12.83	0.797		19.01	14.06	0.809	
8	18.69	..	0.701		16.84	..	0.911		0.790		18.00	..	0.738		18.95	..	0.685	

THIRD SEASON, 1873. (Samples collected from November 10 to November 14.)

1	17.62	12.73	0.924		16.64	11.20	0.947		16.76	11.33	0.965		18.80	10.21	1.267		16.88	11.64	0.887	
2	18.49	13.02	0.847		16.35	10.75	0.978		16.54	11.59	0.951		13.39	10.29	0.905		16.33	11.52	0.960	
3	18.96	13.84	0.710		16.97	11.89	0.843		18.76	13.07	0.762		16.00	11.24	0.755		17.94	14.20	0.735	
4	18.80	13.81	0.796	0.132	17.97	12.06	0.934	0.181	18.31	13.11	0.877	0.161	16.67	11.21	0.974	0.187	18.30	13.18	0.861	0.149
5	19.25	14.27	0.679	0.121	16.89	11.50	0.847	0.184	18.24	13.17	0.604	0.186	16.66	11.65	0.734	0.227	18.93	13.48	0.664	0.160
6	19.64	14.35	0.757	0.119	17.94	12.49	0.810	0.169	18.42	13.21	0.894	0.140	17.56	11.89	0.906	0.212	18.22	12.97	0.845	0.148
7	19.63	14.43	0.747		17.42	11.71	0.907		18.81	13.72	0.858		17.68	12.11	0.870		19.00	13.09	0.852	
8	20.22	14.66	0.742		16.50	10.90	0.917		18.47	13.20	0.756		16.54	10.83	0.782		18.06	13.07	0.695	

FOURTH SEASON, 1874 (1). Mineral Manures as in 1872 and 1873; but no Farmyard Manure, or cross-dressings of Nitrate Soda, Ammonia-salts, or Rape-cake. (Samples collected in the middle of November.)

1	14.66	11.15	1.100		14.27	10.16	1.089		14.35	9.79	1.112		13.53	10.24	1.029		14.39	10.85	0.972	
2	15.00	12.75	1.022		13.84	9.93	1.082		14.24	10.11	1.081		14.59	10.11	0.970		14.34	10.88	0.933	
3	17.45	13.20	0.792		15.60	10.17	0.990		16.05	11.69	0.863		15.54	11.44	0.861		15.04	11.16	0.864	
4	18.54	13.10	0.721		14.00	9.73	0.840		16.70	12.41	0.921		17.17	11.62	1.026		14.98	12.55	1.027	
5	18.06	13.01	0.668		14.91	9.78	0.898		16.87	12.42	0.893		14.89	11.55	0.746		16.26	10.82	0.796	
6	17.83	12.99	0.752		15.95	10.50	0.859		16.70	13.69	0.865		15.30	12.05	0.938		16.29	11.04	0.879	
7	16.88	..	0.730		15.56	..	0.903		17.74	..	0.784		16.08	..	0.907		15.50	..	0.868	
8	18.76	..	0.726		15.30	..	0.890		17.35	..	0.771		15.48	..	0.841		16.51	..	0.772	

FIFTH SEASON, 1875. Mineral Manures as in 1872, 1873, and 1874; but no Farmyard Manure, or cross-dressings of Nitrate Soda, Ammonia-salts, or Rape-cake. (Samples collected in the middle of November.)

1	16.02	11.71	0.749		16.16	11.85	0.751		16.33	11.51	0.814		16.29	12.02	0.840		16.13	11.57	0.780	
2	16.08	11.72	0.784		15.67	11.22	0.687		15.43	10.77	0.863		15.70	10.90	0.770		15.92	11.71	0.793	
3	17.29	12.78	0.671		15.66	11.52	0.720		17.52	12.80	0.675		15.90	11.45	0.652		16.48	12.12	0.641	
4	16.67	12.11	0.773	0.103	16.10	12.06	0.751	0.112	17.07	12.32	0.755		16.56	11.89	0.758	0.125	16.24	11.69	0.775	0.121
5	16.94	12.99	0.686	0.107	16.53	12.09	0.722	0.125	16.55	12.08	0.683	0.122	15.34	11.20	0.682	0.152	15.86	11.81	0.622	0.123
6	18.04	12.66	0.782	0.127	16.78	12.47	0.762	0.123	16.19	12.21	0.752	0.136	16.21	11.58	0.777	0.158	16.53	12.09	0.759	0.141
7	17.51	..	0.750		16.22	..	0.874		16.50	..	0.802		15.88	..	0.856		16.38	..	0.866	
8	16.81	..	0.770		16.01	..	0.812		16.56	..	0.767		15.96	..	0.768		15.86	..	0.658	

(1) Owing to the deficiency of Rain for some time after sowing a large proportion of the plants failed. Some were transplanted on plots 1, but not on the other plots; and eventually the plant was (excepting on plots 1) upon the whole very deficient and irregular, the remaining plants being larger than usual.

EXPERIMENTS ON SUGAR BEET—BARN FIELD—continued.

SUMMARY OF THE COMPOSITION OF THE SUGAR-BEET ROOTS.

As it will be some time before we shall be able to report fully the results obtained illustrating the influence of different manures, and different seasons, on the composition of Sugar-beet, an abstract of the analytical results obtained is given below. In interpreting the figures it must be borne in mind that with forty different experiments each year, and in each year 4 or 5 or more times as much produce on some plots as on others, it would be impossible to sample each at its best, and all in the same condition of ripeness. Each year the seed was sown on all the Plots at the same time; and the samples (each consisting of the vertical fourths of 10 or 15 roots) were taken from all within a period of about a week, beginning with the ripest. It is obvious, however, that the smaller crops would be much riper than the larger ones. The dry matter, ash, and nitrogen, as given in the table, are determined in the roots themselves; but they have generally been determined in the expressed juice also. The sugar is determined in the juice; and calculated into its percentage in the roots, on the assumption that they contain uniformly 95 per cent. of juice. But, with roots varying so much in character of growth, size, and ripeness, this will not be the case. Nevertheless, the results so calculated, approximately, and usefully, represent both the actual and relative amounts of sugar in the various roots. According to recent experiments of Schiebeler, and others, however, the percentage of the juice in the roots, reckoned from the determined percentage of dry matter in the juice, and in the roots, respectively, has been over-estimated. According to these new results, the amount of true juice will average more nearly 90, than 95 per cent. If this be established, the percentage of sugar in the roots will be less (perhaps $\frac{1}{5}$ to $\frac{1}{10}$ less) than given in the Table below. It need only further be observed that although, in comparable cases, the larger crops generally give a juice containing a lower percentage of sugar, and higher percentages of mineral matter and of nitrogen, yet the larger crops yielded very much more sugar over a given area of land.

For Manures and Produce, see facing page.	CROSS-DRESSED MANURES, PER ACRE, PER ANNUM.																			
	SERIES 1. No Cross-dressing.				SERIES 2. As Series 1, and Cross-dressed with 550 lbs. Nitrate Soda.				SERIES 3. As Series 1, and Cross-dressed with 400 lbs. "Ammonia-salts."				SERIES 4. As Series 1, and Cross-dressed with 2000 lbs. Rape-cake, and 400 lbs. "Ammonia-salts."				SERIES 5. As Series 1, and Cross-dressed with 2000 lbs. Rape-cake.			
FIRST SEASON, 1871. (Results in all cases the means of determinations made on two samples, collected at the end of October, and the end of November, respectively.)																				
Mean Per Cent. Total Dry Matter, Sugar, Mineral Matter (Crude Ash), and Nitrogen in the Roots.																				
PLOTS.	SERIES 1.				SERIES 2.				SERIES 3.				SERIES 4.				SERIES 5.			
	Dry Matter.	Sugar.	Ash.	Nitrogen.	Dry Matter.	Sugar.	Ash.	Nitrogen.	Dry Matter.	Sugar.	Ash.	Nitrogen.	Dry Matter.	Sugar.	Ash.	Nitrogen.	Dry Matter.	Sugar.	Ash.	Nitrogen.
1	17.04	11.77	0.821	0.142	14.83	9.76	0.945	0.184	16.07	11.05	0.934	0.246	14.73	9.36	1.021	0.244	15.44	10.25	0.892	0.192
2	17.24	11.91	0.826	0.146	15.03	9.80	0.970	0.200	15.12	9.95	0.977	0.213	14.80	9.23	0.988	0.249	16.11	10.80	0.909	
3	17.47	12.51	0.711		15.36	10.37	0.861		17.75	10.98	0.901		16.71	9.66	0.915		16.95	11.72	0.758	
4	18.07	12.99	0.738		15.72	10.81	0.828		18.68	11.87	0.907		16.87	9.90	1.002		16.61	11.69	0.767	
5	17.89	13.23	0.746		15.93	11.07	0.787		16.36	11.44	0.754		14.63	9.28	0.843		16.84	11.85	0.722	
6	18.09	13.00	0.778		15.29	10.47	0.856		16.33	11.51	0.843		15.28	9.71	0.956		17.05	12.08	0.812	
7	17.97	13.17	0.762		15.86	10.49	0.901		16.71	11.50	0.826		15.99	10.23	0.904		17.57	12.30	0.782	
8	18.32	13.02	0.791		15.98	11.07	0.856		16.08	10.88	0.764		14.90	9.33	0.806		16.73	11.93	0.747	
SECOND SEASON, 1872. (Samples collected early in November.)																				
1	18.23	12.97	0.874		17.07	12.04	0.973		17.07	11.95	0.962		17.17	12.07	0.930		17.75	12.35	0.925	
2	18.07	13.04	0.822		15.97	11.12	1.000		16.04	10.43	0.982		17.07	11.81	0.965		17.95	12.82	0.875	
3	19.22	13.99	0.767		17.83	12.78	0.823		19.62	14.38	0.691		17.87	12.60	0.720		19.12	13.95	0.683	
4	19.08	14.16	0.778	0.110	16.97	12.19	0.860	0.148	18.55	13.32	0.800	0.128	18.49	12.66	0.965	0.184	18.67	13.38	0.795	0.139
5	18.67	13.92	0.712	0.101	16.37	11.16	0.866	0.167	18.40	13.02	0.734	0.167	15.82	10.40	0.918	0.250	18.07	13.22	0.705	0.159
6	18.83	13.81	0.772	0.098	17.08	11.88	0.891	0.167	18.70	13.46	0.837	0.166	17.38	12.15	0.879	0.173	18.41	13.17	0.780	0.162
7	19.03	13.94	0.742		16.66	11.22	0.937		18.71	13.35	0.787		17.98	12.83	0.797		19.01	14.06	0.809	
8	18.69	..	0.701		16.84	..	0.911		0.790		18.00	..	0.738		18.95	..	0.685	
THIRD SEASON, 1873. (Samples collected from November 10 to November 14.)																				
1	17.62	12.73	0.824		16.64	11.20	0.947		16.76	11.33	0.965		18.80	10.21	1.267		16.88	11.64	0.887	
2	18.49	13.02	0.847		16.35	10.75	0.973		16.54	11.59	0.951		13.39	10.29	0.905		16.33	11.52	0.960	
3	18.96	13.84	0.710		16.97	11.89	0.843		18.76	13.07	0.762		16.00	11.24	0.755		17.94	14.20	0.735	
4	18.80	13.81	0.796	0.132	17.97	12.06	0.934	0.181	18.31	13.11	0.877	0.161	16.67	11.21	0.974	0.187	18.30	13.18	0.861	0.149
5	19.25	14.27	0.679	0.121	16.89	11.50	0.847	0.184	18.24	13.17	0.604	0.186	16.66	11.65	0.734	0.227	18.93	13.48	0.664	0.160
6	19.64	14.35	0.757	0.119	17.94	12.49	0.810	0.169	18.42	13.21	0.894	0.140	17.56	11.89	0.906	0.212	18.22	12.97	0.845	0.148
7	19.63	14.43	0.747		17.42	11.71	0.907		18.81	13.72	0.858		17.68	12.11	0.870		19.00	13.09	0.852	
8	20.22	14.66	0.742		16.50	10.90	0.917		18.47	13.20	0.756		16.54	10.83	0.782		18.06	13.07	0.695	
FOURTH SEASON, 1874 (*). Mineral Manures as in 1872 and 1873; but no Farmyard Manure, or cross-dressings of Nitrate Soda, Ammonia-salts, or Rape-cake. (Samples collected in the middle of November.)																				
1	14.66	11.15	1.100		14.27	10.16	1.089		14.35	9.79	1.112		13.53	10.24	1.029		14.39	10.85	0.972	
2	15.00	12.75	1.022		13.84	9.93	1.082		14.24	10.11	1.081		14.59	10.11	0.970		14.34	10.88	0.933	
3	17.45	13.20	0.792		15.60	10.17	0.990		16.05	11.69	0.863		15.54	11.44	0.861		15.04	11.16	0.864	
4	18.54	13.10	0.721		14.00	9.73	0.840		16.70	12.41	0.921		17.17	11.62	1.026		14.98	12.55	1.027	
5	18.06	13.01	0.668		14.91	9.78	0.898		16.87	12.42	0.833		14.89	11.55	0.746		16.26	10.82	0.796	
6	17.83	12.99	0.752		15.95	10.50	0.859		16.70	13.69	0.865		15.30	12.05	0.938		16.29	11.04	0.879	
7	16.88	..	0.730		15.56	..	0.903		17.74	..	0.784		16.08	..	0.907		15.50	..	0.868	
8	18.76	..	0.726		15.30	..	0.890		17.35	..	0.771		15.48	..	0.841		16.61	..	0.772	
FIFTH SEASON, 1875. Mineral Manures as in 1872, 1873, and 1874; but no Farmyard Manure, or cross-dressings of Nitrate Soda, Ammonia-salts, or Rape-cake. (Samples collected in the middle of November.)																				
1	16.02	11.71	0.749		16.16	11.85	0.751		16.33	11.51	0.814		16.29	12.02	0.840		16.13	11.57	0.780	
2	16.08	11.72	0.784		15.67	11.22	0.687		15.43	10.77	0.863		15.70	10.90	0.770		15.92	11.71	0.793	
3	17.29	12.78	0.671		15.66	11.52	0.720		17.52	12.80	0.675		15.90	11.45	0.652		16.48	12.12	0.641	
4	16.67	12.11	0.773	0.103	16.10	12.06	0.751	0.112	17.07	12.32	0.755		16.56	11.89	0.758	0.125	16.24	11.69	0.775	0.121
5	16.94	12.99	0.686	0.107	16.53	12.09	0.722	0.125	16.55	12.08	0.683	0.122	15.34	11.20	0.682	0.152	15.86	11.81	0.622	0.123
6	18.04	12.66	0.782	0.127	16.78	12.47	0.762	0.123	16.19	12.21	0.752	0.136	16.21	11.58	0.777	0.158	16.33	12.09	0.759	0.141
7	17.51	..	0.730		16.22	..	0.874		16.50	..	0.802		15.88	..	0.856		16.38	..	0.866	
8	16.81	..	0.770		16.01	..	0.812		16.56	..	0.767		15.96	..	0.768		15.86	..	0.658	

(* Owing to the deficiency of Rain for some time after sowing a large proportion of the plants failed. Some were transplanted on plots 1, but not on the other plots; and eventually the plant was (excepting on plots 1) upon the whole very deficient and irregular, the remaining plants being larger than usual.

EXPERIMENTS ON MANGOLD WURZEL.—BARN FIELD (after SUGAR-BEET); commencing 1876.

The arrangement of the Plots is precisely the same as previously for Sugar-beet, excepting that Plot 9, which was unmanured for Sugar-beet, and also previously for Swedes, is now added as a manured Plot. With this exception, the manures are also substantially the same as previously for Sugar-beet; in fact, precisely the same as for the Sugar-beet in 1872 and 1873. Seed, Yellow Globe; dibbled on ridges, rows 26 inches apart; plants 11 inches apart in the rows (?). Area under experiment about 8 acres. Roots all carted off; Leaves weighed, spread on the respective Plots, and ploughed in.

PLOTS.	SERIES 1.	MANURES PER ACRE PER ANNUM.				SERIES 2. As Series 1, and Cross-dressed with 550 lbs. Nitrate Soda.	SERIES 3. As Series 1, and Cross-dressed with 400 lbs. "Ammonia- salts."	SERIES 4. As Series 1, and Cross-dressed with 2000 lbs. Rape-cake and 400 lbs. "Am- monia-salts."	SERIES 5. As Series 1, and Cross-dressed with 2000 lbs. Rape-cake.																																																																																																																																																																																																																															
		PRODUCE PER ACRE.																																																																																																																																																																																																																																						
FIRST SEASON, 1876. Seed dibbled, May 22-26. Crop taken up, Nov. 3-17.																																																																																																																																																																																																																																								
<table border="1"> <thead> <tr> <th colspan="2">Roots.</th> <th colspan="2">Leaves.</th> <th colspan="2">Roots.</th> <th colspan="2">Leaves.</th> <th colspan="2">Roots.</th> <th colspan="2">Leaves.</th> </tr> <tr> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>Farmyard Manure (14 tons)</td> <td>19</td> <td>12</td> <td>4</td> <td>9</td> <td>25</td> <td>2</td> <td>7</td> <td>5</td> <td>29</td> <td>19</td> <td>7</td> <td>12</td> <td>31</td> <td>9</td> <td>10</td> <td>5</td> <td>24</td> <td>9</td> <td>5</td> <td>19</td> </tr> <tr> <td>2</td> <td>Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..</td> <td>19</td> <td>13</td> <td>4</td> <td>6</td> <td>27</td> <td>13</td> <td>7</td> <td>3</td> <td>29</td> <td>8</td> <td>7</td> <td>10</td> <td>30</td> <td>18</td> <td>9</td> <td>16</td> <td>29</td> <td>19</td> <td>6</td> <td>12</td> </tr> <tr> <td>3</td> <td>Without Manure (1846, and since)</td> <td>6</td> <td>10</td> <td>1</td> <td>14</td> <td>20</td> <td>13</td> <td>5</td> <td>12</td> <td>14</td> <td>3</td> <td>4</td> <td>10</td> <td>19</td> <td>19</td> <td>7</td> <td>7</td> <td>17</td> <td>4</td> <td>4</td> <td>15</td> </tr> <tr> <td>4</td> <td>{ 3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride { Sodium (common salt), 200 lbs. Sulphate Magnesia</td> <td>8</td> <td>8</td> <td>1</td> <td>15</td> <td>25</td> <td>1</td> <td>6</td> <td>0</td> <td>19</td> <td>19</td> <td>4</td> <td>9</td> <td>30</td> <td>8</td> <td>8</td> <td>13</td> <td>25</td> <td>8</td> <td>5</td> <td>10</td> </tr> <tr> <td>5</td> <td>3½ cwt. Superphosphate</td> <td>7</td> <td>10</td> <td>1</td> <td>14</td> <td>21</td> <td>0</td> <td>5</td> <td>14</td> <td>13</td> <td>10</td> <td>5</td> <td>1</td> <td>17</td> <td>2</td> <td>7</td> <td>14</td> <td>17</td> <td>17</td> <td>5</td> <td>17</td> </tr> <tr> <td>6</td> <td>3½ cwt. Superphosphate, 500 lbs. Sulphate Potass</td> <td>6</td> <td>16</td> <td>1</td> <td>12</td> <td>21</td> <td>2</td> <td>5</td> <td>8</td> <td>17</td> <td>15</td> <td>4</td> <td>13</td> <td>26</td> <td>8</td> <td>9</td> <td>0</td> <td>20</td> <td>10</td> <td>5</td> <td>4</td> </tr> <tr> <td>7</td> <td>3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)</td> <td>8</td> <td>13</td> <td>2</td> <td>3</td> <td>22</td> <td>11</td> <td>5</td> <td>14</td> <td>19</td> <td>2</td> <td>5</td> <td>11</td> <td>27</td> <td>2</td> <td>9</td> <td>9</td> <td>20</td> <td>12</td> <td>5</td> <td>15</td> </tr> <tr> <td>8</td> <td>Unmanured, 1853, and since; previously part Unman., part Superphos.</td> <td>5</td> <td>9</td> <td>1</td> <td>10</td> <td>15</td> <td>16</td> <td>5</td> <td>3</td> <td>11</td> <td>17</td> <td>4</td> <td>16</td> <td>18</td> <td>2</td> <td>7</td> <td>11</td> <td>15</td> <td>12</td> <td>4</td> <td>18</td> </tr> <tr> <td>9</td> <td>Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>25</td> <td>14</td> <td>7</td> <td>6</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> </tr> </tbody> </table>											Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.		Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	1	Farmyard Manure (14 tons)	19	12	4	9	25	2	7	5	29	19	7	12	31	9	10	5	24	9	5	19	2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..	19	13	4	6	27	13	7	3	29	8	7	10	30	18	9	16	29	19	6	12	3	Without Manure (1846, and since)	6	10	1	14	20	13	5	12	14	3	4	10	19	19	7	7	17	4	4	15	4	{ 3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride { Sodium (common salt), 200 lbs. Sulphate Magnesia	8	8	1	15	25	1	6	0	19	19	4	9	30	8	8	13	25	8	5	10	5	3½ cwt. Superphosphate	7	10	1	14	21	0	5	14	13	10	5	1	17	2	7	14	17	17	5	17	6	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass	6	16	1	12	21	2	5	8	17	15	4	13	26	8	9	0	20	10	5	4	7	3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)	8	13	2	3	22	11	5	14	19	2	5	11	27	2	9	9	20	12	5	15	8	Unmanured, 1853, and since; previously part Unman., part Superphos.	5	9	1	10	15	16	5	3	11	17	4	16	18	2	7	11	15	12	4	18	9	Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)	25	14	7	6
Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.																																																																																																																																																																																																																														
Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.																																																																																																																																																																																																																													
1	Farmyard Manure (14 tons)	19	12	4	9	25	2	7	5	29	19	7	12	31	9	10	5	24	9	5	19																																																																																																																																																																																																																			
2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..	19	13	4	6	27	13	7	3	29	8	7	10	30	18	9	16	29	19	6	12																																																																																																																																																																																																																			
3	Without Manure (1846, and since)	6	10	1	14	20	13	5	12	14	3	4	10	19	19	7	7	17	4	4	15																																																																																																																																																																																																																			
4	{ 3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride { Sodium (common salt), 200 lbs. Sulphate Magnesia	8	8	1	15	25	1	6	0	19	19	4	9	30	8	8	13	25	8	5	10																																																																																																																																																																																																																			
5	3½ cwt. Superphosphate	7	10	1	14	21	0	5	14	13	10	5	1	17	2	7	14	17	17	5	17																																																																																																																																																																																																																			
6	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass	6	16	1	12	21	2	5	8	17	15	4	13	26	8	9	0	20	10	5	4																																																																																																																																																																																																																			
7	3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)	8	13	2	3	22	11	5	14	19	2	5	11	27	2	9	9	20	12	5	15																																																																																																																																																																																																																			
8	Unmanured, 1853, and since; previously part Unman., part Superphos.	5	9	1	10	15	16	5	3	11	17	4	16	18	2	7	11	15	12	4	18																																																																																																																																																																																																																			
9	Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)	25	14	7	6																																																																																																																																																																																																																			
SECOND SEASON, 1877. Seed dibbled, June 4-6 (Plots 8 and 9, June 11th). Crop taken up, Nov. 14-23.																																																																																																																																																																																																																																								
<table border="1"> <thead> <tr> <th colspan="2">Roots.</th> <th colspan="2">Leaves.</th> <th colspan="2">Roots.</th> <th colspan="2">Leaves.</th> <th colspan="2">Roots.</th> <th colspan="2">Leaves.</th> </tr> <tr> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>Farmyard Manure (14 tons)</td> <td>15</td> <td>7</td> <td>2</td> <td>1</td> <td>24</td> <td>13</td> <td>3</td> <td>14</td> <td>27</td> <td>1</td> <td>4</td> <td>4</td> <td>30</td> <td>5</td> <td>5</td> <td>5</td> <td>25</td> <td>18</td> <td>3</td> <td>4</td> </tr> <tr> <td>2</td> <td>Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..</td> <td>16</td> <td>14</td> <td>1</td> <td>19</td> <td>26</td> <td>8</td> <td>3</td> <td>12</td> <td>26</td> <td>18</td> <td>4</td> <td>6</td> <td>28</td> <td>15</td> <td>5</td> <td>9</td> <td>24</td> <td>12</td> <td>2</td> <td>19</td> </tr> <tr> <td>3</td> <td>Without Manure (1846, and since)</td> <td>5</td> <td>9</td> <td>1</td> <td>0</td> <td>16</td> <td>17</td> <td>3</td> <td>14</td> <td>8</td> <td>16</td> <td>3</td> <td>0</td> <td>13</td> <td>9</td> <td>3</td> <td>19</td> <td>13</td> <td>17</td> <td>2</td> <td>10</td> </tr> <tr> <td>4</td> <td>{ 3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride { Sodium (common salt), 200 lbs. Sulphate Magnesia</td> <td>6</td> <td>16</td> <td>1</td> <td>3</td> <td>21</td> <td>10</td> <td>3</td> <td>10</td> <td>16</td> <td>10</td> <td>2</td> <td>2</td> <td>27</td> <td>9</td> <td>3</td> <td>8</td> <td>21</td> <td>14</td> <td>1</td> <td>17</td> </tr> <tr> <td>5</td> <td>3½ cwt. Superphosphate</td> <td>6</td> <td>1</td> <td>0</td> <td>19</td> <td>20</td> <td>5</td> <td>3</td> <td>1</td> <td>12</td> <td>2</td> <td>2</td> <td>10</td> <td>15</td> <td>3</td> <td>3</td> <td>8</td> <td>18</td> <td>5</td> <td>3</td> <td>2</td> </tr> <tr> <td>6</td> <td>3½ cwt. Superphosphate, 500 lbs. Sulphate Potass</td> <td>5</td> <td>8</td> <td>0</td> <td>18</td> <td>20</td> <td>19</td> <td>2</td> <td>18</td> <td>15</td> <td>6</td> <td>1</td> <td>16</td> <td>24</td> <td>18</td> <td>3</td> <td>16</td> <td>19</td> <td>3</td> <td>1</td> <td>12</td> </tr> <tr> <td>7</td> <td>3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)</td> <td>7</td> <td>0</td> <td>1</td> <td>3</td> <td>22</td> <td>2</td> <td>3</td> <td>16</td> <td>16</td> <td>13</td> <td>2</td> <td>7</td> <td>25</td> <td>15</td> <td>5</td> <td>0</td> <td>20</td> <td>13</td> <td>2</td> <td>8</td> </tr> <tr> <td>8</td> <td>Unmanured, 1853, and since; previously part Unman., part Superphos.</td> <td>3</td> <td>19</td> <td>1</td> <td>3</td> <td>9</td> <td>17</td> <td>5</td> <td>4</td> <td>7</td> <td>4</td> <td>3</td> <td>10</td> <td>11</td> <td>9</td> <td>4</td> <td>11</td> <td>10</td> <td>3</td> <td>3</td> <td>3</td> </tr> <tr> <td>9</td> <td>Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>13</td> <td>17</td> <td>4</td> <td>0</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> </tr> </tbody> </table>											Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.		Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	1	Farmyard Manure (14 tons)	15	7	2	1	24	13	3	14	27	1	4	4	30	5	5	5	25	18	3	4	2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..	16	14	1	19	26	8	3	12	26	18	4	6	28	15	5	9	24	12	2	19	3	Without Manure (1846, and since)	5	9	1	0	16	17	3	14	8	16	3	0	13	9	3	19	13	17	2	10	4	{ 3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride { Sodium (common salt), 200 lbs. Sulphate Magnesia	6	16	1	3	21	10	3	10	16	10	2	2	27	9	3	8	21	14	1	17	5	3½ cwt. Superphosphate	6	1	0	19	20	5	3	1	12	2	2	10	15	3	3	8	18	5	3	2	6	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass	5	8	0	18	20	19	2	18	15	6	1	16	24	18	3	16	19	3	1	12	7	3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)	7	0	1	3	22	2	3	16	16	13	2	7	25	15	5	0	20	13	2	8	8	Unmanured, 1853, and since; previously part Unman., part Superphos.	3	19	1	3	9	17	5	4	7	4	3	10	11	9	4	11	10	3	3	3	9	Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)	13	17	4	0
Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.																																																																																																																																																																																																																														
Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.																																																																																																																																																																																																																													
1	Farmyard Manure (14 tons)	15	7	2	1	24	13	3	14	27	1	4	4	30	5	5	5	25	18	3	4																																																																																																																																																																																																																			
2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..	16	14	1	19	26	8	3	12	26	18	4	6	28	15	5	9	24	12	2	19																																																																																																																																																																																																																			
3	Without Manure (1846, and since)	5	9	1	0	16	17	3	14	8	16	3	0	13	9	3	19	13	17	2	10																																																																																																																																																																																																																			
4	{ 3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride { Sodium (common salt), 200 lbs. Sulphate Magnesia	6	16	1	3	21	10	3	10	16	10	2	2	27	9	3	8	21	14	1	17																																																																																																																																																																																																																			
5	3½ cwt. Superphosphate	6	1	0	19	20	5	3	1	12	2	2	10	15	3	3	8	18	5	3	2																																																																																																																																																																																																																			
6	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass	5	8	0	18	20	19	2	18	15	6	1	16	24	18	3	16	19	3	1	12																																																																																																																																																																																																																			
7	3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)	7	0	1	3	22	2	3	16	16	13	2	7	25	15	5	0	20	13	2	8																																																																																																																																																																																																																			
8	Unmanured, 1853, and since; previously part Unman., part Superphos.	3	19	1	3	9	17	5	4	7	4	3	10	11	9	4	11	10	3	3	3																																																																																																																																																																																																																			
9	Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)	13	17	4	0																																																																																																																																																																																																																			
THIRD SEASON, 1878. Seed dibbled, June 8-9 (Plot 9, June 11th). Crop taken up, Nov. 7-20.																																																																																																																																																																																																																																								
<table border="1"> <thead> <tr> <th colspan="2">Roots.</th> <th colspan="2">Leaves.</th> <th colspan="2">Roots.</th> <th colspan="2">Leaves.</th> <th colspan="2">Roots.</th> <th colspan="2">Leaves.</th> </tr> <tr> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>Farmyard Manure (14 tons)</td> <td>13</td> <td>5</td> <td>2</td> <td>16</td> <td>18</td> <td>15</td> <td>4</td> <td>4</td> <td>20</td> <td>11</td> <td>5</td> <td>6</td> <td>22</td> <td>4</td> <td>6</td> <td>3</td> <td>17</td> <td>1</td> <td>3</td> <td>13</td> </tr> <tr> <td>2</td> <td>Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..</td> <td>14</td> <td>16</td> <td>2</td> <td>19</td> <td>21</td> <td>4</td> <td>4</td> <td>15</td> <td>19</td> <td>15</td> <td>5</td> <td>3</td> <td>20</td> <td>18</td> <td>5</td> <td>17</td> <td>18</td> <td>17</td> <td>3</td> <td>15</td> </tr> <tr> <td>3</td> <td>Without Manure (1846, and since)</td> <td>3</td> <td>10</td> <td>1</td> <td>4</td> <td>10</td> <td>2</td> <td>2</td> <td>16</td> <td>4</td> <td>7</td> <td>2</td> <td>11</td> <td>6</td> <td>11</td> <td>3</td> <td>7</td> <td>6</td> <td>3</td> <td>2</td> <td>17</td> </tr> <tr> <td>4</td> <td>{ 3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride { Sodium (common salt), 200 lbs. Sulphate Magnesia</td> <td>5</td> <td>9</td> <td>1</td> <td>7</td> <td>18</td> <td>10</td> <td>4</td> <td>6</td> <td>14</td> <td>3</td> <td>2</td> <td>12</td> <td>21</td> <td>2</td> <td>4</td> <td>14</td> <td>15</td> <td>19</td> <td>3</td> <td>2</td> </tr> <tr> <td>5</td> <td>3½ cwt. Superphosphate</td> <td>4</td> <td>14</td> <td>1</td> <td>8</td> <td>14</td> <td>11</td> <td>3</td> <td>18</td> <td>8</td> <td>2</td> <td>3</td> <td>6</td> <td>8</td> <td>4</td> <td>3</td> <td>3</td> <td>8</td> <td>1</td> <td>3</td> <td>6</td> </tr> <tr> <td>6</td> <td>3½ cwt. Superphosphate, 500 lbs. Sulphate Potass</td> <td>3</td> <td>18</td> <td>1</td> <td>3</td> <td>15</td> <td>1</td> <td>3</td> <td>7</td> <td>12</td> <td>0</td> <td>2</td> <td>14</td> <td>15</td> <td>3</td> <td>4</td> <td>11</td> <td>12</td> <td>5</td> <td>3</td> <td>3</td> </tr> <tr> <td>7</td> <td>3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)</td> <td>5</td> <td>8</td> <td>1</td> <td>9</td> <td>13</td> <td>18</td> <td>3</td> <td>1</td> <td>11</td> <td>18</td> <td>2</td> <td>18</td> <td>14</td> <td>0</td> <td>4</td> <td>5</td> <td>11</td> <td>19</td> <td>3</td> <td>8</td> </tr> <tr> <td>8</td> <td>Unmanured, 1853, and since; previously part Unman., part Superphos.</td> <td>2</td> <td>13</td> <td>1</td> <td>4</td> <td>11</td> <td>19</td> <td>4</td> <td>7</td> <td>6</td> <td>13</td> <td>3</td> <td>5</td> <td>6</td> <td>12</td> <td>4</td> <td>10</td> <td>6</td> <td>4</td> <td>3</td> <td>5</td> </tr> <tr> <td>9</td> <td>Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>15</td> <td>17</td> <td>5</td> <td>9</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> </tr> </tbody> </table>											Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.		Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	1	Farmyard Manure (14 tons)	13	5	2	16	18	15	4	4	20	11	5	6	22	4	6	3	17	1	3	13	2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..	14	16	2	19	21	4	4	15	19	15	5	3	20	18	5	17	18	17	3	15	3	Without Manure (1846, and since)	3	10	1	4	10	2	2	16	4	7	2	11	6	11	3	7	6	3	2	17	4	{ 3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride { Sodium (common salt), 200 lbs. Sulphate Magnesia	5	9	1	7	18	10	4	6	14	3	2	12	21	2	4	14	15	19	3	2	5	3½ cwt. Superphosphate	4	14	1	8	14	11	3	18	8	2	3	6	8	4	3	3	8	1	3	6	6	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass	3	18	1	3	15	1	3	7	12	0	2	14	15	3	4	11	12	5	3	3	7	3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)	5	8	1	9	13	18	3	1	11	18	2	18	14	0	4	5	11	19	3	8	8	Unmanured, 1853, and since; previously part Unman., part Superphos.	2	13	1	4	11	19	4	7	6	13	3	5	6	12	4	10	6	4	3	5	9	Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)	15	17	5	9
Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.																																																																																																																																																																																																																														
Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.																																																																																																																																																																																																																													
1	Farmyard Manure (14 tons)	13	5	2	16	18	15	4	4	20	11	5	6	22	4	6	3	17	1	3	13																																																																																																																																																																																																																			
2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..	14	16	2	19	21	4	4	15	19	15	5	3	20	18	5	17	18	17	3	15																																																																																																																																																																																																																			
3	Without Manure (1846, and since)	3	10	1	4	10	2	2	16	4	7	2	11	6	11	3	7	6	3	2	17																																																																																																																																																																																																																			
4	{ 3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride { Sodium (common salt), 200 lbs. Sulphate Magnesia	5	9	1	7	18	10	4	6	14	3	2	12	21	2	4	14	15	19	3	2																																																																																																																																																																																																																			
5	3½ cwt. Superphosphate	4	14	1	8	14	11	3	18	8	2	3	6	8	4	3	3	8	1	3	6																																																																																																																																																																																																																			
6	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass	3	18	1	3	15	1	3	7	12	0	2	14	15	3	4	11	12	5	3	3																																																																																																																																																																																																																			
7	3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)	5	8	1	9	13	18	3	1	11	18	2	18	14	0	4	5	11	19	3	8																																																																																																																																																																																																																			
8	Unmanured, 1853, and since; previously part Unman., part Superphos.	2	13	1	4	11	19	4	7	6	13	3	5	6	12	4	10	6	4	3	5																																																																																																																																																																																																																			
9	Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)	15	17	5	9																																																																																																																																																																																																																			
FOURTH SEASON, 1879. Seed dibbled, May 13-15. Crop taken up, Nov. 11-20.																																																																																																																																																																																																																																								
<table border="1"> <thead> <tr> <th colspan="2">Roots.</th> <th colspan="2">Leaves.</th> <th colspan="2">Roots.</th> <th colspan="2">Leaves.</th> <th colspan="2">Roots.</th> <th colspan="2">Leaves.</th> </tr> <tr> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>Farmyard Manure (14 tons)</td> <td>6</td> <td>3</td> <td>1</td> <td>15</td> <td>9</td> <td>8</td> <td>2</td> <td>9</td> <td>12</td> <td>6</td> <td>3</td> <td>11</td> <td>13</td> <td>16</td> <td>3</td> <td>15</td> <td>10</td> <td>14</td> <td>2</td> <td>12</td> </tr> <tr> <td>2</td> <td>Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..</td> <td>6</td> <td>13</td> <td>1</td> <td>16</td> <td>11</td> <td>11</td> <td>2</td> <td>18</td> <td>11</td> <td>12</td> <td>3</td> <td>9</td> <td>14</td> <td>1</td> <td>3</td> <td>17</td> <td>9</td> <td>18</td> <td>2</td> <td>11</td> </tr> <tr> <td>3</td> <td>Without Manure (1846, and since)</td> <td>1</td> <td>12</td> <td>0</td> <td>12</td> <td>4</td> <td>17</td> <td>1</td> <td>19</td> <td>3</td> <td>12</td> <td>2</td> <td>4</td> <td>7</td> <td>17</td> <td>3</td> <td>3</td> <td>6</td> <td>8</td> <td>1</td> <td>17</td> </tr> <tr> <td>4</td> <td>{ 3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride { Sodium (common salt), 200 lbs. Sulphate Magnesia</td> <td>2</td> <td>2</td> <td>0</td> <td>14</td> <td>8</td> <td>13</td> <td>2</td> <td>8</td> <td>7</td> <td>10</td> <td>1</td> <td>15</td> <td>12</td> <td>10</td> <td>2</td> <td>19</td> <td>7</td> <td>7</td> <td>1</td> <td>14</td> </tr> <tr> <td>5</td> <td>3½ cwt. Superphosphate</td> <td>1</td> <td>18</td> <td>0</td> <td>14</td> <td>8</td> <td>5</td> <td>2</td> <td>9</td> <td>5</td> <td>0</td> <td>1</td> <td>16</td> <td>9</td> <td>13</td> <td>3</td> <td>5</td> <td>6</td> <td>11</td> <td>1</td> <td>12</td> </tr> <tr> <td>6</td> <td>3½ cwt. Superphosphate, 500 lbs. Sulphate Potass</td> <td>1</td> <td>15</td> <td>0</td> <td>13</td> <td>7</td> <td>16</td> <td>2</td> <td>7</td> <td>6</td> <td>9</td> <td>1</td> <td>12</td> <td>11</td> <td>11</td> <td>3</td> <td>5</td> <td>7</td> <td>17</td> <td>1</td> <td>13</td> </tr> <tr> <td>7</td> <td>3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)</td> <td>1</td> <td>18</td> <td>0</td> <td>14</td> <td>8</td> <td>2</td> <td>2</td> <td>6</td> <td>6</td> <td>7</td> <td>1</td> <td>14</td> <td>11</td> <td>2</td> <td>3</td> <td>6</td> <td>8</td> <td>4</td> <td>2</td> <td>0</td> </tr> <tr> <td>8</td> <td>Unmanured, 1853, and since; previously part Unman., part Superphos.</td> <td>1</td> <td>3</td> <td>0</td> <td>11</td> <td>5</td> <td>16</td> <td>2</td> <td>7</td> <td>3</td> <td>10</td> <td>1</td> <td>16</td> <td>9</td> <td>2</td> <td>3</td> <td>14</td> <td>6</td> <td>9</td> <td>2</td> <td>5</td> </tr> <tr> <td>9</td> <td>Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>9</td> <td>7</td> <td>2</td> <td>19</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> </tr> </tbody> </table>											Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.		Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	1	Farmyard Manure (14 tons)	6	3	1	15	9	8	2	9	12	6	3	11	13	16	3	15	10	14	2	12	2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..	6	13	1	16	11	11	2	18	11	12	3	9	14	1	3	17	9	18	2	11	3	Without Manure (1846, and since)	1	12	0	12	4	17	1	19	3	12	2	4	7	17	3	3	6	8	1	17	4	{ 3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride { Sodium (common salt), 200 lbs. Sulphate Magnesia	2	2	0	14	8	13	2	8	7	10	1	15	12	10	2	19	7	7	1	14	5	3½ cwt. Superphosphate	1	18	0	14	8	5	2	9	5	0	1	16	9	13	3	5	6	11	1	12	6	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass	1	15	0	13	7	16	2	7	6	9	1	12	11	11	3	5	7	17	1	13	7	3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)	1	18	0	14	8	2	2	6	6	7	1	14	11	2	3	6	8	4	2	0	8	Unmanured, 1853, and since; previously part Unman., part Superphos.	1	3	0	11	5	16	2	7	3	10	1	16	9	2	3	14	6	9	2	5	9	Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)	9	7	2	19
Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.																																																																																																																																																																																																																														
Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.																																																																																																																																																																																																																													
1	Farmyard Manure (14 tons)	6	3	1	15	9	8	2	9	12	6	3	11	13	16	3	15	10	14	2	12																																																																																																																																																																																																																			
2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..	6	13	1	16	11	11	2	18	11	12	3	9	14	1	3	17	9	18	2	11																																																																																																																																																																																																																			
3	Without Manure (1846, and since)	1	12	0	12	4	17	1	19	3	12	2	4	7	17	3	3	6	8	1	17																																																																																																																																																																																																																			
4	{ 3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride { Sodium (common salt), 200 lbs. Sulphate Magnesia	2	2	0	14	8	13	2	8	7	10	1	15	12	10	2	19	7	7	1	14																																																																																																																																																																																																																			
5	3½ cwt. Superphosphate	1	18	0	14	8	5	2	9	5	0	1	16	9	13	3	5	6	11	1	12																																																																																																																																																																																																																			
6	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass	1	15	0	13	7	16	2	7	6	9	1	12	11	11	3	5	7	17	1	13																																																																																																																																																																																																																			
7	3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)	1	18	0	14	8	2	2	6	6	7	1	14	11	2	3	6	8	4	2	0																																																																																																																																																																																																																			
8	Unmanured, 1853, and since; previously part Unman., part Superphos.	1	3	0	11	5	16	2	7	3	10	1	16	9	2	3	14	6	9	2	5																																																																																																																																																																																																																			
9	Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)	9	7	2	19																																																																																																																																																																																																																			
FIFTH SEASON, 1880. Seed dibbled, April 22-23 (Plot 9, April 24th). Crop taken up, Nov. 2-11.																																																																																																																																																																																																																																								
<table border="1"> <thead> <tr> <th colspan="2">Roots.</th> <th colspan="2">Leaves.</th> <th colspan="2">Roots.</th> <th colspan="2">Leaves.</th> <th colspan="2">Roots.</th> <th colspan="2">Leaves.</th> </tr> <tr> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> <th>Tons.</th> <th>cwts.</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>Farmyard Manure (14 tons)</td> <td>18</td> <td>11</td> <td>2</td> <td>14</td> <td>26</td> <td>8</td> <td>3</td> <td>5</td> <td>25</td> <td>4</td> <td>5</td> <td>10</td> <td>27</td> <td>3</td> <td>6</td> <td>1</td> <td>27</td> <td>5</td> <td>4</td> <td>1</td> </tr> <tr> <td>2</td> <td>Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..</td> <td>17</td> <td>8</td> <td>2</td> <td>0</td> <td>27</td> <td>16</td> <td>3</td> <td>14</td> <td>25</td> <td>15</td> <td>5</td> <td>10</td> <td>26</td> <td>0</td> <td>5</td> <td>12</td> <td>27</td> <td>9</td> <td>4</td> <td>3</td> </tr> <tr> <td>3</td> <td>Without Manure (1846, and since)</td> <td>4</td> <td>10</td> <td>0</td> <td>18</td> <td>14</td> <td>0</td> <td>2</td> <td>13</td> <td>9</td> <td>17</td> <td>2</td> <td>11</td> <td>11</td> <td>4</td> <td>3</td> <td>0</td> <td>12</td> <td>6</td> <td>2</td> <td>9</td> </tr> <tr> <td>4</td> <td>{ 3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride { Sodium (common salt), 200 lbs. Sulphate Magnesia</td> <td>5</td> <td>17</td> <td>0</td> <td>19</td> <td>23</td> <td>6</td> <td>3</td> <td>3</td> <td>19</td> <td>14</td> <td>2</td> <td>18</td> <td>30</td> <td>11</td> <td>5</td> <td>12</td> <td>24</td> <td>4</td> <td>3</td> <td>6</td> </tr> <tr> <td>5</td> <td>3½ cwt. Superphosphate</td> <td>5</td> <td>3</td> <td>0</td> <td>16</td> <td>18</td> <td>6</td> <td>2</td> <td>4</td> <td>9</td> <td>18</td> <td>2</td> <td>13</td> <td>12</td> <td>9</td> <td>2</td> <td>18</td> <td>14</td> <td>8</td> <td>2</td> <td>13</td> </tr> <tr> <td>6</td> <td>3½ cwt. Superphosphate, 500 lbs. Sulphate Potass</td> <td>4</td> <td>15</td> <td>0</td> <td>14</td> <td>21</td> <td>10</td> <td>2</td> <td>11</td> <td>18</td> <td>12</td> <td>3</td> <td>4</td> <td>27</td> <td>4</td> <td>5</td> <td>11</td> <td>21</td> <td>8</td> <td>2</td> <td>7</td> </tr> <tr> <td>7</td> <td>3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)</td> <td>7</td> <td>0</td> <td>0</td> <td>19</td> <td>21</td> <td>10</td> <td>2</td> <td>6</td> <td>19</td> <td>6</td> <td>2</td> <td>19</td> <td>26</td> <td>0</td> <td>5</td> <td>6</td> <td>23</td> <td>2</td> <td>2</td> <td>11</td> </tr> <tr> <td>8</td> <td>Unmanured, 1853, and since; previously part Unman., part Superphos.</td> <td>4</td> <td>0</td> <td>0</td> <td>17</td> <td>11</td> <td>14</td> <td>3</td> <td>5</td> <td>5</td> <td>19</td> <td>2</td> <td>17</td> <td>12</td> <td>4</td> <td>3</td> <td>1</td> <td>12</td> <td>1</td> <td>2</td> <td>15</td> </tr> <tr> <td>9</td> <td>Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>20</td> <td>19</td> <td>4</td> <td>0</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> <td>..</td> </tr> </tbody> </table>											Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.		Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	1	Farmyard Manure (14 tons)	18	11	2	14	26	8	3	5	25	4	5	10	27	3	6	1	27	5	4	1	2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..	17	8	2	0	27	16	3	14	25	15	5	10	26	0	5	12	27	9	4	3	3	Without Manure (1846, and since)	4	10	0	18	14	0	2	13	9	17	2	11	11	4	3	0	12	6	2	9	4	{ 3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride { Sodium (common salt), 200 lbs. Sulphate Magnesia	5	17	0	19	23	6	3	3	19	14	2	18	30	11	5	12	24	4	3	6	5	3½ cwt. Superphosphate	5	3	0	16	18	6	2	4	9	18	2	13	12	9	2	18	14	8	2	13	6	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass	4	15	0	14	21	10	2	11	18	12	3	4	27	4	5	11	21	8	2	7	7	3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)	7	0	0	19	21	10	2	6	19	6	2	19	26	0	5	6	23	2	2	11	8	Unmanured, 1853, and since; previously part Unman., part Superphos.	4	0	0	17	11	14	3	5	5	19	2	17	12	4	3	1	12	1	2	15	9	Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)	20	19	4	0
Roots.		Leaves.		Roots.		Leaves.		Roots.		Leaves.																																																																																																																																																																																																																														
Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.	Tons.	cwts.																																																																																																																																																																																																																													
1	Farmyard Manure (14 tons)	18	11	2	14	26	8	3	5	25	4	5	10	27	3	6	1	27	5	4	1																																																																																																																																																																																																																			
2	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?) ..	17	8	2	0	27	16	3	14	25	15	5	10	26	0	5	12	27	9	4	3																																																																																																																																																																																																																			
3	Without Manure (1846, and since)	4	10	0	18	14	0	2	13	9	17	2	11	11	4	3	0	12	6	2	9																																																																																																																																																																																																																			
4	{ 3½ cwt. Superphosphate, 500 lbs. Sulphate Potass, 200 lbs. Chloride { Sodium (common salt), 200 lbs. Sulphate Magnesia	5	17	0	19	23	6	3	3	19	14	2	18	30	11	5	12	24	4	3	6																																																																																																																																																																																																																			
5	3½ cwt. Superphosphate	5	3	0	16	18	6	2	4	9	18	2	13	12	9	2	18	14	8	2	13																																																																																																																																																																																																																			
6	3½ cwt. Superphosphate, 500 lbs. Sulphate Potass	4	15	0	14	21	10	2	11	18	12	3	4	27	4	5	11	21	8	2	7																																																																																																																																																																																																																			
7	3½ cwt. Superphos., 500 lbs. Sulphate Potass, 36½ lbs. Am.-salts (?)	7	0	0	19	21	10	2	6	19	6	2	19	26	0	5	6	23	2	2	11																																																																																																																																																																																																																			
8	Unmanured, 1853, and since; previously part Unman., part Superphos.	4	0	0	17	11	14	3	5	5	19	2	17	12	4	3	1	12	1	2	15																																																																																																																																																																																																																			
9	Farmyard Manure (14 tons), 3½ cwt. Superphosphate (?)	20	19	4	0																																																																																																																																																																																																																			

(1) "Superphosphate of Lime"—in all cases made from 200 lbs. Bone-ash, 150 lbs. Sulphuric acid, sp. gr.: 1.7 (and water).
 (2) "Ammonia-salts"—in each case equal parts Sulphate and Muriate of Ammonia of Commerce.
 (3) Plot 9 sown on the flat instead of on ridges; plants ridged up afterwards; rows 22 inches apart, plants 10 inches apart in the rows.

EXPERIMENTS ON POTATOS.—HOOS FIELD; commencing 1876.

The Land had been under experiments with Wheat, differently manured, from 1856 to 1874; and was fallowed in 1875.

Plots 1, 2, 3, and 4 had been unmanured for the Wheat. Plots 5 and 6 had received the same quantity of Ammonia-salts alone every year for the Wheat, as Plot 5 now receives for potatos; Plot 6 now receiving the same amount of nitrogen, but as Nitrate of Soda, instead of Ammonia-salts. Plots 7 and 8 received the same amount of complex mineral manure, and Ammonia-salts, for the Wheat, as Plot 7 now receives for potatos; and Plot 8 now receives the same complex mineral manures, and the same amount of nitrogen, but as Nitrate of Soda instead of Ammonia-salts. Plots 9 and 10 received the same complex mineral manures alone for the Wheat as Plot 10 now receives for potatos; Plot 9 now receives superphosphate only (?). Description of Potatos, in 1876, 1877, 1878, and 1879, the "Rock;" and in those years the rows were 25 inches apart; with 12 inches from plant to plant in the rows. In 1880 and 1881, the description was the "Champion;" and the rows were 25 inches apart; with 14 inches from plant to plant in the rows.

PLOTS.	MANURES PER ACRE PER ANNUM.	PRODUCE PER ACRE.				
		Tubers.				Tops.
		Good.	Small.	Diseased.	TOTAL.	
FIRST SEASON, 1876. Potatos planted, June 10-13; Crop taken up, Oct. 30-31.						
1	Unmanured	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Withered, not weighed, each lot spread on its own Plot and ploughed in.
2	Farmyard Manure (14 tons)	3 6½	0 5½	0 3½	3 17½	
3	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?)	3 18½	0 4	0 5½	4 5½	
4	Farmyard Manure (14 tons), 3½ cwt. Superphosphate, and 550 lbs. Nitrate of Soda	4 14½	0 6½	0 5½	5 6½	
5	400 lbs. Ammonia-salts (?)	5 9½	0 5½	0 19½	6 14½	
6	550 lbs. Nitrate of Soda	2 5½	0 6½	0 6	2 18	
7	400 lbs. Ammonia-salts, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	3 2	0 5½	0 9½	3 17½	
8	550 lbs. Nitrate of Soda, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	6 12½	0 9½	1 0	8 2	
9	3½ cwt. Superphosphate	6 17½	0 10	1 8½	8 15½	
10	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass., 100 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia	4 18½	0 8½	0 13½	6 1	
		5 3½	0 6½	0 13½	6 3½	
SECOND SEASON, 1877. Potatos planted, April, 27-28; Crop taken up, Oct. 8-10.						
1	Unmanured	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Withered, not weighed, each lot spread on its own Plot, (Oct. 14th) blew all off, before ploughing.
2	Farmyard Manure (14 tons)	2 11½	0 6½	0 2½	3 0½	
3	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?)	5 0½	0 11½	0 6	5 18	
4	Farmyard Manure (14 tons), 3½ cwt. Superphosphate, and 550 lbs. Nitrate of Soda	4 13½	0 7½	0 4	5 4½	
5	400 lbs. Ammonia-salts (?)	6 18½	0 7	0 17½	8 3½	
6	550 lbs. Nitrate of Soda	3 9½	0 7½	0 4	4 1	
7	400 lbs. Ammonia-salts, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	4 14½	0 6½	0 5½	5 7½	
8	550 lbs. Nitrate of Soda, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	6 12	0 11½	0 14½	7 17½	
9	3½ cwt. Superphosphate	7 8½	0 8½	0 16½	8 13½	
10	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass., 100 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia	2 12½	0 11½	0 1½	3 6	
		3 6½	0 7½	0 1½	3 15½	
THIRD SEASON, 1878. Potatos planted, April 29. Crop taken up, Sept. 18-21; Tops weighed, and spread on the Plots.						
1	Unmanured	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Withered, not weighed, each lot spread on its own Plot, (Oct. 14th) blew all off, before ploughing.
2	Farmyard Manure (14 tons)	2 6½	0 8½	0 2	2 17½	
3	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?)	4 11	0 12½	0 8½	5 11½	
4	Farmyard Manure (14 tons), 3½ cwt. Superphosphate, and 550 lbs. Nitrate of Soda	5 18½	0 14½	0 13½	7 6	
5	400 lbs. Ammonia-salts (?)	6 11½	0 11½	1 6½	8 9½	
6	550 lbs. Nitrate of Soda	2 16½	0 8½	0 5½	3 10½	
7	400 lbs. Ammonia-salts, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	3 16½	0 7	0 9½	4 13½	
8	550 lbs. Nitrate of Soda, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	7 6½	0 9½	1 1	8 17½	
9	3½ cwt. Superphosphate	7 11½	0 9	1 3½	9 4½	
10	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass., 100 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia	3 5½	0 9½	0 3½	3 18½	
		3 8	0 9	0 4½	4 1½	
FOURTH SEASON, 1879. Potatos planted, May 2; Crop taken up, Oct. 13-16.						
1	Unmanured	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Withered, not weighed, each lot spread on its own Plot and ploughed in.
2	Farmyard Manure (14 tons)	0 11½	0 4	0 0½	0 16½	
3	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?)	1 13½	0 4½	0 10½	2 8½	
4	Farmyard Manure (14 tons), 3½ cwt. Superphosphate, and 550 lbs. Nitrate of Soda	1 14	0 6	0 10½	2 10½	
5	400 lbs. Ammonia-salts (?)	2 16	0 5½	0 12½	3 14½	
6	550 lbs. Nitrate of Soda	0 17½	0 4	0 1½	1 3	
7	400 lbs. Ammonia-salts, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	0 14½	0 4½	0 2	1 0½	
8	550 lbs. Nitrate of Soda, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	2 4½	0 5	0 6	2 15½	
9	3½ cwt. Superphosphate	1 18½	0 4½	0 6½	2 9	
10	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass., 100 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia	0 17½	0 3½	0 1½	1 2	
		0 16½	0 3	0 1½	1 1½	
FIFTH SEASON, 1880. Potatos planted, April 13; Crop taken up, Plots 5 and 6, Sept. 9th; other Plots, Sept. 28-30.						
1	Unmanured	Tons. cwt.	Tons. cwt.	Tons. cwt.	Tons. cwt.	Withered, not weighed, each lot spread on its own Plot and ploughed in.
2	Farmyard Manure (14 tons)	0 14½	0 6½	0 0½	1 1½	
3	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (?)	4 13½	0 6	0 5	5 4½	
4	Farmyard Manure (14 tons), 3½ cwt. Superphosphate, and 550 lbs. Nitrate of Soda	5 6½	0 5½	0 10½	6 2½	
5	400 lbs. Ammonia-salts (?)	5 4	0 5½	1 1½	6 10½	
6	550 lbs. Nitrate of Soda	0 8½	0 9½	0 0	0 17½	
7	400 lbs. Ammonia-salts, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	0 11½	0 10	0 0	1 1½	
8	550 lbs. Nitrate of Soda, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	5 16½	0 5½	0 13	6 14	
9	3½ cwt. Superphosphate	6 3½	0 6½	1 1	7 11½	
10	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass., 100 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia	3 9	0 6½	0 3½	3 19	
		3 7½	0 6	0 3½	3 16½	

(1) "Superphosphate of Lime"—in all cases made from 200 lbs. Bone-ash, 150 lbs. Sulphuric acid, sp. gr. 1.7 (and water).

(2) "Ammonia-salts"—in each case equal parts Sulphate and Nitrate Ammonia of Commerce.

(3) The complex mineral manure having been sown in October, 1874, but the Wheat not put in, and therefore no crop taken in 1875, no mineral manures are sown afresh on Plots 7, 8, 9, and 10, for the first crop of potatos, 1876.

EXPERIMENTS ON POTATOS.—HOOS FIELD—continued.

SUMMARY OF THE COMPOSITION OF THE "GOOD" TUBERS.

As it will be some time before we shall be able to report fully the results obtained, or to be yet obtained, illustrating the influence of different manures, and of different seasons, on the composition of Potatos, an abstract of some of the analytical results at present at command is given below. The specific gravity of the tubers is also given. In the tubers the dry matter, nitrogen, and ash have been determined; and in some cases complete analyses of the ash have been made. Besides the results obtained relating to the composition of the tubers themselves, the dry matter, the sugar, the nitrogen, and the ash in the expressed juice have in many cases been determined; in some cases the amount of the nitrogen existing as albumenoids has been determined; and in some, complete analyses of the ash of the juice have been made. It may be remarked, that by far the larger proportion of both the mineral matter, and the nitrogen, is found to exist in the juice; and of the nitrogen in the juice, as a rule, not much more than half exists as albuminoids. In the majority of cases, the small potatoes have been submitted to the same methods of analysis as the good potatoes. And in a large number of cases, similar methods of examination have been applied to the still white, and also to the separated discoloured portions of the diseased potatoes. With regard to these latter results, it may be observed, that whilst the juice of the white portion of the diseased potatoes contained approximately the normal amount of nitrogen, that of the discoloured portion contained very much less. On the other hand, the washed, or exhausted "mark" of the white portion, contained very little nitrogen, whilst that of the discoloured portion contained very much more. The distribution of the mineral matter was much in the same order as that of the nitrogen. It was obvious that the juice had suffered exhaustion of much of both its nitrogen and its mineral matter, in the development of the fungus. There was an increased amount of sugar found in the diseased potatoes, the result of diseased action, and it probably also contributed to the development of the fungus. The results given in the Table relate to the "good" potatoes only. In interpreting the figures it must be borne in mind that in each year, the seed was planted on all the plots at the same time, and that all the crops were taken up at the same time; and as there was several times as much produce in some cases as in others, it is obvious that the crops would not each be at its best, and all in the same condition of maturity, when taken up. Then, again, the analyses were not performed immediately after taking up the crops, but after weighed samples had been kept in a cool place for some weeks or months; and in the following only preliminary statement of results, no correction is made for any change from the original weight of the samples, the results being calculated upon the fresh weights as finally taken for analysis.

Plots.	MANURES PER ACRE, PER ANNUM. (For Produce, see facing page.)	Specific Gravity of the Tubers.	Composition of the "Good" Tubers.				
			Dry Matter.	Mineral Matter (Ash).		Nitrogen.	
				In Fresh Tubers.	In Dry Matter.	In Fresh Tubers.	In Dry Matter.
FIRST SEASON, 1876.							
1	Unmanured	1.097	23.9	0.84	3.53	0.273	1.14
2	Farmyard Manure (14 tons)	1.091	23.4	0.96	4.11	0.226	0.97
3	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (1)	1.097	23.5	1.00	4.27	0.193	0.83
4	Farmyard Manure (14 tons), 3½ cwt. Superphosphate, and 550 lbs. Nitrate of Soda	1.085	21.2	0.83	3.92	0.299	1.41
5	400 lbs. Ammonia-salts (?)	1.087	22.1	0.81	3.67	0.337	1.52
6	550 lbs. Nitrate of Soda	1.091	22.0	0.79	3.59	0.332	1.51
7	400 lbs. Ammonia-salts, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	1.090	20.9	0.98	4.71	0.270	1.29
8	550 lbs. Nitrate of Soda, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	1.088	21.9	0.98	4.46	0.296	1.35
9	3½ cwt. Superphosphate	1.103	23.5	1.10	4.72	0.201	0.86
10	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass., 100 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia	1.102	22.9	1.06	4.64	0.173	0.76
SECOND SEASON, 1877.							
1	Unmanured	1.119	33.0	1.05	3.17	0.302	0.91
2	Farmyard Manure (14 tons)	1.109	26.5	1.06	4.00	0.212	0.80
3	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (1)	1.103	26.0	1.11	4.26	0.207	0.80
4	Farmyard Manure (14 tons), 3½ cwt. Superphosphate, and 550 lbs. Nitrate of Soda	1.112	27.2	1.06	3.90	0.301	1.11
5	400 lbs. Ammonia-salts (?)	1.107	22.0	0.67	3.07	0.281	1.28
6	550 lbs. Nitrate of Soda	1.116	25.9	0.74	2.85	0.301	1.16
7	400 lbs. Ammonia-salts, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	1.103	28.4	1.23	4.33	0.270	0.95
8	550 lbs. Nitrate of Soda, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	1.112	27.3	1.16	4.26	0.268	0.98
9	3½ cwt. Superphosphate	1.109	26.5	1.18	4.44	0.203	0.76
10	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass., 100 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia	1.109	26.8	1.21	4.52	0.208	0.78
THIRD SEASON, 1878.							
1	Unmanured	1.107	26.0	0.85	3.26	0.228	0.88
2	Farmyard Manure (14 tons)	1.100	24.4	1.02	4.20	0.209	0.86
3	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (1)	1.090	23.8	1.03	4.55	0.205	0.86
4	Farmyard Manure (14 tons), 3½ cwt. Superphosphate, and 550 lbs. Nitrate of Soda	1.078	21.9	0.97	4.45	0.269	1.23
5	400 lbs. Ammonia-salts (?)	1.099	24.9	0.78	3.12	0.310	1.25
6	550 lbs. Nitrate of Soda	1.105	25.5	0.67	2.64	0.326	1.28
7	400 lbs. Ammonia-salts, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	1.093	23.6	1.08	4.57	0.223	0.95
8	550 lbs. Nitrate of Soda, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	1.097	24.4	1.08	4.41	0.228	0.94
9	3½ cwt. Superphosphate	1.097	24.1	1.14	4.74	0.165	0.68
10	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass., 100 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia	1.098	23.7	1.16	4.90	0.167	0.71
FOURTH SEASON, 1879.							
1	Unmanured	1.103	24.3	0.96	3.95	0.242	1.00
2	Farmyard Manure (14 tons)	1.103	23.7	0.99	4.16	0.220	0.93
3	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (1)	1.099	24.0	1.02	4.26	0.218	0.91
4	Farmyard Manure (14 tons), 3½ cwt. Superphosphate, and 550 lbs. Nitrate of Soda	1.102	24.6	0.91	3.69	0.254	1.04
5	400 lbs. Ammonia-salts (?)	1.103	24.6	0.76	3.06	0.270	1.10
6	550 lbs. Nitrate of Soda	1.104	25.0	0.76	3.05	0.300	1.20
7	400 lbs. Ammonia-salts, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	1.098	23.1	0.95	4.13	0.241	1.05
8	550 lbs. Nitrate of Soda, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	1.102	23.9	1.04	4.36	0.272	1.14
9	3½ cwt. Superphosphate	1.099	23.6	1.10	4.65	0.219	0.93
10	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass., 100 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia	1.099	23.5	1.15	4.89	0.211	0.90
FIFTH SEASON, 1880.							
1	Unmanured	1.123	28.8	0.77	2.66	0.382	1.33
2	Farmyard Manure (14 tons)	1.114	27.6	0.98	3.56	0.287	1.04
3	Farmyard Manure (14 tons), and 3½ cwt. Superphosphate (1)	1.117	27.8	0.98	3.52	0.275	0.99
4	Farmyard Manure (14 tons), 3½ cwt. Superphosphate, and 550 lbs. Nitrate of Soda	1.102	25.2	0.88	3.48	0.357	1.41
5	400 lbs. Ammonia-salts (?)	1.114	28.5	0.84	2.95	0.430	1.51
6	550 lbs. Nitrate of Soda	1.117	28.8	0.88	3.06	0.415	1.44
7	400 lbs. Ammonia-salts, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	1.097	25.9	0.97	3.73	0.327	1.26
8	550 lbs. Nitrate of Soda, 3½ cwt. Superphos., 300 lbs. Sulph. Potass., 100 lbs. Sulph. Soda, 100 lbs. Sulph. Mag.	1.118	26.7	0.96	3.59	0.318	1.19
9	3½ cwt. Superphosphate	1.114	27.2	1.03	3.81	0.247	0.91
10	3½ cwt. Superphosphate, 300 lbs. Sulphate Potass., 100 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia	1.116	27.3	1.06	3.80	0.236	0.87

(1) "Superphosphate of Lime"—in all cases made from 200 lbs. Bone-ash, 150 lbs. Sulphuric acid, sp. gr. 1.7 (and water).
 (2) "Ammonia-salts"—in each case equal parts Sulphate and Muriate Ammonia of Commerce.

AGDELL FIELD.

EXPERIMENTS ON AN ACTUAL COURSE OF ROTATION—TURNIPS, BARLEY, LEGUMINOUS CROP (OR FALLOW), AND WHEAT. These Experiments were commenced in 1848; so that the present crop (1881) is the 34th experimental one, or the second crop of the Ninth Course. One-third of the land has been continuously unmanured; one-third manured with Superphosphate of Lime alone once every four years, that is for the turnip-crop commencing each course; and one-third manured (also for the turnip-crop only) with a complex manure, as described in the foot-note, No. 2. In the Second, Third, and Fourth Courses, clover was sown, but failed; and in them, and in the Fifth and Sixth Courses, beans were taken instead, on half of each plot, and the other half left fallow; for the third crop of the Seventh Course clover was again sown (spring 1873), on half of each plot, and gave three cuttings in 1874; the other half of each being left fallow. In the Eighth Course beans were again grown. For the Ninth Course (Spring 1881) clover has been sown. From half of each of the three plots the whole turnip-crop (roots and leaves) was removed; and on the other half the roots were eaten on the land by sheep, and the uneaten leaves spread and ploughed in. In the case of all the other crops, the total produce was removed from the land. The abstract of the results given below relates to the portions of each plot from which the turnip-crops were entirely removed; and on which, in the second, third, fourth, fifth, sixth, and eighth courses, beans (not fallow) replaced the clover. (Area under experiment, about 2½ acres.)

1 lb. (pound avoird.) per acre .. = (about) 1.12 Kilogramme per Hectare, or 0.57 Zollverein Pfund. per Prussian Morgen.
 1 cwt. (hundredweight) per acre = (about) 125.5 Kilogrammes per Hectare, or 0.64 Centner per Pr. Morgen.

Years.	Description of Crop.	PRODUCE PER ACRE.								
		Plot 1. Unmanured continuously.			Plot 2. Superphosphate of Lime, ¹ alone, for the Turnip Crops only.			Plot 3. Complex Manure, ² for the Turnip Crops only.		
		Corn ³ (or Roots).	Straw (or Leaf).	Total Produce. ⁴	Corn ³ (or Roots).	Straw (or Leaf).	Total Produce. ⁴	Corn ³ (or Roots).	Straw (or Leaf).	Total Produce. ⁴
1ST COURSE, 1848-51.										
1848	Norfolk White Turnips	63½ cwt.	45½ cwt.	111½ cwt.	23½ cwt.	106½ cwt.	32 cwt.	218 cwt.	151½ cwt.	369½ cwt.
1849	Barley	44½ bush.	293½ lbs.	565½ lbs.	29½ bush.	211½ lbs.	384½ lbs.	38½ bush.	208½ lbs.	374½ lbs.
1850	Clover (calcd. as hay)	5½ cwt.	57½ cwt.	53 cwt.
1851	Wheat	2½ bush.	343½ lbs.	539 lbs.	2½ bush.	337½ lbs.	533½ lbs.	2½ bush.	352½ lbs.	530 lbs.
2ND COURSE, 1852-55.										
1852	Swedish Turnips	26 cwt.	4½ cwt.	30½ cwt.	22¾ cwt.	90½ cwt.	243½ cwt.	39½ cwt.	36½ cwt.	433 cwt.
1853	Barley	34½ bush.	2450 lbs.	4465 lbs.	29½ bush.	1873 lbs.	5560 lbs.	38½ bush.	2604 lbs.	4873 lbs.
1854	Beans	5½ bush.	1655 lbs.	1445 lbs.	5½ bush.	1103 lbs.	1534 lbs.	9½ bush.	1353 lbs.	2065 lbs.
1855	Wheat	33½ bush.	3619 lbs.	5839 lbs.	33½ bush.	3395 lbs.	5789 lbs.	37½ bush.	3942 lbs.	6371 lbs.
3RD COURSE, 1856-59.										
1856	Swedish Turnips	32 cwt.	2½ cwt.	34½ cwt.	136 cwt.	7½ cwt.	147½ cwt.	333½ cwt.	12½ cwt.	346½ cwt.
1857	Barley	48½ bush.	2600 lbs.	4718 lbs.	337 lbs.	1475 lbs.	3076 lbs.	48 bush.	2435 lbs.	5165 lbs.
1858	Beans	6½ bush.	1100 lbs.	1515 lbs.	6½ bush.	1155 lbs.	1605 lbs.	12 bush.	1320 lbs.	2357 lbs.
1859	Wheat	35½ bush.	4030 lbs.	6262 lbs.	34½ bush.	3930 lbs.	6120 lbs.	39½ bush.	4610 lbs.	7154 lbs.
4TH COURSE, 1860-63.										
1860	Swedish Turnips	1 cwt.	(64 lbs.)	1 cwt.	2¾ cwt.	1½ cwt.	30½ cwt.	27½ cwt.	3½ cwt.	90½ cwt.
1861	Barley	38½ bush.	2522 lbs.	4718 lbs.	304 bush.	2000 lbs.	3775 lbs.	6½ bush.	3040 lbs.	7391 lbs.
1862	Beans	29 bush.	1840 lbs.	3661 lbs.	29 bush.	2150 lbs.	4640 lbs.	12 bush.	1320 lbs.	2357 lbs.
1863	Wheat	34½ bush.	3467 lbs.	5825 lbs.	34½ bush.	3390 lbs.	5619 lbs.	48½ bush.	3303 lbs.	5990 lbs.
5TH COURSE, 1864-67.										
1864	Swedish Turnips	8½ cwt.	0½ cwt.	9½ cwt.	68 cwt.	4½ cwt.	72½ cwt.	176½ cwt.	8½ cwt.	185 cwt.
1865	Barley	39 bush.	2154 lbs.	4182 lbs.	331 bush.	1618 lbs.	3394 lbs.	47½ bush.	2395 lbs.	5148 lbs.
1866	Beans	10½ bush.	1013 lbs.	1629 lbs.	7½ bush.	975 lbs.	1463 lbs.	20½ bush.	1990 lbs.	3143 lbs.
1867	Wheat	21 bush.	2143 lbs.	3473 lbs.	19½ bush.	1866 lbs.	3222 lbs.	23½ bush.	3003 lbs.	4567 lbs.
6TH COURSE, 1868-71.										
1868	Swedish Turnips	Failed, and ploughed up.			Failed, and ploughed up.			Failed, and ploughed up.		
1869	Barley	24½ bush.	1948 lbs.	3338 lbs.	284 bush.	2025 lbs.	3686 lbs.	42½ bush.	3309 lbs.	5800 lbs.
1870	Beans	19½ bush.	738 lbs.	1391 lbs.	15½ bush.	768 lbs.	1778 lbs.	24½ bush.	1656 lbs.	2664 lbs.
1871	Wheat	20½ bush.	2799 lbs.	4092 lbs.	23½ bush.	3098 lbs.	4521 lbs.	24 bush.	3440 lbs.	4942 lbs.
7TH COURSE, 1872-75.										
1872	Swedish Turnips	34½ cwt.	8½ cwt.	42½ cwt.	170½ cwt.	17½ cwt.	188 cwt.	339½ cwt.	33½ cwt.	375½ cwt.
1873	Barley	23½ bush.	1343 lbs.	2717 lbs.	20½ bush.	1555 lbs.	2875 lbs.	31½ bush.	1723 lbs.	3573 lbs.
1874	Clover	31½ cwt.	52½ cwt.	84½ cwt.
1875	Wheat	21½ bush.	2430 lbs.	3784 lbs.	28½ bush.	3536 lbs.	5328 lbs.	31½ bush.	4685 lbs.	6639 lbs.
8TH COURSE, 1876-79.										
1876	Swedish Turnips	17½ cwt.	5 cwt.	22½ cwt.	18¾ cwt.	284 cwt.	216½ cwt.	356 cwt.	55½ cwt.	411½ cwt.
1877	Barley	23½ bush.	1291 lbs.	2623 lbs.	24½ bush.	1174 lbs.	2538 lbs.	34½ bush.	1918 lbs.	3890 lbs.
1878	Beans	8½ bush.	740 lbs.	1301 lbs.	7½ bush.	1045 lbs.	1557 lbs.	20½ bush.	1655 lbs.	2963 lbs.
1879	Wheat	10½ bush.	1324 lbs.	1987 lbs.	14½ bush.	1771 lbs.	2729 lbs.	13 bush.	1658 lbs.	2493 lbs.
9TH COURSE, 1880-83.										
1880	Swedish Turnips	14 cwt.	2½ cwt.	16½ cwt.	199½ cwt.	11½ cwt.	211½ cwt.	439½ cwt.	43½ cwt.	482½ cwt.
1881	Barley
1882	Clover or Beans
1883	Wheat

SUMMARY—AVERAGE OF THE FIRST 8 COURSES, 1848-1879.

1848, '52, '56, '60, '64, '72, '78	} Swedish Turnips . .	26½ cwt.	9½ cwt.	35½ cwt.	148½ cwt.	26½ cwt.	175½ cwt.	272½ cwt.	43½ cwt.	315½ cwt.
1849, '53, '57, '61, '65, '69, '73, '77		} Barley	34½ bush.	2159 lbs.	4132 lbs.	284 bush.	1730 lbs.	3346 lbs.	41½ bush.	2577 lbs.
1850, '54, '58, '62, '66, '70, '74, '78	} Clover, 1850 and '74 } { (calcd. as hay) . . }	42½ cwt.	55 cwt.	73½ cwt.
1851, '55, '59, '63, '67, '71, '75, '79		} Beans	12½ bush.	1081 lbs.	1867 lbs.	12½ bush.	1200 lbs.	1996 lbs.	21½ bush.	1809 lbs.
	} Wheat	25½ bush.	2905 lbs.	4559 lbs.	27½ bush.	3067 lbs.	4823 lbs.	30½ bush.	3693 lbs.	5669 lbs.

(1) First Course—100 lbs. Bone-ash, and 100 lbs. Sulphuric Acid (sp. gr. 1.7); Second Course—160 lbs. Bone-ash, 120 lbs. Sulphuric Acid; Third, Fourth, Fifth, Sixth, Seventh, Eighth, and Ninth Courses—200 lbs. Bone-ash, and 150 lbs. Sulphuric Acid, per acre.
 (2) First Course—100 lbs. Pearl-ash, 100 lbs. Bone-ash, 100 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 1000 lbs. Rape-Cake; Second Course—300 lbs. Sulphate of Potass, 100 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 160 lbs. Bone-ash, 120 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake; Third, Fourth, Fifth, Sixth, Seventh, Eighth, and Ninth Courses—300 lbs. Sulphate of Potass, 200 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake, per acre.
 (3) The quantities given in Bushels represent the Dressed Corn only.
 (4) The "Total Produce" of the Corn-crops includes Dressed Corn, Offal Corn, Straw, and Chaff.

EXPERIMENTS WITH DIFFERENT DESCRIPTIONS OF WHEAT, IN 1881, AND SUMMARY OF RESULTS OBTAINED IN PREVIOUS YEARS.

Season 1881. RICKYARD FIELD, 1½ Cwt. Nitrate Soda, after Mangolds (with Dung and Guano) Carted off.	1871: Sawpit Field; 3 cwt. Guano; after Mangolds, carted off.	1872: Foster's Field; 2 cwt. Super- phosphate, 2 cwt. Nitrate Soda; after Hoops, carted off.	1873: Long Hoops Field; 1½ cwt. Nitrate; after Mangolds, (with Dung), carted off.	1874: Upton Harpenden Field; 2 cwt. Nitrate after Mangolds (with Dung), carted off.	1875: Little Knot- wood Field; 1½ cwt. Nitrate Soda; after Mangolds (with Dung), 1874, carted off.	1876: Harpenden Field; 2 cwt. Nitrate Soda; after Mangolds (with Dung), 1875, carted off.	1877: Sawpit Field; 1½ cwt. Nitrate Soda; after Mangolds (with Dung), 1876, carted off.	1878: Foster's Field; 2 cwt. Nitrate, after White Turnips (with Dung, as at Alden, 1877 cart. off. part carted off.	1879; () Little Knot- wood Field; 2 cwt. Nitrate, after Clover. First and second crops upwards; Feet.	1880; () Harpenden Field; 50 bushels of Soot; after Clover announced. Oats as at Hay; (Foster's wards Fed. 1880) carted off.	1882; Blackwood Field; 1½ cwt. Nitrate Soda; after Mangolds (with Dung, and Guano 1880) carted off.	() Averages, 1879 inclusive.	Nos.
1. White-chaff (Red)	82½	42½	38½	50½	40½	49½	48½	59	22½	28½	46½	48½	1
2. Rivet's (Red)	48½	48½	42½	49½	66½	42½	23½	51½	58½	2
3. Chubb Wheat (White)	28½	40	38½	35½	38½	40½	41½	55½	20½	14½	41½	51½	3
4. Red-chaff (White)	32½	37	48½	43½	34½	43½	41	89	4	
5. Brownick (Red)	33½	30½	38½	38½	39½	40½	40½	49½	24	19½	41½	41½	5
6. Red Wonder	31½	43½	37½	37½	33½	44½	39	52½	22	28½	39½	42½	6
7. Burwell (Old Red Lammas)	31½	41½	43½	39½	47½	38½	44½	46½	27	27	42½	43½	7
8. Bristol Red	29½	44½	39½	38½	31½	43½	44½	52½	21½	30½	42½	49½	8
9. Red Nunsey	34½	44½	27½	34½	39	37½	40½	47½	20½	27½	41½	39½	9
10. Red Langham	30½	43½	34½	34½	42½	42½	42½	50½	25½	28½	44½	41½	10
11. Woolly Ear (White)	31½	42½	37½	37½	36½	46½	37½	48½	46	21	41½	41½	11
12. Handcastle (White)	42	44½	42½	44	54	21½	24½	44½	44½	12
13. Golden Drop (Red), Hallett's	39½	40½	44½	44½	38½	48½	49½	52½	18½	18½	45½	45½	13
14. Victoria White, Hallett's	33½	41½	38½	38½	41½	43½	43½	43½	14½	15½	40½	40½	14
15. Hunter's White, Hallett's	29½	39½	43½	39½	28½	45½	40	42½	17½	22½	37½	37½	15
16. Original Red, Hallett's	30	35½	39½	39½	26	44½	44½	42½	37½	37½	16
17. White Chiddian	29½	38½	31½	31½	42	37½	42	49½	11½	11½	37½	37½	17
18. Red Rostock	37	49½	46½	46½	37½	46½	46½	47½	18½	28½	45½	45½	18
19. Casey's White	29½	42	37½	37½	39	46	43	47	15½	23	42	42	19
20. Golden Rough-chaff (Red)	33	39½	38½	38½	38½	38	39	46½	31½	31½	44	44	20
21. Hunter's White (Red)	33	42	43½	43½	44	44	43	52½	31	28	44	44	21
22. Club Wheat (Red)	36	45½	45½	45½	40	49½	49½	61	16	16	49	49	22
23. Main's Standing White	50	33½	33½	50	50	23
24. Main's Rough-chaff (White)	50	24	18	52	52	24
25. Belgian (White)	52½	21½	37	53	53	25
26. Webb's Challenge (White)	53	...	30	59	59	26
Means	51½	21½	29½	45½	45½	Means
1. White-chaff (Red)	60½	62½	59½	61½	60½	63½	60½	62	53½	56½	61½	61½	I
2. Rivet's (Red)	58½	61	63	60½	60½	51½	54½	57½	57½	2
3. Chubb Wheat (Red)	60½	61½	59½	59½	59½	62	60½	58½	49½	52	57	57	3
4. Red-chaff (White)	61	62½	60½	60½	60½	62½	60½	61½	53	53	60½	60½	4
5. Brownick (Red)	60	61½	59½	59½	59½	60½	60½	62	52½	54	61	61	5
6. Red Wonder	59	60½	60	60	60½	61½	61½	64	52½	52	61	61	6
7. Burwell (Old Red Lammas)	62	63	61½	61½	61½	64	61½	64	55	55	62	62	7
8. Bristol Red	60	61	60½	60½	60½	62	60½	63	54	54	61	61	8
9. Red Nunsey	63	65	62	62	60	66	58	62	57½	57	63	63	9
10. Red Langham	60	61	60½	60½	60	62	61½	63	54½	56	61	61	10
11. Woolly Ear (White)	61	61	61	61	61	63	61	63	54	54	61	61	11
12. Handcastle (White)	61	62	60	60	60	63	61	63	52	52	61	61	12
13. Golden Drop (Red), Hallett's	61	63	62	62	61	64	61	64	52	52	62	62	13
14. Victoria White, Hallett's	59	61	59½	59½	59	63	61	63	52	52	62	62	14
15. Hunter's White, Hallett's	58	60	58½	58½	58	63	61	63	51	51	61	61	15
16. Original Red, Hallett's	62	63	61½	61½	61	64	61	62	55	55	62	62	16
17. White Chiddian	60	61	60½	60½	60	63	61	64	55	55	62	62	17
18. Red Rostock	62	63	61½	61½	61½	64	61	65	55	55	62	62	18
19. Casey's White	60	61	60½	60½	60	64	61	65	54	54	62	62	19
20. Golden Rough-chaff (Red)	61	62	60½	60½	60	64	61	65	54	54	62	62	20
21. Bolet's Prolific (Red)	61	62	60½	60½	60	65	61	65	54	54	62	62	21
22. Club Wheat (Red)	60	61	60	60	61	65	61	65	52	52	61	61	22
23. Main's Standing White	62	52	52	61	61	23
24. Main's Rough-chaff (White)	61	56	56	61	61	24
25. Belgian (White)	60½	51	51	61	61	25
26. Webb's Challenge (White)	62	51	51	61	61	26
Means	62	53	56	61	61	Means

(*) All the crops were more or less affected by wire-worm, large bare patches appearing on many plots; and much grain was immature and blighted.

(†) Owing doubtless in great part to the imperfect development of the crop of 1880 did not germinate at all, and of that which did come up a great deal was afterwards destroyed by wire-worm, so that up to the end of March it was a question whether there would be a plant left in the field at all.

(‡) With the exception of the crops of 1879 and 1880, the seed for which was obtained direct from the seedsmen, not grown on the farm, as were the others. The white wheats appeared to suffer most either from imperfectly developed seed, wire-worm, or blight. The most satisfactory crop was "Webb's Challenge," the seed for which was obtained direct from the seedsmen, not grown on the farm, as were the others.

ROTHAMSTED

MAY

SUMMARY STATEMENT OF THE PRESENT AND PREVIOUS

(13 Years, 1869-1881,

PREVIOUS CROPPING

Name of Field.	Acres.	1869.	1870.	1871.	1872.	1873.	1874.	1875.
Thirty Acres	30	Wheat, 2 cwt. Guano.	Oats, 2 cwt. Guano.	Barley, 2 cwt. superphos., 2 cwt. Nitrate Soda.	Barley, 2½ cwt. superphos., 2½ cwt. Nitr. Soda, (2½ acres experiment).	Barley (½ with Grass-seeds). 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Grass (½), Folded, and 1 cwt. Nitrate. Barley (¾), 2 cwt. superphosphate, 2½ cwt. Nitrate Soda.	Grass (¾), Sheep-folded. Tares (½) Dung.
Harpenden	22	Swedes, Dung and various Artificial Manures.	Wheat, 3 cwt. Guano.	Oats, 3 cwt. Guano, 1 cwt. Nitrate Soda. Tares, Dung.	Oats, 2½ cwt. superphos., 2½ cwt. Nitr. Soda, Tares, Dung.	Barley, After Oats—2 cwt. super- phosphate; 2 cwt. Nitrate. After Tares—1 cwt. super- phosphate; 1 cwt. Nitrate.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Mangolds, Dung, and 2 cwt. Guano. (Carted off.)
Little Hoos	9	Barley, 1 cwt. dried Blood, ½ cwt. Sulph. Ammonia, 1 cwt. superphosphate.	Barley, 2½ cwt. Guano.	Barley, 3 cwt. superphos., 2½ cwt. Nitrate Soda.	Barley (with Clover). 2½ cwt. superphos., 2½ cwt. Nitr. Soda.	Barley (¾), Unmanured. Clover (¼), Unmanured.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda (1 acre Unmanured).	Barley, where Barley 1873, 2 cwt. superphosphate, 2 cwt. Nitrate of Soda, where Clover 1873, Half quantities.
Fosters'	18	Barley, 1 cwt. dried Blood, ½ cwt. Sulph. Ammonia, 1 cwt. superphosphate.	Oats, 2 cwt. Guano, 3 cwt. Blood Manure.	Roots, Tares, and Rape, Dung and Artificial.	Wheat, Varieties of Wheat, 2 cwt. superphos., 2 cwt. Nitr. Soda, ¾ Sheep-folded.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda, (2 acres experiment).	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Barley, (1) 3½ cwt. Guano, (2) 2½ cwt. superphosphate, 2½ cwt. Nitrate Soda, (3) 1½ cwt. Guano, 1½ Nitrate.
Knott Wood	30	Wheat, 3 cwt. Guano (one-half), Unmanured (one-half), after Swedes ploughed up and Followed.	Oats, 3 cwt. Guano.	Oats, 3 cwt. Guano, 1 cwt. Nitrate Soda.	Oats, 2½ cwt. superphos., 2½ cwt. Nitr. Soda.	Tares (¾), Dung. Swedes (¾), Dung, 2 cwt. superphosph.; 2 cwt. Nitrate Soda.	Barley, After Roots and Tares carted, 2 cwt. superphosphate, 2 cwt. Nitrate Soda, After Tares fed, 1 cwt. each.	Barley, 2½ cwt. superphosphate, 2½ cwt. Nitrate Soda.
Little Knott Wood	14	Mangolds, 12 tons Dung, 3 cwt. Guano.	Wheat, 3 cwt. Guano.	Oats, 3 cwt. Guano, 1 cwt. Nitrate Soda.	Oats, ¾ Sheep-folded. All, 2½ cwt. super, 2½ cwt. Nitr. Soda.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Mangolds, Dung, (Carted off.)	Wheat (Varieties), 1½ cwt. Nitrate Soda.
Sawpit	14	Wheat, 3 cwt. Guano.	Mangolds, Dung and 3 cwt. Guano.	Wheat, 3 cwt. Guano.	Oats, 2½ cwt. superphos., 2½ cwt. Nitr. Soda.	Oats, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Barley, 2½ cwt. superphosphate, 2½ cwt. Nitrate Soda.
Rick-yard	8	Tares, Dung.	Barley, 1 cwt. Guano.	Mangolds, Dung and 4 cwt. Cotton Cake.	Wheat, Unmanured.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Tares, Dung, ½ followed by Turnips, 1 cwt. superphosphate, 1 cwt. Nitrate Soda.	Barley, 1 cwt. Nitrate Soda.
Six Acres	6	Wheat, 2 cwt. Guano, 1 cwt. Nitrate of Soda.	Barley, 2½ cwt. Guano.	Barley, 3 cwt. superphos., 2½ cwt. Nitrate Soda.	Barley, 2½ cwt. superphos., 2½ cwt. Nitr. Soda.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Barley, 2 cwt. superphosphate, 2½ cwt. Nitrate Soda.	Barley, 2 cwt. superphosphate, 2½ cwt. Nitrate Soda.
Clay-Croft	12	Oats, 2 cwt. Guano, 1 cwt. dried Blood, ½ cwt. Sulph. Ammonia.	Turnips, Dung and 3 cwt. super- phosphate.	Wheat, Unmanured.	Oats, 2½ cwt. superphos., 2½ cwt. Nitr. Soda.	Clover, Unmanured.	Wheat, 2 cwt. Nitrate Soda.	Oats, 2½ cwt. superphosphate, 2½ cwt. Nitrate Soda.
Ten Acres	10	Wheat, 2 cwt. Guano.	Oats, 3 cwt. Guano.	Mangolds, Dung and 4 cwt. Cotton Cake.	Wheat, Unmanured.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda (5 acres experiment).	Oats, 2 cwt. superphosphate, 2½ cwt. Nitrate Soda.	Oats, 2½ cwt. superphosphate, 2½ cwt. Nitrate Soda.
Agdell	9	Barley, Unmanured.	Barley, 1½ cwt. Guano, 1½ cwt. super- phosphate.	Mangolds, Dung and 4 cwt. Cotton Cake.	Wheat, Unmanured (and part Roots).	Clover, Unmanured. Barley, Experiment.	Wheat, 1 cwt. Nitrate Soda (3 acres Experiment, ½ Clover, ½ Fallow).	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda. Wheat, 3 acres, Experiment.
Long Hoos	25	Oats, 2 cwt. Guano, 1 cwt. dried Blood, ½ cwt. Sulph. Ammonia.	Sainfoin, Unmanured.	Sainfoin, Unmanured, (Steam cultivated, July.)	Mangolds, Dung, (Carted off.)	Wheat, (½ Varieties of Wheat), 1½ cwt. Nitrate Soda.	Oats, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Oats, 2½ cwt. superphosphate, 2½ cwt. Nitrate Soda.
Sawyers'	25	Fallow.	Wheat, 4 cwt. Guano.	Wheat, 4 cwt. Guano, 1 cwt. Nitrate Soda.	Barley, 2½ cwt. superphos., 2½ cwt. Nitr. Soda.	Oats, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Mangolds and Swedes, Dung.	Barley after Swedes (¾) 2 cwt. superphosphate, 2 cwt. Nitrate Soda. Wheat after Mangolds (¾) 1½ cwt. Nitrate Soda.
West Barn	30	Wheat, 3 cwt. Guano.	Sainfoin, Unmanured.	Sainfoin, Unmanured.	Sainfoin, Unmanured.	Oats, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Wheat (Oats fed off 1873), 1½ cwt. Nitrate Soda.	Oats, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.

FARM.

1881.

CROPPING, &c., OF THE ARABLE LAND NOT UNDER EXPERIMENT.
inclusive.)

AND MANURING.					Crops, &c., Present Season, 1880-81.	Acres.	Name of Field.
1876.	1877.	1878.	1879.	1880.			
Grass (3/4), Compost. Wheat (1/2), 1 cwt. Nitrate Soda.	Grass (3/4), Cattle Grazed. Barley (1/2), 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.	Grass (3/4), Cattle Grazed with Cotton-Cake. Tares (1/2), Dung.	Grass (3/4), Cattle Grazed with Cotton-Cake. Barley (1/2), 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Grass (3/4), Cattle Grazed with Cotton-Cake. Wheat (1/2), (Varieties), 50 bushels Soot. Mangolds (1/2), 15 tons Dung & 3 cwt. Guano. (Carted off).	Grass (3/4), Cattle Grazed with Cotton-Cake, Hay, and Mangolds. Oats (1/2), 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.	30	Thirty Acres
Wheat (Varieties), 2 cwt. Nitrate Soda.	Barley, 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.	Barley (with Clover), 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.	Clover, Unmanured. One Crop as Hay.	Fallow (1/2), Wheat (1/2), 1 1/2 cwt. Nitrate Soda.	22		Harpenden.
Barley, 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda (1/2 with Clover).	Barley, 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda (1/2 with Clover).	Barley (1/2), 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda. Clover (1/2), Unmanured. Two Crops as Hay.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Barley, 2 1/2 cwt. Guano.		Barley, 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda. (1/2 with Clover).	9
Barley, 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.	White Turnips, Dung, Superphosphate. 1/2 cwt. Nitrate Soda; part fed, part carted.	Wheat (Varieties), 2 cwt. Nitrate Soda.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Barley, 2 1/2 cwt. Guano.	Fallow.	18	
Oats, 2 1/2 cwt. superphosphate, 3 cwt. Nitrate Soda.	Barley, 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.	Roots (1/2), Dung and Artificial. (Carted off). Fallow (1/2).	Wheat (1/2), 2 cwt. Nitrate Soda. Barley (1/2), 2 cwt. superphosphate, 2 cwt. Nitrate Soda (all with Clover).	Barley (1/2), 2 1/2 cwt. Guano. Clover (1/2), Unmanured. (1/2) 2 crops as Hay, (1/2) 1 crop as Hay, aftwda. Fed.	Wheat, 1 1/2 cwt. Nitrate Soda.		30
Oats, 2 1/2 cwt. superphosphate, 3 cwt. Nitrate Soda.	Oats (with Clover), 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.	Clover, Unmanured. First and second Crops as Hay; afterwards fed.	Wheat (Varieties), 2 cwt. Nitrate Soda.	Barley, 2 1/2 cwt. Guano.	Oats, 2 1/2 cwt. Nitrate Soda, 2 1/2 cwt. superphosphate.	14	
Mangolds, 25 tons Dung. (Carted off).	Wheat (Varieties), 1 1/2 cwt. Nitrate Soda.	Barley, 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Fallow.	Mangolds, (1/2) Dung 16 tons in 1880, 4 cwt. Guano in 1881, (1/2) Dung, 18 tons in 1881.		14
Swedes, Dung, and Superphosphate.	Barley, 1 cwt. Nitrate Soda.	Barley, 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.	Barley (1/2), 2 cwt. superphosphate, 2 cwt. Nitrate Soda. Fallow (1/2).	Mangolds, 15 tons Dung, 3 cwt. Guano. (Carted off.) Mangolds, 15 tons Dung, 3 cwt. Guano. (Carted off.)	Wheat (Varieties), 1 1/2 cwt. Nitrate Soda.	8	
Barley, 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.	Barley (with Clover), 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.	Clover, Unmanured. Two Crops as Hay.	Wheat, 2 cwt. Nitrate Soda.	Barley, 2 1/2 cwt. Guano.	Wheat, 1 1/2 cwt. Nitrate Soda.		6
Oats, 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.	Fallow.	Wheat, 2 cwt. Nitrate Soda.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Barley, 2 1/2 cwt. Guano.	Fallow.	12	
Fallow.	Wheat (with Clover), 2 cwt. Nitrate Soda.	Clover, Unmanured. Two Crops as Hay.	Barley, 2 cwt. Nitrate Soda (with Grass Seeds).	Grass, Unmanured.	Grass, Cattle Grazed, with Cotton-Cake. Dung 15 tons.		10
Barley, 2 1/2 cwt. superphosphate, 3 cwt. Nitrate Soda. Swedes, 3 acres Experiment.	Barley, 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda. Barley, 3 acres Experiment.	Potatos, Dung and Artificial. (3 acres Experiment 1/2 Beans, 1/2 Fallow.)	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda (3 acres Experiment, Wheat).	Fallow, (3 acres Experiment, Swedes).	Oats, 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda, (3 acres Experiment, Barley).	9	
Oats (1/2), 2 1/2 cwt. superphosphate, 3 cwt. Nitrate Soda. Tares (1/2), Dung.	Barley, 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.	Barley, 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.	Barley, 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Barley, 2 1/2 cwt. Guano.	Barley (with Clover), 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.		25
Barley (with Clover), 2 cwt. superphosphate, 2 cwt. Nitrate Soda.	Barley (1/2), 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda. Tares (1/2), Dung.	Barley, (1/2) 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda, (1/2) 2 1/2 cwt. Nit. Soda alone.	Roots (1/2), 25 tons Dung, 1 cwt. Nitrate Soda (Carted off); Fallow (1/2).	Wheat, 50 bushels Soot.	Mangolds, Dung 20 tons.	25	
Oats, 2 cwt. superphosphate, (1) 1 1/2 Nitrate Soda, (1) 2 1/2 Nitrate Soda.	Fallow.	Wheat, 2 cwt. Nitrate Soda.	Winter Oats, 2 cwt. Nitrate Soda.	Barley, 50 bushels Soot.	Barley, 2 1/2 cwt. superphosphate, 2 1/2 cwt. Nitrate Soda.		30