Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



## Memoranda of the Field Experiments at Rothamsted: May 1881



Full Table of Content

## **Experiments on Oats; Geescroft Field**

## **Rothamsted Research**

Rothamsted Research (1882) *Experiments on Oats; Geescroft Field ;* Memoranda Of The Field Experiments At Rothamsted: May 1881, pp 12 - 12 - **DOI:** 

https://doi.org/10.23637/ERADOC-1-245

( 12 )

## GEESCROFT FIELD.

EXPERIMENTS ON THE GROWTH OF OATS YEAR AFTER YEAR ON THE SAME LAND; WITHOUT MANDER, AND WITH DIFFERENT KINDS OF MANDER.

Previous Cropping—1847 and 1848, Clover, Experimental Manures; 1849—1859, Beans, Experimental Manures; 1860, Fallow; 1861 and 1862, Wheat, Unmanured; 1866, Beans, Unmanured; 1867 and 1868, Wheat, Unmanured. The first Experimental Oat Crop was in 1869; the last in 1878, since which time, owing to the wetness and the foulness of the land, it has been left fallow.

(Area under Experiment, 4 acre.)

Produce per Acre,	AVERAGE PER ANNUM 5 YEARS, 1869-1873.	Total Straw.		cwts. 103	133	283	411	273	35	
		Dressed Corn.	Weight per Bushel.	1bs. 832	 	357	37	353	352	
			Quantity.	Bushels.	24 +48	47	29	471	573	H
	5тн Ѕвавом, 1873.	Total Straw.		cwts.	1000	162	273	164	24	
		Dressed Corn.	Weight per Bushel.	1bs. 271	12 88 88	325	343	303	83 83 83	A 10
			Quantity.	Bushels.	17	364	484	393	638	TOTAL
	4TH SEASON, 1872.	Total Straw.		cwts.	103	308	453	208	24	PA MOM
		Dressed Corn.	Weight per Bushel.	lbs. 364	373	373	393	365	374	0 4
			Quantity.	Bushels.	193	553	623	421	448	ONI.V H
	3RD SEASON, 1871.	Total Straw.		cwts.	131	403	50	343	488 8	S. a.
		Corn.	Weight per Bushel,	1bs. 333	354	363	353	363	60 64	THE ATTE
		Dressed Corn.	Quantity.	Bushels.	22	57 <sub>3</sub>	5888	55	₹09	N dwa k
	2nd Season, 1870.	Total Straw.		cwta.	938	174	288	23	288	TTA-SAT.TS
		Dressed Corn.	Weight per Bushel,	1bs.	351	347	36	353	35.4	AMMON
			Quantity.	Bushels.	191	80	508	361	20	впеове
	869.	Total Straw.		cwts. 194	243	367	54	423	497	RE BE A B
,	1sr Season, 1869.	Dressed Corn.	Weight per Bushel.	lbs. 363	583	373	\$68	383	188	T. MAN
			Quantity.	Bushels.	45	563	15\$	€24	869	WINER
MANURES, PER ACRE, PER ANNUM.			Unmanured	(200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ owts. Superphosphate of Lime 00	400 lbs. Ammonia-salts (2)	(400 lbs. Ammonia-salts, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ cwts. Superphosphate	550 lbs. Nitrate of Soda (3)	(550 lbs. Nitrate of Soda, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3‡ owts. Superphosphate	SECOND 5 VEARS: WINERAL WANTERS AS REFORE AMMONIA-SAIMS AND NITHEAR OF SONA ONIX HAIR AS WHICH AS DEPARTMENTS	
	Prots.			1	61	တ	4	10	9	

6, and 8	cwts.	63	141	50	111	14}
AVERAGE FER ANNUM YEARS, 1874, 5, 6, and	1bs. 314	318	\$88	853	318	348
AYERAGE FER ANNUM 4 YEARS, 1874, 5, 6, and 8.	Bushels.	131	281	38	263	283
1878.	cwts.	144	123	223	123	173
10TH SEASON, 1878.	1bs.	354	32₹	37	341	361
10тн	Bushels.	172	30	458	341	87
77 (%).	cwts.	:	:	:		
9TH SEASON, 1877 (*). FALLOW.	lbs.	1	;	:	:	:
9TH SE.	Bushels.	:			;	;
76 (*).	cwts.	23	9	124	38	00
8TH SEASON, 1876 (*).	1bs. 32	30	343	353	303	83.
STH SE	Bushels.	75	178	293	123	198
1875.	cwts. 5g	63	153	\$0 <b>\$</b>	113 (+)	33 (4) 14 (4)
7TH SEASON, 1875.	1bs.	298	327	342	314 (*)	33% (4)
7TH 8	Bushels.	181	303	30\$	234 (4)	28§ (4)
1874.	cwts.	₹9	223	245	16½ (1)	16§ (4)
6тн Ѕвавои, 1874.	1bs.	814	331	348	30 (4)	33½ (4)
6тн 8	Bushels.	138	874	463	351 (4)	284 (4) 332 (4) 168 (4)
	Unmanured	(200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda,) 100 lbs. Sulphate Magnesia, and 3½ owts.) Superphosphate of Lime (¹)	200 lbs. Ammonia-salts (2)	(2001bs. Ammonia-salts, 2001bs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 3½ cwts. Superphosphate	275 lbs. Nitrate of Soda (3)	275 lbs. Nitrate of Soda, 200 lbs. Sulphate Potass, 100 lbs. Sulphate Soda, 100 lbs. Sulphate Magnesia, and 34 owts. Superphosphate
	1	63	8	4	10	9

o contain the same amount of Nitrogen as 400 lbs, "Ammonia-salts," of Nitrate of Soda had been applied year after year, the land, though more worked, was so wet that it could not be got into favourable condition for sowing, and the plant was very irregular, of Nitrate of Soda had been applied year after year, the land, though more worked, was so wet that it could not be got into favourable condition for sowing, and the plant was very irregular. (1) "Superphosphate of Lime"—in all cases, made from 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid sp. gr. 1.7 (and water).
(2) A Ammoini-sells "—in each case, equal parts Shiphate and Muritee of Ammonia of Commerce.
(3) 550 lbs. Nitrate of Soda is redecaned to contain the same amount of Nitrogen as 400 lbs. "Ammonia-salts."
(4) On these piots, where large quantities of Nitrate of Soda had been applied year after year, the land, though more worked, was so wet that it could not be got if on the september of the land, especially on the Nitrate piots, it was not sown until April 6, and than with a very unfavourable seed be up very irregularly, and much of the paished from standing surface-water.
(2) Owing to the very wet winter, 1876-7, the land could not be worked in time for sowing, and was therefore left fallow in 1877; no manures being applied.