Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Memoranda of the Field Experiments at Rothamsted: May 1881



Full Table of Content

Experiments on Rotation; Agdell Field

Rothamsted Research

Rothamsted Research (1882) *Experiments on Rotation; Agdell Field;* Memoranda Of The Field Experiments At Rothamsted: May 1881, pp 24 - 24 **- DOI:**

https://doi.org/10.23637/ERADOC-1-245

(24)

AGDELL FIELD.

EXPERIMENTS ON AN ACTUAL COURSE OF ROTATION—TURNIPS, BARLEY, LEGUMINOUS CROP (OR FALLOW), AND WHEAT.

Experiments on an actual Course of Rotation—Turnips, Barley, Leguminous Crop (or Fallow), and Wheat.

These Experiments were commenced in 1848; so that the present crop (1881) is the 34th experimental one, or the second crop of the Ninth Course. One-third of the land has been continuously unmanured; one-third manured with Superphosphate of Lime alone once every four years, that is for the turnip-crop commencing each course; and one-third manured (also for the turnip-crop only) with a complex manure, as described in the foot-note, No. 2.

In the Second, Third, and Fourth Courses, clover was sown, but failed; and in them, and in the Fifth and Sixth Courses, beans were taken instead, on half of each plot, and the other half left fallow; for the third crop of the Seventh Course clover was again sown (spring 1873), on half of each plot, and gave three cuttings in 1874; the other half of each being left fallow. In the Eighth Course beans were again grown. For the Ninth Course (Spring 1881) clover has been sown.

From half of each of the three plots the whole turnip-crop (roots and leaves) was removed; and on the other half the roots were eaten on the land by sheep, and the uneaten leaves spread and ploughed in. In the case of all the other crops, the total produce was removed from the land. The abstract of the results given below relates to the portions of each plot from which the turnip-crops were entirely removed; and on which, in the second, third, fourth, fifth, sixth, and eighth courses, beans (not fallow) replaced the clover.

	1 lb. (pound avoir, 1 cwt. (hundredwe) per acre ight) per acre	= (about) = (about)) 1:12 Kilog) 125:5 Kilog	ramme per He rammes per H	ctare, or 0.5 ectare, or 0.6	7 Zollverein Pfi 4 Centner per I	ind, per <mark>Pru</mark> ssi Pr. Morgen.	an Morgen.	1 12
					1	RODUCE PER AC	PRE.			
Years.	Description of Grop.	PLOT 1. Unmanured continuously.			PLOT 2. Superphosphate of Lime, l alone, for the Turnip Crops only.			PLOT 3. Complex Manure,2 for the Turnip Crops only.		
		Corn 3 (or Roots).	Straw (or Leaf).	Total Produce,+	Corn ³ (or Roots),	Straw (or Leaf),	Total Produce.4	Corn 3 (or Roots).	Straw (or Leaf).	Total Produce,
				1st Cou	RSE, 1848-5		1			
1848 1849 1850 1851	Norfolk White Turnips Barley. Clover (calcd, as hay) . Wheat.	65½ cwts. 44½ bush. 28½ bush.	454 cwts. 2983 lbs. 3431 lbs.	111‡ cwts. 5656 lbs. 54 cwts. 5389 lbs.	225% cwts. 29% bush. 28 bush.	106½ cwts. 2111 lbs. 3371 lbs.	332 cwts. 3841 lbs. 574 cwts. 5253 lbs.	218 cwts. 28½ bush. 28½ bush.	1512 cwts. 2088 lbs. 3552 lbs.	3694 cv 3794 lb 63 cv 5500 lb
				2nd Cou	rse, 1852-5	Ď.			THE	
1852 1853 1854 1855	Swedish Turnips. Barley Beans. Wheat	26 cwts. 34% bush. 5% bush. 35% bush.	4½ cwts. 2430 lbs. 1955 lbs. 3619 lbs.	30½ cwts. 4465 lbs. 1445 lbs. 5859 lbs.	2234 cwts. 284 bush. 5% bush. 354 bush.	204 cwts. 1873 lbs. 1103 lbs. 3525 lbs.	243‡ cwts. 3560 lbs. 1534 lbs. 5789 lbs.	396‡ cwts. 38‡ bush. 9½ bush. 37‡ bush.	36½ cwts. 2604 lbs. 1355 lbs. 3942 lbs.	433 cw 4873 lbs 2065 lbs 6371 lbs
				3rd Cou	rse, 1856-5	9.			-	***
1856 1857 1858 1859	Swedish Turnips, Barley Beaus, Wheat	32 cwts. 48‡ bush. 6‡ bush. 35‡ bush.	2½ cwts. 2600 lbs. 1100 lbs. 4030 lbs.	34‡ cwts. 5337 lbs. 1515 lbs. 6262 lbs.	136 cwts. 28½ bush. 6½ bush. 34½ bush.	7½ cwts. 1475 lbs. 1155 lbs. 3930 lbs.	142½ cwts. 3076 lbs. 1605 lbs. 6120 lbs.	3334 cwts. 48 bush. 12% bush. 39% bush.	12½ cwts. 2435 lbs. 1520 lbs. 4610 lbs.	3464 cwts 5168 lbs. 2357 lbs. 7154 lbs.
				4тн Соп	rse, 1860-6	3.			7,00	
1860 1961 1862 1863	Swedish Turnips Barley	1 cwt. 38§ bush. 29 bush. 34‡ bush.	(64 lbs.) 2522 lbs. 1840 lbs. 3467 lbs.	I cwt. 4718 lbs. 3661 lbs. 5625 lbs.	294 cwts. 308 bush. 295 bush. 346 bush.	1½ cwt. 2000 lbs. 2150 lbs. 3590 lbs.	30% cwts. 3775 lbs. 4040 lbs. 5619 lbs.	87½ cwts. 60% bush. 43% bush. 46% bush.	34 cwts. 3940 lbs. 3280 lbs. 4697 lbs.	904 cwt 7391 lbs. 5990 lbs. 7626 lbs.
				5TH Cou	rse, 1864-6	7.				
1864 1865 1866 1867	Swedish Turnips. Barley. Beans. Wheat	84 cwts. 39 bush. 104 bush. 21 bush.	0% cwt. 2154 lbs. 1013 lbs. 2143 lbs.	9½ cwts, 4182 lbs. 1689 lbs, 3473 lbs.	68 cwts. 334 bush. 78 bush. 194 bush.	4% cwts, 1615 lbs. 978 lbs. 1966 lbs.	72% cwts. 3394 lbs. 1463 lbs. 3222 lbs.	1764 cwts, 47½ bush, 20% bush, 23% bush,	8% cwts. 2595 lbs. 1990 lbs. 3003 lbs.	185 cwt 5148 lbs 3343 lbs 4567 lbs
				6тн Соп	rse, 1868-7	1.		1	110	
1868 1869 1870 1871	Swedish Turnips Barley Beans Wheat	Faile 24g bush. 13g bush. 20g bush.	d, and ploughed 1948 lbs. 738 lbs. 2799 lbs.	up. 3358 lbs. 1591 lbs. 4092 lbs.	Faile 284 bush. 154 bush. 234 bush.	ed, and ploughed 2025 lbs. 768 lbs. 3048 lbs.	up. 3686 lbs. 1778 lbs. 4521 lbs.	Fail 427 bush. 248 bush. 24 bush.	ed, and ploughed 3309 lbs. 1056 lbs. 3440 lbs.	5800 lbs 2664 lbs 4942 lbs
MIDS		1/1		7тн Соп	se, 1872-7	j				
1872 1873 1874 1875	Swedish Turnips Barley Clover Wheat	34½ cwfs. 23½ bush. 21½ bush.	84 cwts. 1343 lbs. 2430 lbs.	42% cwts. 2717 lbs. 31% cwts. 3784 lbs.	1704 cwts. 204 bush. 284 bush.	17\$ cwts. 1565 lbs. 3536 lbs.	188 cwfs. 2875 lbs. 52½ cwts. 5328 lbs.	339 cwts. 314 bush. 315 bush.	354 cwts. 1723 lbs. 4685 lbs.	375§ cw 3573 lbs 84½ cw 6699 lbs.
				STH COU	RSE, 1876-79),		, ,		71
1876 1877 1878 1879	Swedish Turnips Barley Beans Wheat	17‡ cwts. 23‡ bush. 8‡ bush. 10≩ bush.	5 ewts. 1291 lbs. 740 lbs. 1324 lbs.	224 cwts. 2623 lbs. 1301 lbs. 1987 lbs.	1884 cwts. 244 bush. 74 bush. 148 bush.	28½ cwts. 1174 lbs. 1045 lbs. 1771 lbs.	2164 cwts. 2558 lbs. 1557 lbs. 2729 lbs.	356 cwts. 344 bush. 204 bush. 13 bush.	55‡ cwts. 1918 lbs. 1655 lbs. 1658 lbs.	4114 cwts 3890 lbs. 2963 lbs. 2493 lbs.
				9тн Соц	se, 1880-8	3.				41
1880 1881 • 1882 1883	Swedish Turnips Barley Clover or Beans Wheat	14 cwts.	24 cwts.	16‡ cwts.	199½ cwts.	11% cwts.	2114 cwts.	439½ cwts.	43‡ cwts.	4824 cwts.
		Sur	MMARY-AV	ERAGE OF THI	First 8 Co	urses, 1848	-1879.			
8, '52, '56, 0, '64, '72, '76 9, '53, '57, '61, 5, '69, '73, '77 0, '54, '58, '62, 6, '70, '74, '78	(calcd as hay)	263 cwts. 344 bush.	9} cwts. 2159 lbs.	35 cwts. 4132 lbs. 42 cwts. 1867 lbs.	1484 cwts. 284 bush.	26§ cwts. 1730 lbs.	1753 cwts. 3346 lbs. 55 cwts. 1996 lbs.	272½ cwts. 41½ bush.	43% cwts. 2577 lbs.	3157 cw 4955 lbs. 738 cwt 3230 lbs.
1,'55,'59,'63, 7,'71,'75,'79	Wheat	25% bush.	2905 lbs.	4559 lbs.	274 bush.	3067 lbs.	1996 10s. 4823 lbs.	30 g bush.	3698 lbs.	3230 lbs

(0) First Course—100 lbs. Bone-ash, and 100 lbs. Sulphuric Acid (sp. gr. 1-7;) Second Course—160 lbs. Bone-ash, 120 lbs. Sulphuric Acid; Thirt, Fourth, Fifth, Sixth, Serenth, Eighth, and Ninth Courses—200 lbs. Bone-ash, and 150 lbs. Sulphuric Acid, per acre.

(2) First Course—100 lbs. Pearl-ash, 100 lbs. Bone-ash, 100 lbs. Sulphuric Acid, 100 l

Ammonia, and 2000 lbs. Rape-cake; Third, Fourth, Fifth, Sixth, Seventh, Eighth, and Ninth Courses—300 lbs. Sulphate of Potass, 200 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 200 lbs. Done-sah, 150 lbs. Sulphate of Jamonia, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake, per acre.

(5) The quantities give in Builder tepresent the Dressed Corn only.

(6) The "Total Produce" of the Corn-crops includes Dressed Corn, Offial Corn, Straw, and Chaff.