Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



## Yields of the Field Experiments 1898



Full Table of Content

## Wheat Alternated With Fallow, and Wheat Grown Continuously

## **Rothamsted Research**

Rothamsted Research (1899) Wheat Alternated With Fallow, and Wheat Grown Continuously; Yields Of The Field Experiments 1898, pp 32 - 33 - DOI: https://doi.org/10.23637/ERADOC-1-228

## AND WHEAT GROWN CONTINUOUSLY, FALLOW. WHEAT ALTERNATED WITH EXPERIMENTS ON

acre, on the half acre of wheat after fallow; and in the second column the produce per acre obtained in the adjoining field (Broadbalk), where wheat is grown year after year on the same land. Lastly, in the third column of each of the vertical divisions is given the amount of produce after fallow, + or - that grown year after year on the same land. The results for the individual years show that during the earlier years of the experiments on alternate wheat and fallow, when the accumulations due to previous The results given in the following Table show the produce of Wheat obtained on the Rothamsted soil for many years in succession, after bare fallow, compared with that of wheat grown continuously year after year on the same land, without the intervention Rothamsted soil for many years in succession, after bare wheat grown continuously year after year on the same la of fallow; in both cases without manure.

Hoos-field, in which the experiments or

Hoos-field, in which the experiments on alternate wheat and fallow are conducted, adjoins Broadbalk-field, in which wheat has now been grown continuously without plot of that field, is compared with that grown in manure (also with different descriptions of manure), for 55 years in succession; and the produce of the unmanured plot of that field, is com-alternation with fallow, also without manure, in Hoos-field.

During the first or preliminary period of 5 years, 1851-1855, the cropping of the acre set apart for the experiment on wheat alternated with fallow was as follows:— Fallow (after wheat in 1850); 1852, Wheat; 1853, Fallow; 1854, Wheat; half Fallow, and half Wheat. From that time to the present the respective 1855, half Fallow, and half Wheat. From that time to the present the respective halves have been alternately fallow and wheat, giving therefore a crop of wheat suc-The description of seed sown has been the same in the two fields in the corresponding years; namely—for the crop of 1852 "Red Cluster"; for 28 years, 1854 to 1881 inclusive, "Red Rostock"; and for 1882, and since, "Club" or "Square Head" (Red).

In the upper division of the Table are given the results for each of the five years of the upper division of the main division are recorded the results for each individual year of the exact experiment, from 1856 up to the present time. In the first column of each main vertical division of the Table is given the produce per ceeding fallow, on half the acre each year.

also more straw, per acre per annum, than where the crop is grown continuously. On the other hand, if the produce after fallow is reckoned (as in the bottom division) at the yield per acre of the whole area, half in crop and half fallow, it gives several bushels less grain, and also less straw, per acre per annum, than where the crop is grown year after year on the same land

Referring to the two sets of averages at the foot of the Table, it is seen that if (as in the upper of the two divisions), the produce after fallow is reckoned at the yield per acre of the half in crop each year, it gives on the average several bushels more grain, and

treatment were less exhausted, the produce after fallow was more in excess of that grown in the adjoining field year after year on the same land than afterwards.

of a crop every other year; and that a given area of land yields more when the crop is grown year after year than when alternated with fallow. The explanation doubtless is, that much of the nitrogen brought into an available condition under the influence of the fallow, is lost by drainage during the long period that the land is without a crop. The conclusion to be drawn is, that although there is an increase of produce after fallow compared with that of wheat grown continuously, it is obtained at the sacrifice (Area under experiment, 1 acre.

			1851 1852 1853 1854 1854	or.	1856	1858 1859	1860	1862	1863	1864	1865
nd Straw).	After Fallow + or – after Wheat.		1bs. -2710 +4565 -1772 +3758 -45		+1051	+1287	- 274 +1227	+ 946	+2263	+2323	+1729
Fotal Produce (Grain and Straw).	Wheat after Wheat each year.		1bs. 27110 2457 1772 3496 2859		2450 2813	3226	2197 1990	2709	2727	2428	1861
Total Produ	Wheat after Fallow each year.		1bs. 7022 Fallow 7254 2814		3501 5374	4038 5662	1923 3217	3655	4990	4751	3590
	After Fallow + or - after Wheat.		1bs. -1627 +3337 -1413 +2408 - 53		+ 555 +1498	+ 798	+ 818	+ 581	+1300	+1396	+1117
Fotal Straw.	Wheat after Wheat each year.		1627 1597 1418 2137 1787		1558	2175	1459 1254	1713	1600	1350	1033
1	Wheat after Fallow each year.	٥.	1bs. 4934 Fallow 4545 1734	COMPARISON.	2113	3686	1226 2072	2294	2900	2746	7150
The state of the s	After Fallow + or - after Wheat,	Y PERIOD	1bs. -1083 +1228 - 359 +1350 + 8		+ 496 +1063	+ 925	- 41 + 409	+ 365	+ 963	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	+ 612
Total Grain.	Wheat after Wheat each year.	PRELIMINARY	1083 860 359 1359 1072	OF EXACT	892 1236	1051	738	966	1127	1078	878
n water)	Wheat after Fallow each year.	PR	1bs. Fallow 2088 Fallow 2709 1080	Period	1388	1630	697 1145	1361	2090	2002	1440
r Bushel.	Wheat after Wheat each year.		lbs. 61·1 56·6 45·9 60·6 59·2		54.3	60.4 52.5	52·6 57·4	8.12	62.7	62.0	9.09
Weight per Bushel	Wheat after Fallow each year.		lbs. 53 · 0 Fallow 60 · 5 54 · 0		60.0 58.4	55.0	54.8	57.1	61.4	61.7	9.76
	After Fallow + or – after Wheat.		Bushels. — 157 + 234 - 534 + 21 + 03		+ 74 + 18	+153	1+	+	+155	十143	+111
Dressed Grain.	Wheat after Wheat each year.		Bushels, 157, 153, 254, 21, 17		144 20	20 St	124	16	174	16½	13‡
9	Wheat after Fallow each year.		Bushels. Fallow 37 Fallow 42 173		38	255 4 4 4	12½ 177	222	323	318	243
			1851 1852 1853 1853 1854 1855		1856	1858 1859	1860	1862	1863	1864	1865

1866 1867	1868	1870	1871	1872	1874	1875	1876	1878	1879	1880	1881	1883	1884	1885	1886	1887	1888	1889	1890	1891	1893	1894	1896		1	5 yrs. 1851-'55	10 yrs. 1856-'65	10 yrs. 1866-75	10 yrs. 1876-'85 10 yrs. 1886-'95	40 mm 1050 30F	TO 318, 1000-30	LOW.	5 yrs. 1851-'55	10 mm 10K0 30F	10 yrs 1866 775	10 yrs. 1876–'85	10 yrs. 1886-'93
- 247 + 237	+ 2027	+ 381	+ 177	F 1	+1686	+1143	+ 283	1 368	+ 94	+ 764	1 364	+ 583	+1055	+1101	+ 111	+ 564	+ 459	19 +	+ 892	+ 414 + 414	+ 473	- 172 + 745	- 64	1	- 1	+ 759			+ +		1 1 1	HALF FALLOW.	- 950	488	689	599	COC
1505	2027	2002	1715	1603	1684	C/CI	1142	1857	1093	1838	2009	1878	1729	2062	1134	1801	1515	1645	1853	1425	1251	$\frac{2608}{1384}$	2396	EACH YEAR.		2659	2521	1821	1676	1991		AND	2659	- 9591	1821	1667	TOTAL
1742	400 <del>4</del>	2383	1892	1056	3370	2/18	1425	2825	1187	2602	1804	2461	2784	3163	1245	2365	1974	1712	2745	1839	1724	2436 2129	2332	CROP		3418	4076	2278	2182	9868		F IN CROP	1709	9038	1139	1068	TOOT
+ 153	+ 1425	+ 236	+ 187	- 27	+1010	/T/ +	+ 148 - 23	+ 573	+ 45	+ 516	- 245 10	+ 295	+ 639	+ 675	+ 87	+ 317	+ 338	+	+ 653	+ 272	+ 245	+ 431	+ 3	HA	m	+ 531		+ 341	+ 328	+ 468		AREA, HALF	- 590	- 303	- 367	- 348	2007
973	1350	1046	1159	905	066	TONO	642	1081	763	1149	1095	1006	902	1137	570	895	901	905	1314	836	609	1487 720	1309	E OF THE		1712	1539	1076	924	1127		WHOLE A	1712	1539	1076	967	
1126	1019	1282	1807	875	2000	07/1	062 889	1654	808	1665	1085	1301	1544	1812	657	1212	1239	916	1657	1108	854	1483	1312	PER ACRE	0700	2243	2473	1917	1252	1595		OF THE W	1122	1236	602	619	
1 +-	+ 007   193	+ 145	1 + 12	- 520	949 +	07F T	+ + 109	+ 395	+ 49	+ 248	1 + 1 04	+ 288	+ 416	+ 456	+ 24	+ 247	+ 121	+	+ 259 + 576	+ 142	+ 228	+ 314	- 67 - 132	YLELD	4	4 228	+ 621	1100	+ 178	+ 278	- 1	ACRE	- 360	- 180	- 315	- 251 - 287	0 20
532	848	956	705	701	694	100	543	977	330	689	629	872	824	925	564	906	614	743	828 828	589	642	664	1087	D AT THE	041	34.1	982	7007	752	795		YIELD PER	947	982	745	700	100
616	655	1011	280	181	1370	200	643 649	1171	379	937	719	1160	1240	1351	588	1153	735	1000	1404	731	870	978	1020	RECKONED	1172	CVII	1603	899	930	1073		AT THE Y	587	802	430	449 465	963
56-1	56.1	61.8	59.0	57.0	28.0		58.9	59.0	52.5	56.9	58.7	61.2	62.1	0.60	61.5	20.00	20.00	23.8	57.4	59-6	62.7	62.5	61.4	FALLOW	50.7	1 00	57.9	0 00	60.5	58.8		ECKONED 4					
286.8	60.1	62.5	58 6	42.0	60.0	1 0 1	60.5	6.75	55.6	54.6	58.6	61.2	60.2	6.70	62.5	59-9 56-1	200.	50.0	58.0	60.2	62.4	62.2	60·7 59·5	AFTER	и п.	0 00	28.2	58.4	1.09	58.7		FALLOW RE				D	İ
+ + 200%	-   g.41	+ 23	4	6 	+10	2 6	+ + \$200							- 1					- +			1 + 1 10 10 40	175	-Рворисв	1 43	62 5	+ TO\$	₩ cc	+	+ 45		AFTER HA	- 5½	- 2 <sub>3</sub>	ا ي ي	 	41
888	143	15	108	118	11. \$2.55	9 5	0000	124	4, -	40 0% -1 00	113	133	n :	801	6	- 1. - 1. - 1.	101	47	13%	ರಾ ೧	2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,	10	16 <del>2</del> 83	AVERAGES	143	444	10% 1174	8 T	$12^{\frac{1}{3}}$	123	- 11	PRODUCE A	143	157	211	121	198
1 00 TC	$10\frac{1}{4}$	171	123	C7 8 <del>14</del>	21 <del>\$</del> 161	101	103	194	161	124	113	00°	20 <del>4</del>	67	Q (	108	421	173	23,4	11. 84.	152	152	16g 7	Av	191	400	7.0g 1.31	1 L 2 4 L 2 8 4 L	151	173			95	13	0 84	nio To	ŝ
1867	1869	1870	1872	1873	1875	9781	1877	1878	1878	1881	1882	1883	1884	COOT	1886	7007	1880	1890	1891	1892	1893	1895	1896 1897		5 vrs. 1851-755		10 yrs 1856-'05	VI'S	yrs.	40 yrs. 1856-'95		AVERAGES	yrs.	yre.	yrs	10 yrs. 1876–'85 10 yrs. 1886–'95	40 vrs 1856 '95