


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1901

[Full Table of Content](#)



Rotation; Agdell Field


Rothamsted Research

Rothamsted Research (1902) *Rotation; Agdell Field* ; Yields Of The Field Experiments 1901, pp 110 - 121 - DOI: <https://doi.org/10.23637/ERADOC-1-229>

PLAN OF THE PLOTS IN AGDELL FIELD,
ON WHICH EXPERIMENTS HAVE BEEN MADE
ON FOUR-COURSE ROTATION.

54 years, commencing 1848.

[For a brief summary of results and conclusions, see opposite page.]



Total area of ploughed land about 3 acres.
Area of each of the 12 divisions $\frac{1}{2}$ acre.

- The 4 lower divisions, Unmanured continuously (Plot 1).
- The 4 middle divisions, Mineral Manure, for the Roots, each Course (Plot 2).
- The 4 upper divisions, Mineral and Nitrogenous Manure, for the Roots, each Course (Plot 3).
- The 6 left-hand divisions, Clover (or Beans), 3rd year each Course.
- The 6 right-hand divisions, Fallow, 3rd year each Course.

The double lines indicate division paths between plot and plot.

[For details of the manuring and produce, see pp. 112-121.]

RESULTS OF EXPERIMENTS MADE IN AGDELL FIELD, ON
THE ROTATION OF CROPS.

The experiments were commenced in 1848; so that 1901 is the 54th year of their continuance, and the second year of the 14th Course. In the experiments in other fields, some of the most important crops of rotation have been grown, each separately, for many years in succession—without manure, with farmyard manure, and with various artificial manures. But besides such experiments, others have been made on the growth of the crops in an actual course of rotation, without manure, and with different manures. The results with the individual crops throw much light on the characteristic requirements of each particular crop; whilst those on the growth of the crops in rotation serve to confirm and control those with the individual crops.

The rotation selected for investigation was the well known and typical four-course rotation of—1. Turnips; 2. Barley; 3. Leguminous Crops (or Fallow); 4. Wheat; that is, an alternation of Root-crops and of Leguminous Crops with cereals; which is the basis of most of the various rotations adopted in different parts of our own country, and also in many other countries. One portion of the land was left entirely without manure each course; another received mineral manure only, for the turnips of each course; and a third mixed mineral and nitrogenous manures, also only for the turnips of each course.

1. *The Swedish Turnips commencing each Course.*—When various root-crops were grown year after year on the same land without manure, they soon reverted to the uncultivated condition; and the experiments on rotation show that the Swedish turnips grown once in four years in unmanured rotation, came down to only about 1 ton per acre. The results further show, that mineral manures alone applied for the root-crops gave considerable increase, but that mineral and nitrogenous manures together gave more still. Without manure, the average produce of roots was less over the last 4 than over the preceding 8 courses; but with mineral manure alone (including potash in the last 4 courses) it was higher, and with mineral and nitrogenous manures together much higher, over the last 4 courses; the result being, however, largely due to more favourable seasons. Indeed, in 1888 and 1892, the years of root-crop in the 11th and 12th courses, although the produce without manure was less, that by each of the two descriptions of manure was considerably more than the average of the preceding courses; that is, both the reversion to the uncultivated condition without manure, and the increased growth with suitable manures, were very marked. In fact, without manure the produce of roots was as restricted in rotation as in continuous growth; with purely mineral manure it was greater in rotation than in continuous growth, the exhaustion of the available nitrogen of the soil being less under rotation; and with the mixed mineral and nitrogenous manure much more produce was obtained under rotation than with continuous growth. Lastly, the results conclusively show how artificial a product is the cultivated root-crop, and how dependent it is for its successful growth on an abundant supply of available food—nitrogenous as well as mineral—within the soil.

2. *The Barley Crops.*—Barley, without manure, succeeded the differently manured Swedish turnip crops of each course. Although the average produce of the root-crops was greater over the last 4 (10th, 11th, 12th, and 13th) than over the preceding 8 courses, the succeeding barley crops were much less over the last 4 courses. This was the case, not only where the root-crops had been carted off, but also where they had not been so removed. As, however, the produce of barley in the 4 years in question (1885, 1889, 1893, and 1897) was also less than the average in Hoos Field where the crop is grown year after year, the result is doubtless mainly due to the seasons. Then, the average produce of barley over the 8 courses was actually less after the carted off roots grown by mineral manure (super-phosphate) than after those grown without manure. The explanation is—that as there was practically no produce of roots without manure the unmanured plot was practically fallow for the barley; whilst with the mineral manure fair crops of roots were grown and removed, leaving the surface soil the more exhausted of its available nitrogen and other constituents. In the later years, however, after such long continued exhaustion, the unmanured plot has yielded less barley after the removal of the roots than the mineral manured plot. On the other hand, where the roots were not removed from the land, the mineral manured plot has generally yielded more barley than the unmanured. Further, under all conditions of treatment, the plots with mixed mineral and nitrogenous manure have yielded more barley than those with the mineral manure alone. In fact, the effects of the manurial and other treatment of the first crop of the course are clearly manifested in the produce of the second crop. Lastly, both without manure, and with the mineral manure alone, there was more produce of barley in rotation than in continuous growth, but with mixed mineral and nitrogenous manure there was more produce when the crop was grown continuously, the supply of nitrogen in that case being somewhat larger and annually applied for the crop.

3. *The Leguminous Crops (or Fallow).*—Under equal conditions as to manuring, the Leguminous crops, especially the clover, bring much more nitrogen into the course than either of the other crops. Further, the amount of nitrogen so brought into the rotation is much greater under the influence of mineral manures, and especially of potash manures, than without manure; whilst under the influence of the mixed mineral and nitrogenous manure the yield of nitrogen is greater still, the leguminous crop utilising the unexhausted nitrogenous manure- and crop-residue. For the successful growth of leguminous crops, however, a liberal supply of available mineral constituents within the soil, especially potash and lime, is essential. Judging from comparable cases, the amount of nitrogen accumulated by the Leguminous crops was much greater when they were grown in rotation, than is only occasionally, than when grown continuously. With fallow instead of a Leguminous crop, there is very much less nitrogen yielded in the rotation, and more liability to loss of it by drainage, and hence so much less brought into the circulation of the farm for food or manure. Lastly, most of the nitrogen of the leguminous crop is retained on the farm; and there is more or less, and sometimes much nitrogenous crop-residue left in the soil for succeeding crops.

4. *The Wheat Crops.*—There was very much more produce of wheat both without manure and with mineral manure, and considerably more with the mineral and nitrogenous manure, when it was grown in rotation than under comparable conditions continuously. Taking the quantities of produce by the mixed mineral and nitrogenous manure the result was that the two cereal crops produced approximately equal amounts of dry substance, and each considerably more than either of the assumed restorative crops—the roots or the leguminous crops. The supply of nitrogen within the soil available to the wheat crop is increased both by fallow and by the growth of a leguminous crop, especially of clover; and the accumulation is the greater when the soil and subsoil are not abnormally exhausted of organic nitrogen.

Upon the whole the results show that the benefits of rotation are very various. They depend on the varying requirements, habits of growth, and capabilities of gathering and assimilating the necessary constituents, of the different crops. The difference in the amounts available within the soil of the various mineral constituents, is one element in the explanation; but the facts relating to the amount, and to the sources, of the nitrogen of the different crops, are of still greater significance. The uses of the different crops have also to be taken into account. The cereals yield more produce for sale in the season of growth in rotation than when grown continuously. The crops alternated with them accumulate very much more of mineral constituents and of nitrogen in their produce; but by far the greater proportion of those constituents remains in circulation in the manure of the farm, whilst the remainder yields highly valuable products for sale in meat and milk. Again, with a variety of crops, the operations of the farm are better distributed over the year, and are therefore more economically performed. Lastly, the opportunities which alternate cropping afford for cleaning the land constitute a prominent element of advantage.

For details of the manuring and produce of the different plots, see pages 112-121.

AGDELL FIELD.

(Area under experiment, about 3 acres.)

EXPERIMENTS ON AN ACTUAL COURSE OF ROTATION—TURNIPS, BARLEY, LEGUMINOUS CROP (OR FALLOW), AND WHEAT.

These Experiments were commenced in 1848; so that the present season (1901) is the 54th, and the growing crop (Barley) is the second of the Fourteenth Course. One-third of the land has been continuously unmanured. One-third has, for the first Nine Courses, or 36 years, 1848-83, been manured with Superphosphate of Lime alone, once every four years, that is for the turnip-crop commencing each course; but for the Tenth, Eleventh, Twelfth, Thirteenth, and Fourteenth Courses, a complex mineral manure has been applied, as described in foot-note, No. 2. Lastly, one-third has been manured also for the turnip-crop only, with a complex mineral and Nitrogenous manure, as described in the foot-note No. 3. From half of each of the three differently manured plots the turnip-crops (roots and leaves) are removed; and on the other half they are either consumed on the land by sheep, or spread and ploughed in. In the case of all the other crops, the total produce is removed from the land. In the First Course, clover was sown over the whole of each of the three differently manured plots; but in the subsequent courses, a leguminous crop was grown on only half of each of the three plots, the other half being left fallow, in the third year of each course. In the Second, Third, and Fourth Courses, clover was sown, but failed; and in them, and in the Fifth and Sixth Courses, beans were taken instead. In the Seventh Course, clover was sown (spring 1873), and gave three cuttings in 1874. In the Eighth Course beans were grown. In the Ninth Course clover was sown (in the spring of 1881), and gave two cuttings in 1882. In the Tenth Course clover was sown (in the spring of 1885), and yielded two cuttings in 1886. In the Eleventh Course clover was sown (with the barley) in 1889, but failed during the winter, and in 1890 beans were grown instead. In the Twelfth Course clover was again sown in April 1893, and gave two cuttings in 1894. In the Thirteenth Course clover was sown (with the barley), April 1897, but failed during the winter, and in 1898 beans were grown instead. In the Fourteenth Course clover was sown (with the barley), May 4, 1901.

TABLE I. (below), gives the results relating to the portions of each plot from which the turnip-crops were entirely removed; and on which clover or beans were grown.

Years.	Description of Crop.	PRODUCE PER ACRE.									
		Plot 1. Unmanured continuously.			Plot 2. Superphosphate of Lime alone. (1) Courses 1-9, Complex Mineral Manure (2), Courses 10-14, for the Turnip Crops only.			Plot 3. Complex Mineral and Nitrogenous Manure, (3) for the Turnip Crops only.			Total Produce. (5)
		Corn (4) (or Roots).	Straw (or Leaf).	Total Produce. (5)	Corn (4) (or Roots).	Straw (or Leaf).	Total Produce. (5)	Corn (4) (or Roots).	Straw (or Leaf).	Total Produce. (5)	
1848	Norfolk White Turnips	65½ cwts.	45½ cwts.	111½ cwts.	225½ cwts.	106½ cwts.	320 cwts.	218 cwts.	15½ cwts.	368½ cwts.	
1849	Barley	44½ bush.	2983 lbs.	5656 lbs.	20½ bush.	2111 lbs.	3941 lbs.	28½ bush.	2083 lbs.	3794 lbs.	
1850	Clover (calcd. as hay) (6)	28½ bush.	3431 lbs.	52½ cwts.	28 bush.	3371 lbs.	523½ lbs.	28½ bush.	3582 lbs.	614 cwts.	
1851	Wheat	26 cwts.	44 cwts.	5389 lbs.	26 cwts.	3371 lbs.	523½ lbs.	28½ bush.	3582 lbs.	5500 lbs.	
1852	Swedish Turnips	26 cwts.	44 cwts.	30½ cwts.	223½ cwts.	20½ cwts.	243½ cwts.	39½ cwts.	364 cwts.	433 cwts.	
1853	Barley	34½ bush.	2430 lbs.	4464 lbs.	28½ bush.	1873 lbs.	3560 lbs.	34½ bush.	2604 lbs.	4873 lbs.	
1854	Beans	34 bush.	1055 lbs.	1445 lbs.	34 bush.	1103 lbs.	1534 lbs.	34 bush.	1353 lbs.	2065 lbs.	
1855	Wheat	34 bush.	3619 lbs.	5559 lbs.	35½ bush.	3325 lbs.	5789 lbs.	37½ bush.	3942 lbs.	6371 lbs.	
1856	Swedish Turnips	32 cwts.	24 cwts.	34½ cwts.	136 cwts.	7½ cwts.	143½ cwts.	333½ cwts.	12½ cwts.	346½ cwts.	
1857	Barley	48½ bush.	2600 lbs.	5337 lbs.	28½ bush.	1475 lbs.	3076 lbs.	48 bush.	2435 lbs.	5165 lbs.	
1858	Beans	64 bush.	1100 lbs.	1515 lbs.	64 bush.	1165 lbs.	1605 lbs.	12½ bush.	150 lbs.	2357 lbs.	
1859	Wheat	34 bush.	4030 lbs.	6262 lbs.	34½ bush.	2930 lbs.	6120 lbs.	30½ bush.	4610 lbs.	7154 lbs.	
1860	Swedish Turnips	1 cwt.	(64 lbs.)	1 cwt.	294 cwts.	1½ cwt.	30½ cwts.	8½ cwts.	2½ cwts.	90½ cwts.	
1861	Barley	28½ bush.	2522 lbs.	4718 lbs.	30½ bush.	2000 lbs.	3775 lbs.	64 bush.	3940 lbs.	7381 lbs.	
1862	Beans	23 bush.	1840 lbs.	3661 lbs.	29½ bush.	2150 lbs.	4040 lbs.	43½ bush.	3280 lbs.	5580 lbs.	
1863	Wheat	34 bush.	3448 lbs.	5621 lbs.	34½ bush.	3530 lbs.	5619 lbs.	44½ bush.	499 lbs.	7657 lbs.	
1864	Swedish Turnips	88 cwts.	0½ cwt.	88 cwts.	68 cwts.	4½ cwts.	72½ cwts.	176½ cwts.	8½ cwts.	185 cwts.	
1865	Barley	39 bush.	2154 lbs.	4182 lbs.	33½ bush.	1615 lbs.	3394 lbs.	47½ bush.	2595 lbs.	5148 lbs.	
1866	Beans	104 bush.	1013 lbs.	1629 lbs.	37 bush.	978 lbs.	1463 lbs.	20½ bush.	1990 lbs.	3343 lbs.	
1867	Wheat	21 bush.	2143 lbs.	3473 lbs.	19½ bush.	1966 lbs.	3222 lbs.	23½ bush.	3003 lbs.	4567 lbs.	

1st Course, 1848-51

2nd Course, 1852-55

3rd Course, 1856-59

4th Course, 1860-63

5th Course, 1864-67

Year	Crop	Failed, and ploughed up.	Failed, and ploughed up.	Failed, and ploughed up.	Failed, and ploughed up.
1868 1869 1870 1871	Swedish Turnips	24 bush.	1448 lbs.	3538 lbs.	42 1/2 bush.
	Barley	13 1/2 bush.	738 lbs.	1778 lbs.	38 3/4 bush.
	Beans	20 1/2 bush.	2799 lbs.	4092 lbs.	24 bush.
1872 1873 1874 1875	Swedish Turnips	3 1/2 bush.	8 1/2 cwt.	42 1/2 cwt.	33 3/4 cwt.
	Barley	23 1/2 bush.	1343 lbs.	2717 lbs.	31 1/2 bush.
	Clover (calcd. as hay) (c)	21 1/2 bush.	2430 lbs.	3784 lbs.	31 1/2 bush.
1876 1877 1878 1879	Swedish Turnips	17 1/2 cwt.	5 cwt.	22 1/2 cwt.	35 1/2 cwt.
	Barley	24 bush.	1291 lbs.	2623 lbs.	34 bush.
	Beans	8 1/2 bush.	740 lbs.	1301 lbs.	13 bush.
1880 1881 1882 1883	Swedish Turnips	14 cwt.	24 cwt.	16 1/2 cwt.	43 1/2 cwt.
	Barley	26 1/2 bush.	1484 lbs.	2922 lbs.	35 3/4 bush.
	Clover (calcd. as hay) (c)	29 1/2 bush.	2280 lbs.	4175 lbs.	45 1/2 bush.
1884 1885 1886 1887	Swedish Turnips	5 cwt.	1270 lbs.	1960 lbs.	28 1/2 cwt.
	Barley	12 bush.	603 lbs.	1079 lbs.	34 1/2 bush.
	Clover (weighed as hay) (c)	25 1/2 bush.	1859 lbs.	3433 lbs.	42 1/2 bush.
1888 1889 1890 1891	Swedish Turnips	24 cwt.	17 cwt.	4 1/2 cwt.	47 1/2 cwt.
	Barley	11 bush.	931 lbs.	1510 lbs.	26 1/2 bush.
	Beans	7 bush.	603 lbs.	1079 lbs.	15 1/2 bush.
1892 1893 1894 1895	Swedish Turnips	6 1/2 cwt.	0 1/2 cwt.	7 1/2 cwt.	47 3/4 cwt.
	Barley	16 1/2 bush.	1440 lbs.	2446 lbs.	20 1/2 bush.
	Clover (weighed as hay) (c)	23 1/2 bush.	1713 lbs.	3267 lbs.	39 bush.
1896 1897 1898 1899	Swedish Turnips	7 1/2 cwt.	1 1/2 cwt.	8 1/2 cwt.	34 1/2 cwt.
	Barley	11 1/2 bush.	1251 lbs.	1927 lbs.	30 1/2 bush.
	Beans	24 1/2 bush.	1335 lbs.	2976 lbs.	24 1/2 bush.
1900 1901 1902 1903	Swedish Turnips	16 cwt.	4 1/2 cwt.	20 1/2 cwt.	46 1/2 cwt.
	Barley	16 bush.	1251 lbs.	1927 lbs.	23 1/2 bush.
	Beans	24 1/2 bush.	1335 lbs.	2976 lbs.	24 1/2 bush.

(1) First Course—100 lbs. Bone-ash, and 100 lbs. Sulphuric Acid (sp. gr. 1.7); Second Course—160 lbs. Bone-ash, 120 lbs. Sulphuric Acid; Third, Fourth, Fifth, Sixth, Seventh, Eighth, Ninth, and Tenth Courses—made from high percentage mineral phosphate, and containing 37 per cent. of soluble phosphate.
 (2) For the Tenth Course, in addition to the Superphosphate for the Swedish Turnips—300 lbs. Sulphate of Potash, 200 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia were applied February 25, 1884, and harrowed in; and the same quantities were applied again before the final ploughing and preparation of the land for the sowing of the seed in May. For the Swedish of the Eleventh and Twelfth Courses the same mineral manures (which are the same as the mineral manures of Plot 3 for the third and subsequent Courses) were again applied, but only once for each of these two Courses. For the Swedish of the Thirteenth and Fourteenth Courses—500 lbs. Sulphate of Potash, 100 lbs. Sulphate of Soda, 200 lbs. Sulphate of Magnesia, and 600 lbs. Sulphuric Acid, 100 lbs. Bone-ash, 100 lbs. Sulphate of Ammonia, and 100 lbs. Pearl-ash, 100 lbs. Bone-ash, 100 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, and 100 lbs. Muriate of Ammonia, and 1000 lbs. Rape Cake; Second Course—300 lbs. Sulphate of Potash, 100 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 160 lbs. Bone-ash, 120 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake; Third, Fourth, Fifth, Sixth, Seventh, Eighth, Ninth, and Tenth Courses—300 lbs. Sulphate of Potash, 200 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake, per acre; Eleventh and Twelfth Courses—the same in other respects as in Courses 3-10, but the Superphosphate made from high percentage mineral phosphate, and containing 37 per cent. of soluble phosphate. For the Swedish of the Thirteenth and Fourteenth Courses—500 lbs. Sulphate of Potash, 100 lbs. Sulphate of Soda, 200 lbs. Sulphate of Magnesia, 600 lbs. Basic Slag, 2000 lbs. Rape-cake, 100 lbs. Sulphate of Ammonia, and 100 lbs. Muriate of Ammonia, per acre. (3) The quantities given in Swedish represent the Dressed Corn only.
 (4) The "Total Produce" of the Corn-crops includes Dressed Corn, Offal Corn, Straw, and Chaff. (5) Two cuttings. (6) Three cuttings.

[For Summary Table of the above results, see pp. 120-121.]

A G D E L L F I E L D .

(Area under experiment, about 3 acres.)

EXPERIMENTS ON AN ACTUAL COURSE OF ROTATION—TURNIPS, BARLEY, LEGUMINOUS CROP (OR FALLOW), AND WHEAT.

These Experiments were commenced in 1848; so that the present season, 1901, is the 54th, and the growing crop (Barley) is the second of the Fourteenth Course. One-third of the land has been continuously unmanured. One-third has, for the first Nine Courses, or 36 years, 1848-83, been manured with Superphosphate of Lime alone, once every four years, that is for the turnip-crop commencing each course; but for the Tenth, Eleventh, Twelfth, Thirteenth, and Fourteenth Courses, a complex mineral manure has been applied, as described in foot-note, No. 2. Lastly, one-third has been manured (also for the turnip-crop only), with a complex mineral and Nitrogenous manure, as described in the foot-note, No. 3. From half of each of the three differently manured plots, the turnip-crops (roots and leaves) are removed; and on the other half they are either consumed on the land by sheep, or spread and ploughed in. In the case of all the other crops, the total produce is removed from the land. In the First Course, clover was sown over the whole of each of the three differently manured plots, and on which, in the third year of each course (excepting the first, 1850, when clover was grown), the land was left fallow.

TABLE II. (below), gives the results relating to the portions of each plot from which the turnip-crops were entirely removed; and on which, in the third year of each course (excepting the first, 1850, when clover was grown), the land was left fallow.

Years.	Description of Crop.	PRODUCE PER ACRE.					
		Corn (4) (or Roots).	Straw (or Leaf).	Total Produce (5).	Corn (4) (or Roots).	Straw (or Leaf).	Total Produce (5).
1st Course, 1848-51.	Swedish Turnips	175½ cwts.	19½ cwts.	195 cwts.	292 cwts.	35 cwts.	327 cwts.
	Barley	33½ bush.	2200 lbs.	4149 lbs.	294 bush.	1870 lbs.	3575 lbs.
	Wheat (calcd. as hay) (6)	30½ bush.	3273 lbs.	5290 lbs.	31½ bush.	3497 lbs.	604 cwts.
2nd Course, 1852-55.	Swedish Turnips	37 cwts.	5½ cwts.	42½ cwts.	256½ cwts.	22½ cwts.	279½ cwts.
	Barley	32½ bush.	2187 lbs.	4046 lbs.	32 bush.	2003 lbs.	3876 lbs.
	Wheat	37½ bush.	4295 lbs.	6735 lbs.	38½ bush.	4286 lbs.	6756 lbs.
3rd Course, 1856-59.	Swedish Turnips	45½ cwts.	2½ cwts.	47½ cwts.	170½ cwts.	8 cwts.	178½ cwts.
	Barley	43½ bush.	2330 lbs.	4777 lbs.	30½ bush.	1545 lbs.	3272 lbs.
	Wheat	35½ bush.	4315 lbs.	6582 lbs.	37½ bush.	4310 lbs.	6671 lbs.
4th Course, 1860-63.	Swedish Turnips	1½ cwts.	0½ cwt.	1½ cwts.	53½ cwts.	2 cwts.	55½ cwts.
	Barley	33½ bush.	2190 lbs.	4248 lbs.	32½ bush.	1954 lbs.	3807 lbs.
	Wheat	45 bush.	4563 lbs.	7446 lbs.	46 bush.	4690 lbs.	7626 lbs.
5th Course, 1864-67.	Swedish Turnips	7½ cwts.	0½ cwt.	8 cwts.	52½ cwts.	4½ cwts.	57½ cwts.
	Barley	34½ bush.	1828 lbs.	3659 lbs.	31½ bush.	1509 lbs.	3170 lbs.
	Wheat	27½ bush.	2654 lbs.	4330 lbs.	26½ bush.	2774 lbs.	4420 lbs.

1 lb. (pound avoird.) per acre = (about) 1.12 Kilogramme per Hectare, or 0.57 Zollverein Pfund, per Prussian Morgen.
 1 cwt. (hundredweight) per acre = (about) 125.5 Kilogrammes per Hectare, or 0.64 Centner per Pr. Morgen.

Plot 2. Superphosphate of Lime alone (1), Courses 1-9, Complex Mineral Manure (2), Courses 10-14, for the Turnip Crops only.

Plot 1. Unmanured continuously.

Plot 3. Complex Mineral and Nitrogenous Manure (3), for the Turnip Crops only.

Year	Course	Crop	Failed, and ploughed up.		Failed, and ploughed up.		Failed, and ploughed up.	
			bu.	lbs.	bu.	lbs.	bu.	lbs.
1868	6th Course, 1868-71	Swedish Turnips	21½ bush.	1638 lbs.	25½ bush.	1873 lbs.	39½ bush.	3064 lbs.
1869		Barley	11½ bush.	2075 lbs.	16½ bush.	2128 lbs.	17½ bush.	2628 lbs.
1870		Wheat	5½ cwt.	8½ cwt.	14½ cwt.	14½ cwt.	332 cwt.	344 cwt.
1871		Barley	20½ bush.	1374 lbs.	22½ bush.	1370 lbs.	31½ bush.	1626 lbs.
1872	7th Course, 1872-75	Fallow	24½ bush.	2893 lbs.	25½ bush.	3230 lbs.	28½ bush.	3623 lbs.
1873		Wheat	31½ cwt.	54 cwt.	19½ cwt.	17 cwt.	309½ cwt.	344½ cwt.
1874		Barley	21 cwt.	2602 lbs.	21 bush.	1054 lbs.	304 bush.	1625 lbs.
1875		Fallow	10½ bush.	1493 lbs.	14½ bush.	1956 lbs.	12½ bush.	1691 lbs.
1876	8th Course, 1876-79	Swedish Turnips	32½ cwt.	3½ cwt.	224 cwt.	124 cwt.	450½ cwt.	36 cwt.
1877		Barley	29½ bush.	1556 lbs.	24½ bush.	1239 lbs.	33½ bush.	1755 lbs.
1878		Fallow	33½ bush.	2994 lbs.	38½ bush.	3686 lbs.	37½ bush.	3689 lbs.
1879		Wheat	17½ cwt.	7½ cwt.	159½ cwt.	18½ cwt.	298½ cwt.	55½ cwt.
1880	9th Course, 1880-83	Swedish Turnips	15½ bush.	1786 lbs.	15½ bush.	965 lbs.	19 bush.	1328 lbs.
1881		Barley	34½ bush.	2505 lbs.	41½ bush.	3465 lbs.	39½ bush.	3308 lbs.
1882		Fallow	15 cwt.	22½ cwt.	142½ cwt.	154 cwt.	431½ cwt.	37½ cwt.
1883		Wheat	2941 lbs.	4868 lbs.	36 bush.	3586 lbs.	41 bush.	4288 lbs.
1884	10th Course, 1884-87	Swedish Turnips	9½ cwt.	1614 lbs.	226½ cwt.	4 cwt.	523½ cwt.	15½ cwt.
1885		Barley	19½ bush.	1614 lbs.	13 bush.	1203 lbs.	18½ bush.	1597 lbs.
1886		Fallow	21½ bush.	1630 lbs.	28½ bush.	2188 lbs.	32½ bush.	2368 lbs.
1887		Wheat	154 cwt.	34 cwt.	161 cwt.	8½ cwt.	345 cwt.	85 cwt.
1888	11th Course, 1888-91	Swedish Turnips	114 bush.	944 lbs.	124 bush.	969 lbs.	214 bush.	1465 lbs.
1889		Barley	26½ bush.	3081 lbs.	30½ bush.	3734 lbs.	33½ bush.	4006 lbs.
1890		Fallow	41½ cwt.	54 cwt.	199 cwt.	5½ cwt.	486½ cwt.	11½ cwt.
1891		Wheat	46½ cwt.	46½ cwt.	199 cwt.	5½ cwt.	486½ cwt.	498½ cwt.
1892	12th Course, 1892-95	Swedish Turnips	9½ cwt.	1614 lbs.	226½ cwt.	4 cwt.	523½ cwt.	15½ cwt.
1893		Barley	19½ bush.	1614 lbs.	13 bush.	1203 lbs.	18½ bush.	1597 lbs.
1894		Fallow	21½ bush.	1630 lbs.	28½ bush.	2188 lbs.	32½ bush.	2368 lbs.
1895		Wheat	154 cwt.	34 cwt.	161 cwt.	8½ cwt.	345 cwt.	85 cwt.
1896	13th Course, 1896-99	Swedish Turnips	114 bush.	944 lbs.	124 bush.	969 lbs.	214 bush.	1465 lbs.
1897		Barley	26½ bush.	3081 lbs.	30½ bush.	3734 lbs.	33½ bush.	4006 lbs.
1898		Fallow	41½ cwt.	54 cwt.	199 cwt.	5½ cwt.	486½ cwt.	11½ cwt.
1899		Wheat	46½ cwt.	46½ cwt.	199 cwt.	5½ cwt.	486½ cwt.	498½ cwt.
1900	14th Course, 1900-1903	Swedish Turnips	9½ cwt.	1614 lbs.	226½ cwt.	4 cwt.	523½ cwt.	15½ cwt.
1901		Barley	19½ bush.	1614 lbs.	13 bush.	1203 lbs.	18½ bush.	1597 lbs.
1902		Fallow	21½ bush.	1630 lbs.	28½ bush.	2188 lbs.	32½ bush.	2368 lbs.
1903		Wheat	154 cwt.	34 cwt.	161 cwt.	8½ cwt.	345 cwt.	85 cwt.

(1) First Course—100 lbs. Bone-ash, and 100 lbs. Sulphuric Acid (sp. gr. 1.7); Second Course—169 lbs. Bone-ash, 100 lbs. Sulphuric Acid; Third, Fourth, Fifth, Sixth, Seventh, Eighth, Ninth, and Tenth Courses—200 lbs. Bone-ash, and 150 lbs. Sulphuric Acid, per acre; Eleventh and Twelfth Courses—made from high percentage mineral phosphates, and containing 37 per cent., or more, of soluble phosphate.

(2) For the Tenth Course, in addition to the Superphosphate for the Swedish Turnips—300 lbs. Sulphate of Potash, 200 lbs. Sulphate of Soda, and 100 lbs. Sulphate of Magnesia were applied February 29, 1884, and harrowed in; and the same quantities were applied again before the final ploughing and preparation of the land for the sowing of the seed in May. For the Swedes of the Eleventh and Twelfth Courses the same mineral manures (which are the same as the mineral manures of the Third and subsequent Courses) were again applied, but only once for each of these two Courses. For the Swedes of the Thirteenth and Fourteenth Courses—500 lbs. Sulphate of Potash, 100 lbs. Sulphate of Soda, 200 lbs. Sulphate of Magnesia, and 600 lbs. Basic Slag, per acre.

(3) First Course—100 lbs. Bone-ash, 100 lbs. Sulphuric Acid, 100 lbs. Sulphate of Potash, 100 lbs. Sulphate of Soda, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake; Third, Fourth, Fifth, Sixth, Seventh, Eighth, Ninth, and Tenth Courses—300 lbs. Sulphate of Potash, 200 lbs. Sulphate of Soda, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake, per acre; Eleventh and Twelfth Courses—the same in other respects as in Courses 3-10, but the Superphosphate made from high percentage mineral phosphates, and containing 37 per cent., or more, of soluble phosphate.

(4) The quantities given in *Bushels* represent the *Dressed Corn* only.

(5) The "Total Produce" of the Corn-crops includes Dressed Corn, Offal Corn, Straw, and Chaff.

(6) Two cuttings.

[For Summary Table of the above results, see pp. 120-121.]

AGDELL FIELD.
(Area under experiment, about 3 acres.)

EXPERIMENTS ON AN ACTUAL COURSE OF ROTATION—TURNIPS, BARLEY, LEGUMINOUS CROP (OR FALLOW), AND WHEAT.

These Experiments were commenced in 1848; so that the present season, 1901, is the 54th, and the growing crop (Barley) is the second of the Fourteenth Course. One-third of the land has been continuously unmanured. One-third has, for the first Nine Courses, or 36 years, 1848-83, been manured with Superphosphate of Lime alone, once every four years, that is, for the turnip-crop commencing each course; but for the Tenth, Eleventh, Twelfth, Thirteenth, and Fourteenth Courses, a complex mineral manure has been applied, as described in foot-note, No. 2. Lastly, one-third has been manured (also for the turnip-crop only), with a complex mineral and Nitrogenous manure, as described in the foot-note, No. 3. From half of each of the three differently manured plots, the turnip-crops (roots and leaves) are removed; and on the other half they are either consumed on the land by sheep, or spread and ploughed in. In the case of all the other crops, the total produce is removed from the land. In the First Course, clover was sown over the whole of each of the three differently manured plots; but in each of the subsequent courses a leguminous crop was grown on only half of each of the three plots, the other half being left fallow, in the third year of each course. In the Second, Third, and Fourth Courses, clover was sown, but failed; and in them, and in the Fifth and Sixth Courses, beans were taken instead. In the Seventh Course, clover was sown (spring 1873), and gave three cuttings in 1874. In the Eighth Course beans were grown. In the Ninth Course clover was sown (in the spring of 1881), and gave two cuttings in 1882. In the Tenth Course clover was sown (in the spring of 1885), and yielded two cuttings in 1886. In the Eleventh Course clover was sown (with the barley) in 1889, but failed during the winter, and in 1890 beans were grown instead. In the Twelfth Course clover was again sown in April 1893, and gave two cuttings in 1894. In the Thirteenth Course clover was sown (with the barley), April 1897, but failed during the winter, and in 1898 beans were grown instead. In the Fourteenth Course clover was sown (with the barley), May 4, 1901.

TABLE III. (below), gives the results relating to the portions of each plot on which the turnip-crops were either fed off by sheep, or cut and spread on the land; and on which clover or beans were grown.

Years.	Description of Crop.	PRODUCE PER ACRE.														
		Plot 1. Unmanured continuously.					Plot 2. Superphosphate of Lime alone ⁽¹⁾ , Courses 1-9, Complex Mineral Manure ⁽²⁾ , Courses 10-14, for the Turnip Crops only.					Plot 3. Complex Mineral and Nitrogenous Manure ⁽³⁾ , for the Turnip Crops only.				
		Corn ⁽⁴⁾ (or Roots).	Straw (or Leaf).	Total Produce. ⁽⁵⁾	Corn ⁽⁴⁾ (or Roots).	Straw (or Leaf).	Total Produce. ⁽⁵⁾	Corn ⁽⁴⁾ (or Roots).	Straw (or Leaf).	Total Produce. ⁽⁵⁾	Corn ⁽⁴⁾ (or Roots).	Straw (or Leaf).	Total Produce. ⁽⁵⁾			
1848 1849 1850 1851	Norfolk White Turnips Barley Clover (called as hay) ⁽⁶⁾ Wheat	109 cwts. 48 bush. 39½ bush. 39½ bush.	67½ cwts. 3225 lbs. 3760 lbs.	1763 cwts. 6046 lbs. 5855 lbs.	229½ cwts. 42½ bush. 32 bush.	90 cwts. 3327 lbs. 4014 lbs.	310½ cwts. 5385 lbs. 6176 lbs.	229 cwts. 42½ bush. 31½ bush.	151½ cwts. 3646 lbs. 4035 lbs.	380½ cwts. 6206 lbs. 6169 lbs.	33 cwts. 2981 lbs. 1605 lbs. 4370 lbs.	419 cwts. 5190 lbs. 2544 lbs. 6992 lbs.				
1852 1853 1854 1855	Swedish Turnips Barley Beans Wheat	19½ cwts. 28½ bush. 5½ bush. 34½ bush.	3½ cwts. 2077 lbs. 953 lbs. 3351 lbs.	22½ cwts. 3817 lbs. 1367 lbs. 5526 lbs.	250½ cwts. 38 bush. 10½ bush. 36½ bush.	22 cwts. 2756 lbs. 1378 lbs. 3611 lbs.	272½ cwts. 5053 lbs. 2124 lbs. 5921 lbs.	386 cwts. 35½ bush. 138 bush. 40½ bush.	33 cwts. 2981 lbs. 1605 lbs. 4370 lbs.	419 cwts. 5190 lbs. 2544 lbs. 6992 lbs.	33 cwts. 2981 lbs. 1605 lbs. 4370 lbs.	419 cwts. 5190 lbs. 2544 lbs. 6992 lbs.				
1856 1857 1858 1859	Swedish Turnips Barley Beans Wheat	20½ cwts. 40½ bush. 5½ bush. 30½ bush.	1½ cwts. 2312 lbs. 965 lbs. 3355 lbs.	21½ cwts. 4558 lbs. 1307 lbs. 5265 lbs.	196 cwts. 52½ bush. 8½ bush. 37½ bush.	14½ cwts. 2730 lbs. 1320 lbs. 4320 lbs.	210½ cwts. 5741 lbs. 1895 lbs. 6659 lbs.	341½ cwts. 62½ bush. 143 bush. 39½ bush.	11½ cwts. 3405 lbs. 1760 lbs. 4955 lbs.	353 cwts. 6930 lbs. 2754 lbs. 7417 lbs.	11½ cwts. 3405 lbs. 1760 lbs. 4955 lbs.	353 cwts. 6930 lbs. 2754 lbs. 7417 lbs.				
1860 1861 1862 1863	Swedish Turnips Barley Beans Wheat	1 cwt. 1970 lbs. 27 bush. 50½ bush.	(5 lbs.) 1815 lbs. 3008 lbs.	1 cwt. 3525 lbs. 3546 lbs. 4941 lbs.	38½ cwts. 49½ bush. 30 bush. 41½ bush.	14 cwt. 2553 lbs. 2155 lbs. 3388 lbs.	40½ cwts. 4922 lbs. 4097 lbs. 6562 lbs.	72 cwts. 54½ bush. 41½ bush. 44½ bush.	4½ cwts. 3940 lbs. 2945 lbs. 4919 lbs.	76½ cwts. 7148 lbs. 5520 lbs. 7721 lbs.	4½ cwts. 3940 lbs. 2945 lbs. 4919 lbs.	76½ cwts. 7148 lbs. 5520 lbs. 7721 lbs.				
1864 1865 1866 1867	Swedish Turnips Barley Beans Wheat	8½ cwts. 27½ bush. 8½ bush. 15½ bush.	1 cwt. 1480 lbs. 905 lbs. 1524 lbs.	9½ cwts. 2961 lbs. 2481 lbs. 2506 lbs.	78½ cwts. 41½ bush. 10 bush. 25 bush.	44 cwts. 2244 lbs. 1835 lbs. 2646 lbs.	83½ cwts. 4457 lbs. 2481 lbs. 4242 lbs.	163½ cwts. 433 bush. 24½ bush. 21½ bush.	8½ cwts. 2958 lbs. 2155 lbs. 1654 lbs.	177½ cwts. 5308 lbs. 3732 lbs. 3023 lbs.	8½ cwts. 2958 lbs. 2155 lbs. 1654 lbs.	177½ cwts. 5308 lbs. 3732 lbs. 3023 lbs.				

1 lb. (pound avoird.) per acre = (about) 1.12 Kilogramme per Hectare, or 0.57 Zollverein Pfund. per Prussian Morgen.
1 cwt. (hundredweight) per acre = (about) 125.5 Kilogrammes per Hectare, or 0.64 Centner per Pr. Morgen.

Year	Course	Swedish Turnips	Failed, and ploughed up.	Failed, and ploughed up.	Failed, and ploughed up.
1868	6th Course, 1868-71	Swedish Turnips	25½ bush.	33½ bush.	42½ bush.
1869		Barley	1944 lbs.	2401 lbs.	3229 lbs.
1870		Beans	710 lbs.	878 lbs.	1068 lbs.
1871		Wheat	2855 lbs.	2980 lbs.	3644 lbs.
1872	7th Course, 1872-75	Swedish Turnips	294 cwts.	19½ cwts.	320 cwts.
1873		Barley	1495 lbs.	1841 lbs.	2456 lbs.
1874		Beans	775 lbs.	923 lbs.	1018 lbs.
1875		Wheat	2553 lbs.	3023 lbs.	4385 lbs.
1876	8th Course, 1876-79	Swedish Turnips	21 cwts.	225½ cwts.	359½ cwts.
1877		Barley	1341 lbs.	1994 lbs.	3125 lbs.
1878		Beans	775 lbs.	1350 lbs.	1880 lbs.
1879		Wheat	1219 lbs.	1771 lbs.	2138 lbs.
1880	9th Course, 1880-83	Swedish Turnips	21 cwts.	223½ cwts.	384 cwts.
1881		Barley	1468 lbs.	1430 lbs.	3078 lbs.
1882		Beans	775 lbs.	704 lbs.	834 lbs.
1883		Wheat	2060 lbs.	3275 lbs.	4505 lbs.
1884	10th Course, 1884-87	Swedish Turnips	12 cwts.	206 cwts.	280½ cwts.
1885		Barley	1379 lbs.	2353 lbs.	3386 lbs.
1886		Beans	775 lbs.	42 lbs.	44½ bush.
1887		Wheat	1844 lbs.	3468 lbs.	3645 lbs.
1888	11th Course, 1888-91	Swedish Turnips	8 cwts.	249½ cwts.	417½ cwts.
1889		Barley	865 lbs.	23 cwts.	2030 lbs.
1890		Beans	633 lbs.	1613 lbs.	25½ bush.
1891		Wheat	2318 lbs.	5017 lbs.	4309 lbs.
1892	12th Course, 1892-95	Swedish Turnips	6½ cwts.	254½ cwts.	332½ cwts.
1893		Barley	1358 lbs.	1466 lbs.	2100 lbs.
1894		Beans	775 lbs.	644 cwts.	81 cwts.
1895		Wheat	1619 lbs.	2831 lbs.	2100 lbs.
1896	13th Course, 1896-99	Swedish Turnips	11½ cwts.	249½ cwts.	313½ cwts.
1897		Barley	986 lbs.	2794 lbs.	3353 lbs.
1898		Beans	1323 lbs.	2144 lbs.	1548 lbs.
1899		Wheat	3181 lbs.	4404 lbs.	4509 lbs.
1900	14th Course, 1900-1903	Swedish Turnips	15½ cwts.	275½ cwts.	499½ cwts.
1901		Barley	196 cwts.	8 cwts.	153 cwts.
1902		Beans	775 lbs.	7134 lbs.	5742 lbs.
1903		Wheat	3181 lbs.	4404 lbs.	5071 lbs.

(1) First Course—100 lbs. Bone-ash, and 100 lbs. Sulphuric Acid (sp. gr. 1.7); Second Course—160 lbs. Bone-ash, 120 lbs. Sulphuric Acid, Third, Fourth, Fifth, Sixth, Seventh, Eighth, Ninth, and Tenth Courses—200 lbs. Bone-ash, and 150 lbs. Sulphuric Acid, per acre; Eleventh and Twelfth Courses—made from high percentage mineral phosphates, and containing 37 per cent., or more, of soluble phosphate.

(2) For the Tenth Course, in addition to the Superphosphate for the Swedish Turnips—300 lbs. Sulphate of Potash, 200 lbs. Sulphate of Soda, and 100 lbs. Sulphate of Magnesia were applied. February 29, 1884, and harrowed in; and the same quantities were applied again before the final ploughing and preparation of the land for the sowing of the seed in May. For the Swedes of the Eleventh and Twelfth Courses the same mineral manures (which are the same as the mineral manures of Plot 3 for the Third and subsequent Courses) were again applied, but only once for each of these two Courses. For the Swedes of the Thirteenth and Fourteenth Courses—500 lbs. Sulphate of Potash, 100 lbs. Sulphate of Soda, 200 lbs. Sulphate of Magnesia, and 600 lbs. Basic Sluag, per acre.

(3) First Course—100 lbs. Peat-ash, 100 lbs. Bone-ash, 100 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 1000 lbs. Rape-cake; Second Course—300 lbs. Sulphate of Potash, 100 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 160 lbs. Bone-ash, 120 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake; Third, Fourth, Fifth, Sixth, Seventh, Eighth, Ninth, and Tenth Courses—300 lbs. Sulphate of Potash, 200 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 3000 lbs. Rape-cake, per acre; Eleventh and Twelfth Courses—the same in other respects as in Courses 3-10, but the Superphosphate made from high percentage mineral phosphates, and containing 37 per cent. or more, of soluble phosphate. For the Swedes of the Thirteenth and Fourteenth Courses—500 lbs. Sulphate of Potash, 100 lbs. Sulphate of Soda, 200 lbs. Sulphate of Magnesia, 600 lbs. Basic Sluag, 2000 lbs. Rape-cake, 100 lbs. Sulphate of Ammonia, and 100 lbs. Muriate of Ammonia, per acre.

(4) The quantities given in *italics* represent the *Present* Corn only.

(5) The "Total Produce" of the Corn-crops includes Dressed Corn, Oat Corn, Straw, and Chaff.

(6) Two cuttings.

(7) Three cuttings.

[For Summary Table of the above results, see pp. 120-121.]

AGDELL FIELD.

(Area under experiment, about 3 acres.)

EXPERIMENTS ON AN ACTUAL COURSE OF ROTATION—TURNIPS, BARLEY, LEGUMINOUS CROP (OR FALLOW), AND WHEAT.

These Experiments were commenced in 1843; so that the present season, 1901, is the 54th, and the growing crop (Barley) is the second of the Fourteenth Course.

One-third of the land has been continuously unmanured. One-third has, for the first Nine Courses, or 36 years, 1843-83, been manured with Superphosphate of Lime alone, once every four years, that is for the turnip-crop commencing each course; but for the Tenth, Eleventh, Twelfth, Thirteenth, and Fourteenth Courses, a complex mineral manure has been applied, as described in foot-note, No. 2. Lastly, one-third has been manured (also for the turnip-crop only), with a complex mineral and Nitrogenous manure, as described in the foot-note, No. 3.

From half of each of the three differently manured plots, the turnip-crops (roots and leaves) are removed; and on the other half they are either consumed on the land by sheep, or spread and ploughed in. In the case of all the other crops, the total produce is removed from the land.

In the First Course, clover was sown over the whole of each of the three differently manured

plots; but in each of the subsequent courses, a leguminous crop was grown on only half of each of the three plots, the other half being left fallow, in the third year of each course. In the Second, Third, and Fourth Courses, clover was sown, but failed; and in them, and in the Fifth and Sixth Courses, beans were taken instead. In the Seventh Course, clover was sown (spring 1875), and gave three cuttings in 1874. In the Eighth Course beans were grown. In the Ninth Course clover was sown (in the spring of 1881), and gave two cuttings in 1882. In the Tenth Course clover was sown (in the spring of 1885), and yielded two cuttings in 1886. In the Eleventh Course clover was sown (with the barley), in 1889, but failed during the winter, and in 1890 beans were grown instead. In the Twelfth Course clover was again sown in April 1893, and gave two cuttings in 1894. In the Thirteenth Course clover was sown (with the barley), April 1897, but failed during the winter, and in 1898 beans were grown instead. In the Fourteenth Course clover was sown (with the barley), May 4, 1901.

TABLE IV. (below), gives the results relating to the portions of each plot on which the turnip-crops were either fed off by sheep, or cut and spread on the land; and on which, in the third year of each course (excepting the first, 1850, when clover was grown), the land was left fallow.

1 lb. (pound avoird.) per acre = (about) 1.12 Kilogramme per Hectare, or 0.57 Zollverein Pfund. per Prussian Morgen.
 1 cwt. (hundred weight) per acre = (about) 125.5 Kilogrammes per Hectare, or 0.64 Centner per Pr. Morgen.

Years.	Description of Crop.	PRODUCE PER ACRE.														
		PLOT 1. Unmanured continuously.					PLOT 2. Superphosphate of Lime alone, (1) Courses 1-9, Complex Mineral Manures (2), Courses 10-14; for the Turnip Crops only.					PLOT 3. Complex Mineral and Nitrogenous Manure, (3) for the Turnip Crops only.				
		Corn (1) (or Roots).	Straw (or Leaf).	Total Produce (2)	Corn (4) (or Roots).	Straw (or Leaf).	Total Produce (5)	Corn (4) (or Roots).	Straw (or Leaf).	Total Produce (6)	Corn (4) (or Roots).	Straw (or Leaf).	Total Produce (7)			
1848	Swedish Turnips	17½ cwt.	20½ cwt.	193½ cwt.	345 cwt.	38½ cwt.	384½ cwt.	429 cwt.	46½ cwt.	475½ cwt.	429 cwt.	46½ cwt.	475½ cwt.			
1849	Barley	44½ bush.	3139 lbs.	5785 lbs.	41 bush.	3209 lbs.	5704 lbs.	44½ bush.	3709 lbs.	6344 lbs.	44½ bush.	3709 lbs.	6344 lbs.			
1850	Clover (calc ^d as hay) (1)	31½ bush.	3498 lbs.	624 cwt.	35½ bush.	3834 lbs.	6962 lbs.	27½ bush.	3969 lbs.	65 cwt.	27½ bush.	3969 lbs.	65 cwt.			
1851	Wheat	27½ cwt.	4 cwt.	314 cwt.	273½ cwt.	22½ cwt.	295½ cwt.	390½ cwt.	37½ cwt.	423½ cwt.	390½ cwt.	37½ cwt.	423½ cwt.			
1852	Swedish Turnips	33 bush.	2210 lbs.	4161 lbs.	39½ bush.	2729 lbs.	5110 lbs.	37½ bush.	3323 lbs.	5672 lbs.	37½ bush.	3323 lbs.	5672 lbs.			
1853	Barley	37½ bush.	4070 lbs.	6473 lbs.	37½ bush.	4492 lbs.	6961 lbs.	37½ bush.	5107 lbs.	7499 lbs.	37½ bush.	5107 lbs.	7499 lbs.			
1854	Fallow	34 cwt.	2 cwt.	36 cwt.	193½ cwt.	124 cwt.	206 cwt.	3394 cwt.	124 cwt.	351½ cwt.	3394 cwt.	124 cwt.	351½ cwt.			
1855	Wheat	44½ bush.	2430 lbs.	4912 lbs.	43½ bush.	2595 lbs.	5326 lbs.	68½ bush.	3570 lbs.	7261 lbs.	68½ bush.	3570 lbs.	7261 lbs.			
1856	Swedish Turnips	33½ bush.	4045 lbs.	6270 lbs.	39½ bush.	4720 lbs.	7242 lbs.	40½ bush.	5545 lbs.	8136 lbs.	40½ bush.	5545 lbs.	8136 lbs.			
1857	Barley	14 cwt.	1 cwt.	15 cwt.	40½ cwt.	2 cwt.	42½ cwt.	87 cwt.	57½ cwt.	92½ cwt.	87 cwt.	57½ cwt.	92½ cwt.			
1858	Fallow	33 bush.	2013 lbs.	3871 lbs.	40½ bush.	2475 lbs.	4803 lbs.	57½ bush.	4175 lbs.	7554 lbs.	57½ bush.	4175 lbs.	7554 lbs.			
1859	Wheat	42 bush.	4295 lbs.	6989 lbs.	49½ bush.	5051 lbs.	8194 lbs.	49 bush.	5638 lbs.	8747 lbs.	49 bush.	5638 lbs.	8747 lbs.			
1860	Swedish Turnips	9 cwt.	3 cwt.	12 cwt.	79½ cwt.	5½ cwt.	84½ cwt.	135½ cwt.	94 cwt.	195 cwt.	135½ cwt.	94 cwt.	195 cwt.			
1861	Barley	35½ bush.	1809 lbs.	3695 lbs.	39½ bush.	2043 lbs.	4122 lbs.	46½ bush.	3274 lbs.	5753 lbs.	46½ bush.	3274 lbs.	5753 lbs.			
1862	Fallow	29½ bush.	2638 lbs.	4126 lbs.	27½ bush.	2989 lbs.	4702 lbs.	19½ bush.	2905 lbs.	4180 lbs.	19½ bush.	2905 lbs.	4180 lbs.			
1863	Wheat	29½ bush.	2638 lbs.	4126 lbs.	27½ bush.	2989 lbs.	4702 lbs.	19½ bush.	2905 lbs.	4180 lbs.	19½ bush.	2905 lbs.	4180 lbs.			

Year	Crop	Failed and ploughed up.	Failed and ploughed up.	Failed and ploughed up.	Failed and ploughed up.
1868	Swedish Turnips	21 bush.	1648 lbs.	2343 lbs.	38 3/4 bush.
1869	Barley	14 1/2 bush.	1946 lbs.	2840 lbs.	3244 lbs.
1870	Fallow	49 1/2 cwt.	7 1/2 cwt.	56 1/2 cwt.	2863 lbs.
1871	Wheat	20 1/2 bush.	1311 lbs.	2536 lbs.	17 1/2 bush.
1872	Swedish Turnips	24 1/2 bush.	2851 lbs.	4396 lbs.	331 1/2 cwt.
1873	Barley	32 1/2 cwt.	5 1/2 cwt.	37 1/2 cwt.	47 cwt.
1874	Fallow	22 1/2 bush.	1275 lbs.	2609 lbs.	2796 lbs.
1875	Wheat	11 1/2 bush.	1612 lbs.	2351 lbs.	4085 lbs.
1876	Swedish Turnips	38 1/2 cwt.	4 cwt.	42 1/2 cwt.	30 bush.
1877	Barley	31 1/2 bush.	1668 lbs.	3997 lbs.	37 1/2 cwt.
1878	Fallow	34 1/2 bush.	3231 lbs.	5445 lbs.	44 1/2 bush.
1879	Wheat	20 1/2 cwt.	7 cwt.	27 1/2 cwt.	10 1/2 bush.
1880	Swedish Turnips	23 cwt.	7 1/2 cwt.	30 1/2 cwt.	44 1/2 cwt.
1881	Barley	16 1/2 bush.	996 lbs.	1898 lbs.	47 1/2 bush.
1882	Fallow	31 1/2 bush.	2398 lbs.	4763 lbs.	38 cwt.
1883	Wheat	12 1/2 cwt.	1 cwt.	13 1/2 cwt.	2993 lbs.
1884	Swedish Turnips	19 bush.	1639 lbs.	2768 lbs.	4028 lbs.
1885	Barley	22 1/2 bush.	3231 lbs.	5445 lbs.	41 bush.
1886	Fallow	33 1/2 bush.	3231 lbs.	5445 lbs.	66 1/2 cwt.
1887	Wheat	23 cwt.	7 1/2 cwt.	30 1/2 cwt.	2778 lbs.
1888	Swedish Turnips	16 1/2 bush.	996 lbs.	1898 lbs.	3763 lbs.
1889	Barley	31 1/2 bush.	2398 lbs.	4763 lbs.	36 1/2 cwt.
1890	Fallow	12 1/2 cwt.	1 cwt.	13 1/2 cwt.	4624 lbs.
1891	Wheat	19 bush.	1639 lbs.	2768 lbs.	6410 lbs.
1892	Swedish Turnips	22 1/2 bush.	3231 lbs.	5445 lbs.	45 1/2 cwt.
1893	Barley	33 1/2 bush.	3231 lbs.	5445 lbs.	35 cwt.
1894	Fallow	23 cwt.	7 1/2 cwt.	30 1/2 cwt.	1776 lbs.
1895	Wheat	16 1/2 bush.	996 lbs.	1898 lbs.	4938 lbs.
1896	Swedish Turnips	12 1/2 cwt.	1 cwt.	13 1/2 cwt.	500 1/2 cwt.
1897	Barley	19 bush.	1639 lbs.	2768 lbs.	11 1/2 cwt.
1898	Fallow	22 1/2 bush.	3231 lbs.	5445 lbs.	1979 lbs.
1899	Wheat	23 cwt.	7 1/2 cwt.	30 1/2 cwt.	257 1/2 lbs.
1900	Swedish Turnips	24 1/2 cwt.	4 cwt.	28 1/2 cwt.	32 1/2 bush.
1901	Barley	13 1/2 bush.	1945 lbs.	2840 lbs.	33 1/2 cwt.
1902	Fallow	27 1/2 bush.	3050 lbs.	4778 lbs.	35 1/2 bush.
1903	Wheat	45 1/2 cwt.	5 1/2 cwt.	64 1/2 cwt.	39 bush.

(1) First Course—100 lbs. Bone-ash, and 100 lbs. Sulphuric Acid (sp. gr. 1.7); Second Course—1600 lbs. Bone-ash, 120 lbs. Sulphuric Acid; Third, Fourth, Fifth, Sixth, Seventh, Eighth, Ninth, and Tenth Courses—200 lbs. Bone-ash, and 150 lbs. Sulphuric Acid, per acre; Eleventh and Twelfth Courses—made from high percentage mineral phosphates, and containing 37 per cent., or more, of soluble phosphate.

(2) For the Tenth Course, in addition to the Superphosphate for the Swedish Turnips—300 lbs. Sulphate Potash, 200 lbs. Sulphate Soda, and 100 lbs. Sulphate Magnesia were applied February 29, 1884, and harrowed in; and the same quantities were applied again before the final ploughing and preparation of the land for the sowing of the seed in May. For the Swedes of the Eleventh and Twelfth Courses the same mineral manures (which are the same as the mineral manures of Plot 3 for the Third and subsequent Courses) were again applied, but only once for each of these two Courses. For the Swedes of the Thirteenth and Fourteenth Courses—500 lbs. Sulphate of Potash, 100 lbs. Sulphate of Soda, 200 lbs. Sulphate of Magnesia, and 600 lbs. Basic Slag, per acre.

(3) First Course—100 lbs. Pearl-ash, 100 lbs. Bone-ash, 100 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, and 100 lbs. Muriate of Ammonia, and 1000 lbs. Rape-cake; Second Course—300 lbs. Sulphate of Potash, 100 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 160 lbs. Bone-ash, 120 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake; Third, Fourth, Fifth, Sixth, Seventh, Eighth, Ninth, and Tenth Courses—300 lbs. Sulphate of Potash, 200 lbs. Sulphate of Soda, 100 lbs. Sulphate of Magnesia, 200 lbs. Bone-ash, 150 lbs. Sulphuric Acid, 100 lbs. Sulphate of Ammonia, 100 lbs. Muriate of Ammonia, and 2000 lbs. Rape-cake, per acre; Eleventh and Twelfth Courses—the same in other respects as in Courses 3-10, but the Superphosphate cake, per cent. high percentage mineral phosphates, and containing 37 per cent., or more, of soluble phosphate. For the Swedes of the Thirteenth and Fourteenth Courses—500 lbs. Sulphate of Potash, 100 lbs. Sulphate of Soda, 200 lbs. Sulphate of Magnesia, 600 lbs. Basic Slag, 2000 lbs. Rape-cake, 100 lbs. Sulphate of Ammonia, and 100 lbs. Muriate of Ammonia, per acre.

(4) The quantities given in *Bushels* represent the *Dressed Corn* only.

(5) The "Total Produce" of the Corn-crops includes Dressed Corn, Offal Corn, Straw, and Chaff.

(6) Two cuttings.

[For Summary Table of the above results, see pp. 120-121.]

AGDELL FIELD.

(Area under experiment, about 3 acres.)

EXPERIMENTS ON AN ACTUAL COURSE OF ROTATION—TURNIPS, BARLEY, LEGUMINOUS CROP (OR FALLOW), AND WHEAT.

SUMMARIES OF THE RESULTS GIVEN IN TABLES I, II, III, AND IV. (pp. 112-13, 114-15, 116-17, and 118-19), RESPECTIVELY.

As the Table shows, averages are given for each of the four portions of the experimental land, for which Tables I, II, III, and IV, respectively, give the details. The averages are given first of the produce of the eight intermediate Courses (Courses 2-9, 1852-1883); that is, excluding the First Course, when the land was in somewhat uneven condition, and when (as the detailed Tables show), on some portions Norfolk Whites, and on others Swedish Turnips, were grown; excluding also the Tenth, Eleventh, Twelfth, and

Thirteenth Courses, on account of the change in the Mineral Manures used on Plot 2. Averages are also given of the produce of the Tenth, Eleventh, Twelfth, and Thirteenth Courses, that is, after the change in the Mineral Manures applied to Plot 2. For full particulars of the manures applied to Plot 2, and also of those applied to Plot 3, see Foot-notes 1, 2, and 3, on pages 113, 115, 117, or 119.

1 lb. (pound avoird.) per acre = (about) 1.12 Kilogramme per Hectare, or 0.57 Zollverein Pfund, per Prussian Morgen.
 1 cwt. (hundredweight) per acre = (about) 125.5 Kilogrammes per Hectare, or 0.64 Centner per Pr. Morgen.

Years.	Description of Crop.	PRODUCE PER ACRE.														
		PLOT 1. Unmanured continuously.					PLOT 2. Superphosphate of Lime, alone, Courses 1-9, Complex Mineral Manure, Courses 10-13, for the Turnip Crops only.					PLOT 3. Complex Mineral and Nitrogenous Manure, for the Turnip Crops only.				
		Corn (1) (or Roots).	Straw (or Leaf).	Total Produce. (2)	Corn (1) (or Roots).	Straw (or Leaf).	Total Produce. (2)	Corn (1) (or Roots).	Straw (or Leaf).	Total Produce. (2)	Corn (1) (or Roots).	Straw (or Leaf).	Total Produce. (2)			
AVERAGE OF 8 COURSES (COURSES 2-9), 1852-1883.																
1852, '56, '60, '64, '72, '76, '80	Swedish Turnips	16½ cwt.	3 cwt.	19½ cwt.	126½ cwt.	11½ cwt.	138½ cwt.	266½ cwt.	24½ cwt.	290½ cwt.	206½ cwt.	24½ cwt.	290½ cwt.			
1853, '57, '61, '65, '69, '73, '77, '81	Barley	32½ bush.	1971 lbs.	3790 lbs.	27½ bush.	1623 lbs.	3196 lbs.	42½ bush.	2547 lbs.	4962 lbs.	42½ bush.	2547 lbs.	4962 lbs.			
1854, '58, '62, '66, '70, '74, '78, '82	{ Clover, 1874, and '82 (as hay)	12½ bush.	1081 lbs.	1887 lbs.	12½ bush.	1200 lbs.	1996 lbs.	21½ bush.	1809 lbs.	3230 lbs.	21½ bush.	1809 lbs.	3230 lbs.			
1855, '59, '63, '67, '71, '75, '79, '83	{ Beans	26 bush.	2762 lbs.	4407 lbs.	25½ bush.	3023 lbs.	4841 lbs.	32½ bush.	3753 lbs.	5847 lbs.	32½ bush.	3753 lbs.	5847 lbs.			
	{ Wheat															
AVERAGE OF 4 COURSES (COURSES 10-13), 1884-1899.																
1884, 1888, 1892 and 1896	Swedish Turnips	53 cwt.	1½ cwt.	7½ cwt.	199½ cwt.	15 cwt.	214½ cwt.	393½ cwt.	43½ cwt.	437½ cwt.	393½ cwt.	43½ cwt.	437½ cwt.			
1885, 1889, 1893 and 1897	Barley	12½ bush.	1223 lbs.	1963 lbs.	20 bush.	1448 lbs.	2575 lbs.	25 bush.	2028 lbs.	3633 lbs.	25 bush.	2028 lbs.	3633 lbs.			
1886, 1890, 1894 and 1898	{ Clover, 1886 and 1894 (as hay)	15½ bush.	971 lbs.	2028 lbs.	28 bush.	1894 lbs.	3799 lbs.	19½ bush.	1373 lbs.	2609 lbs.	19½ bush.	1373 lbs.	2609 lbs.			
1887, 1891, 1895 and 1899	{ Beans, 1890 and 1895	27½ bush.	2372 lbs.	4096 lbs.	40½ bush.	3559 lbs.	6104 lbs.	42½ bush.	3749 lbs.	6350 lbs.	42½ bush.	3749 lbs.	6350 lbs.			
	{ Wheat															

SUMMARY OF TABLE I. (pp. 112-13).—Results relating to the portions of each plot from which the turnip-crops were entirely removed; and on which clover or beans were grown.

SUMMARY OF TABLE II. (pp. 114-15).—Results relating to the portions of each plot from which the turnip-crops were entirely removed; and on which, in the third year of each course (excepting the first, 1850, when clover was grown), the land was left fallow.

AVERAGE OF 8 COURSES (COURSES 2-9), 1852-1883.										
	1852, '56, '60, '64, '72, '76, '80	1853, '57, '61, '65, '69, '73, '77, '81	1854, '58, '62, '66, '70, '74, '78, '82	1855, '59, '63, '67, '71, '75, '79, '83	1852, '56, '60, '64, '72, '76, '80	1853, '57, '61, '65, '69, '73, '77, '81	1854, '58, '62, '66, '70, '74, '78, '82	1855, '59, '63, '67, '71, '75, '79, '83	1852, '56, '60, '64, '72, '76, '80	1853, '57, '61, '65, '69, '73, '77, '81
Swedish Turnips	26 cwt.	34 cwt.	29 cwt.	134 cwt.	10 cwt.	144 cwt.	26 cwt.	34 cwt.	29 cwt.	134 cwt.
Barley	30 bush.	1792 lbs.	3497 lbs.	27 cwt.	1568 lbs.	3121 lbs.	30 bush.	1792 lbs.	3497 lbs.	27 cwt.
Fallow	28 cwt.	3153 lbs.	4976 lbs.	30 cwt.	3383 lbs.	5348 lbs.	28 cwt.	3153 lbs.	4976 lbs.	30 cwt.
Wheat	28 cwt.	3153 lbs.	4976 lbs.	30 cwt.	3383 lbs.	5348 lbs.	28 cwt.	3153 lbs.	4976 lbs.	30 cwt.
AVERAGE OF 4 COURSES (COURSES 10-13), 1884-1899.										
Swedish Turnips	14 cwt.	44 cwt.	19 cwt.	172 cwt.	11 cwt.	184 cwt.	14 cwt.	44 cwt.	19 cwt.	172 cwt.
Barley	15 cwt.	1257 lbs.	2146 lbs.	13 cwt.	1045 lbs.	1521 lbs.	15 cwt.	1257 lbs.	2146 lbs.	13 cwt.
Fallow	25 cwt.	2539 lbs.	4352 lbs.	34 bush.	3243 lbs.	5383 lbs.	25 cwt.	2539 lbs.	4352 lbs.	34 bush.
Wheat	25 cwt.	2539 lbs.	4352 lbs.	34 bush.	3243 lbs.	5383 lbs.	25 cwt.	2539 lbs.	4352 lbs.	34 bush.

SUMMARY OF TABLE III. (pp. 116-17).—Results relating to the portions of each plot on which the turnip-crops were either fed off by sheep, or cut and spread on the land; and on which clover or beans were grown.

AVERAGE OF 8 COURSES (COURSES 2-9), 1852-1883.										
	1852, '56, '60, '64, '72, '76, '80	1853, '57, '61, '65, '69, '73, '77, '81	1854, '58, '62, '66, '70, '74, '78, '82	1855, '59, '63, '67, '71, '75, '79, '83	1852, '56, '60, '64, '72, '76, '80	1853, '57, '61, '65, '69, '73, '77, '81	1854, '58, '62, '66, '70, '74, '78, '82	1855, '59, '63, '67, '71, '75, '79, '83	1852, '56, '60, '64, '72, '76, '80	1853, '57, '61, '65, '69, '73, '77, '81
Swedish Turnips	15 cwt.	24 cwt.	17 cwt.	150 cwt.	12 cwt.	163 cwt.	15 cwt.	24 cwt.	17 cwt.	150 cwt.
Barley	29 bush.	1758 lbs.	3351 lbs.	38 bush.	2250 lbs.	4417 lbs.	29 bush.	1758 lbs.	3351 lbs.	38 bush.
Fallow	12 bush.	1026 lbs.	1807 lbs.	14 cwt.	1486 lbs.	2439 lbs.	12 bush.	1026 lbs.	1807 lbs.	14 cwt.
Wheat	23 cwt.	2441 lbs.	3927 lbs.	31 cwt.	3303 lbs.	5307 lbs.	23 cwt.	2441 lbs.	3927 lbs.	31 cwt.
AVERAGE OF 4 COURSES (COURSES 10-13), 1884-1899.										
Swedish Turnips	9 cwt.	27 cwt.	12 cwt.	237 cwt.	17 cwt.	254 cwt.	9 cwt.	27 cwt.	12 cwt.	237 cwt.
Barley	13 cwt.	1147 lbs.	1917 lbs.	29 cwt.	2059 lbs.	3760 lbs.	13 cwt.	1147 lbs.	1917 lbs.	29 cwt.
Fallow	16 bush.	979 lbs.	2067 lbs.	24 cwt.	1887 lbs.	3318 lbs.	16 bush.	979 lbs.	2067 lbs.	24 cwt.
Wheat	26 cwt.	2241 lbs.	3919 lbs.	44 cwt.	3930 lbs.	6731 lbs.	26 cwt.	2241 lbs.	3919 lbs.	44 cwt.

SUMMARY OF TABLE IV. (pp. 118-19).—Results relating to the portions of each plot on which the turnip-crops were either fed off by sheep, or cut and spread on the land; and on which, in the third year of each course (excepting the first, 1850, when clover was grown), the land was left fallow.

AVERAGE OF 8 COURSES (COURSES 2-9), 1852-1883.										
	1852, '56, '60, '64, '72, '76, '80	1853, '57, '61, '65, '69, '73, '77, '81	1854, '58, '62, '66, '70, '74, '78, '82	1855, '59, '63, '67, '71, '75, '79, '83	1852, '56, '60, '64, '72, '76, '80	1853, '57, '61, '65, '69, '73, '77, '81	1854, '58, '62, '66, '70, '74, '78, '82	1855, '59, '63, '67, '71, '75, '79, '83	1852, '56, '60, '64, '72, '76, '80	1853, '57, '61, '65, '69, '73, '77, '81
Swedish Turnips	24 cwt.	2 cwt.	2 cwt.	150 cwt.	11 cwt.	161 cwt.	24 cwt.	2 cwt.	2 cwt.	150 cwt.
Barley	30 cwt.	1784 lbs.	3491 lbs.	38 cwt.	2116 lbs.	4148 lbs.	30 cwt.	1784 lbs.	3491 lbs.	38 cwt.
Fallow	27 cwt.	3081 lbs.	4863 lbs.	31 cwt.	3621 lbs.	5659 lbs.	27 cwt.	3081 lbs.	4863 lbs.	31 cwt.
Wheat	27 cwt.	3081 lbs.	4863 lbs.	31 cwt.	3621 lbs.	5659 lbs.	27 cwt.	3081 lbs.	4863 lbs.	31 cwt.
AVERAGE OF 4 COURSES (COURSES 10-13), 1884-1899.										
Swedish Turnips	20 cwt.	4 cwt.	24 cwt.	194 cwt.	12 cwt.	207 cwt.	20 cwt.	4 cwt.	24 cwt.	194 cwt.
Barley	17 cwt.	1390 lbs.	2414 lbs.	18 bush.	1321 lbs.	2379 lbs.	17 cwt.	1390 lbs.	2414 lbs.	18 bush.
Fallow	23 cwt.	2543 lbs.	4337 lbs.	36 bush.	3474 lbs.	5753 lbs.	23 cwt.	2543 lbs.	4337 lbs.	36 bush.
Wheat	23 cwt.	2543 lbs.	4337 lbs.	36 bush.	3474 lbs.	5753 lbs.	23 cwt.	2543 lbs.	4337 lbs.	36 bush.

(1) The quantities given in *Bushels* represent the *Dressed* Corn only. (2) The "Total Produce" of the Corn-crops includes Dressed Corn, Oat Corn, Straw, and Chaff.