Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1901



Full Table of Content


Summary of Rainfall and Drainage at Rothamsted

Rothamsted Research

Rothamsted Research (1902) *Summary of Rainfall and Drainage at Rothamsted*; Yields Of The Field Experiments 1901, pp 16 - 19 - **DOI:** https://doi.org/10.23637/ERADOC-1-229

16)

THE ROTHAMSTED RAIN GAUGES.



VIEW SHOWING THE COLLECTORS.

No. 1.—Small Funnel-gauge, 5 inches diameter. No. 2.—Small Funnel-gauge, 8 inches diameter.

No. 2.—Small Funnel-gauge, 8 inches diameter.

No. 3.—Large Gauge—

Size — 7 feet 3·12 in. × 6 feet.

Area—One thousandth of an acre.

4 collectors, each holding Rain = 0·500 in.

Gauge-tubes graduated to ... 0·002 in.


Overflow tank to hold Rain = 2·000 ins.

Small cylinder, tube graduated to 0·001 in.

(For quantities less than 0·05 in.)

No. 4.—Stand with level marble top, for measuring.

THE ROTHAMSTED DRAIN GAUGES.



2.000 ins.

(17)

GENERAL SUMMARY OF THE RESULTS RELATING TO RAINFALL AND DRAINAGE AT ROTHAMSTED.

As already explained (p. 6), the Rainfall has been measured at Rothamsted, in gauges of different sizes almost from the commencement of the Field Experiments; and the Drainage through 20 inches, 40 inches, and 60 inches, of unmanured and uncropped soil, in its natural state of consolidation, has been collected from September 1870, up to the present time, a period of nearly 31 Harvest-years. The nitrogen as nitric acid in the drainage waters was determined occasionally during the first 7 years of their collection (1870-1 to 1876-7); but from 1877-8 up to the present time, a period of nearly 24 Harvest-years, it has been determined in proportional samples for each month. A summary of the numerical results relating to rainfall, drainage, and loss of nitrogen by the drainage, will be found in the Tables at pp. 18-19.

The figures show that over the 20 Harvest-years, 1877-8 to 1896-7, there was an average annual loss of nitrogen in the drainage, of 35 07 lbs. through 20 inches, 30 83 lbs. through 40 inches, and 33 87 lbs. through 60 inches depth, of unmanured and uncropped soil; or, taking the average of the three gauges, of about 33 lb. per acre per annum of nitrogen, corresponding to an average annual loss of nearly 2 cwts. of nitrate of soda.

With reference to the large amounts of loss of nitrogen by the drainage thus indicated, it is to be borne in mind, that probably not more than an average of about 5 lbs. would be contributed annually per acre from the atmosphere in rain and the minor aqueous deposits, exclusive of any condensation by the soil. Moreover, the soil of the drain-gauges had been unmanured since 1868; but, being exposed to the access of air from below as well as from above, the oxidation of the nitrogenous matters of the soil and subsoil may be more active than in similar soil in its natural condition. On the other hand, the soil was without any vegetation to arrest the nitric acid formed; whilst, in some cases of fallow-land which had been manured and cropped in the ordinary course, and which would therefore be in a much higher "condition" than the soil of the drain-gauges, as much, or even more nitrogen as nitric acid, has been found to be present in the autumn, to the depth of 18 or 27 inches.

As the three drain-gauges of unmanured and uncropped soil have been in operation since September 1870 that is, for 30 Harvest-years to the end of August 1900-it is obviously of interest to consider whether or not there is evidence of decline in the amount of nitric acid annually formed in the soil over that period. As the drainage waters were not systematically sampled and analysed until 1877, accurate estimates of the amount of nitric acid in the drainage of the first 7 years of the experiments cannot be made; but the indication is, that it averaged more over those earlier than over the subsequent years. Examination of the results for the four 5-yearly periods of the 20 years, 1877-8 to 1896-7, shows that the average annual loss of nitrogen per acre in the drainage was, over the first 5 years considerably more than the average of the 20 years, but over the second 5 it was less than the average, and over the third 5 less than over the second; whilst over the last five years (1892-3 to 1896-7) the loss somewhat exceeded that of the preceding five years. The details show, that the excess over the first 5 years was due to very excessive amounts in the second and fourth years of the twenty; and that these were coincident with very excessive amounts of rain and of drainage. The loss over the second 5 years was considerably less than over the first, and somewhat lower than the average, and so also was the amount of rain, though the drainage was, on the average, slightly higher. The loss over the third 5 years was less than over the second, and less than the average, and so also were the average amounts of rain and drainage. Over the last 5 years again, the distinctly lower than average loss of nitrogen in the drainage, was coincident with lower than average rainfall, and generally lower than average drainage, especially in the first and fourth years.

With this general decline in the loss of nitrogen in the later years, there was, on the other hand, in the 8th year of the 20 considerably more than the average loss, in the 9th about the average, and in the 11th, 17th and 20th years considerably more, and in the 18th year rather more than the average; part of the excess in the 17th year (1893-4) being probably due to accumulation during the preceding year of drought. In each of the last four years of the first 15, however, the loss was considerably below the average, and in the 16th year, a year of drought (1892-3), there was, with nearly the lowest rainfall and drainage, also nearly the lowest loss of nitrogen by drainage. Lastly as to the details:—in October of the fifteenth Harvest-year (1891-2), there was, with a great excess of rain and drainage, very much more than the average amount of nitric acid in the drainage of that month. Indeed, in only one month during the preceding 14 years, namely in September 1880, was there more nitric acid passing through the 20-inch gauge than in October 1891. There was, however, in October of the 17th year, 1893-4, and in November of the 19th year, 1895-6, about as much loss of nitrogen as nitric acid as in October 1891. More pertinent still is the fact, that in September of the Harvest-year, 1896-7, which is the 27th of the parallel rain and drainage records, and the 20th of the monthly analysis of the drainage waters, there was, with the heaviest rainfall and drainage ever recorded at Rothamsted for a single month, also the greatest loss of nitrogen as nitric acid by the drainage, in any one month of the 20 years, and this was the case with the 20-, the 40-, and the 60-inch drain-gauges.

Obviously, the loss of nitrogen as nitric acid through the soil-drain-gauges was very directly dependent on the amount, and on the distribution, of the rain, and of the drainage. Indeed, although there seems to be some indication of a decline in the amount of nitric acid formed over the later than over the earlier years, the evidence is certainly not yet very conclusive. Thus, examination of the figures shows, that the loss of nitrogen was considerably higher in the 20th Harvest-year, 1896-7, than the average of the 20 years. On the other hand, it was actually lower in the 21st year, 1897-8, than in any of the preceding 20 years. In 1898-9 it was rather under, and in 1899-1900 over average. During the 9 months of the current Harvest-year, ending with May 1901, it was about average.

Finally, according to a revised estimate for the 30 Harvest-years ending with August 31, 1900, there has been a loss of nitrogen through the 20-inch gauge of nearly 17, through the 40-inch of 9, and through the 60-inch of more than $7\frac{1}{2}$ per cent., of the total combined nitrogen in the soil and subsoil.

C

(18)

NUMERICAL SUMMARY OF RESULTS RELATING TO RAINFALL AND DRAINAGE AT ROTHAMSTED.

In the Tables (pp. 18-19), are given—the average annual Rainfall over the 19 Harvest-years, 1851-2 to 1869-70; that is, from the commencement of the period for which the average produce, of continuous wheat (pp. 30-1), and of continuous barley (pp. 26-7), is given, up to the time when the experiments with the drain-gauges commenced. Next are given the averages of both rainfall and drainage for the first 7 Harvest-years during which drainage as well as rainfall was collected (1870-1 to 1876-7). Up to that time, the nitric acid in the drainage water had only been occasionally determined; but from 1877-8, up to the present time, it has been determined in proportionally mixed samples for each month; and the Table shows, in inches, the rainfall, and the drainage through each of the three soil-drain-gauges, respectively of 20, 40, and 60 inches depth of soil; also the difference (rainfall collected in the large gauge, \(\frac{1}{10\text{col}}\) acrea area, \(\text{minus}\) drainage), approximately representing evaporation. The Table further shows—the amounts of loss of nitrogen, in 1b, per acre, in the drainage through each of the three drain-gauges, reckoned both as nitrogen, and as nitrate of soda (commercial, 5 per cent. impurity), for each of the 23 Harvest-years 1877-8 to 1899-1900; the maximum and the minimum annual rainfall, with the corresponding drainage, evaporation, and loss of nitrogen, over the 23 Harvest-years; and the averages for each of the four successive five-yearly periods, to 1896-7 inclusive; also the results for the 21st, 22nd, and 23rd, Harvest-years, 1897-8, 1898-9, and 1899-1900. Then are given, averages for four-monthly periods, and for the total Harvest-year, over the 20 years, 1877-8 to 1896-7. There are also given, in the last four divisions of the Table (p. 19)—first the average loss for each month, over the 20 years; secondly, the monthly results for the Harvest-year, 1898-9; thirdly, those for the last complete Harvest-year, 1899-1900; and lastly, similar results for the current Harve

HARVEST-YEARS. September 1 to August 31.	RAINFALL.		Drainage.			DIFFERENCE (1), evaporated (or retained by soil).			Loss of Nitrogen per Acre in Drainage. Reckoned as Nitrate					
									Reckoned as Nitrogen.			of Soda. (2)		
	5-inch Funnel Gauge.	$\frac{\frac{1}{1000}}{\text{Acre}}$ Gauge.	Soil 20 ins. deep.	Soil 40 ins. deep.	Soll 60 ins, deep.	Soil 20 ins. deep.	Soil 40 ins. deep.	Soil 60 ins. deep.	Soil 20 ins. deep.	Soll 40 ins. deep.	Soil 60 ins, deep.	Soil 20 ins. deep.	Soil 40 ins. deep.	Soil 60 in deep
Av. 19 yrs. 1851–2 to '69–70 Av. 7 yrs. 1870–1 to '76–7	inches. 23:80 28:29	inches, 27·04 30·26	inches. 12·29	inches.	inches. 10:86	inches. 17.97	inches.	inches.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
1877-8 1878-9 1879-80 1880-1 1881-2	32·11 40·17 20·88 35·85 31·66	32.65 41.05 21.36 36.77 32.31	14·72 24·44 6·89 22·38 15·81	16·44 26·03 7·39 22·84 16·08	14.84 24.38 6.50 21.26 14.32	17·93 16·61 14·47 14·39 16·50	16·21 15·02 13·97 13·93 16·23	17.81 16.67 14.86 15.51 17.99	44.75 59.36 27.03 57.78 32.93	39·53 46·52 17·87 44·22 31·74	45·92 60·94 20·19 49·95 35·24	256 379 173 369 211	253 297 114 283 203	293 389 129 319 225
$\begin{array}{c} 1882 - 3 \\ 1883 - 4 \\ 1884 - 5 \\ 1885 - 6 \\ 1886 - 7 \end{array}$	$25 \cdot 29 \\ 25 \cdot 90$	34 · 71 25 · 77 26 · 78 31 · 02 23 · 61	20·82 11·86 14·82 17·37 10·64	21·72 12·00 15·14 18·41 12·58	19·72 11·21 13·98 16·57 11·72	13 · 89 13 · 91 11 · 96 13 · 65 12 · 97	12 · 99 13 · 77 11 · 64 12 · 61 11 · 03	14·99 14·56 12·80 14·45 11·89	32·67 29·31 39·55 34·49 25·28	36·08 26·85 36·71 32·27 21·88	38·26 26·89 33·86 34·36 24·98	209 187 253 221 161	231 172 235 206 140	244 172 216 220 160
1887–8 1888–9 1889–90 1890–1 1891–2	29·11 28·79 26·73 22·30 28·45	30·50 30·09 27·43 23·41 29·68	13·96 14·64 13·16 9·95 16·50	15·82 13·60 9·70	14·67 14·33 12·74 9·73 16·47	16·54 15·45 14·27 13·46 13·18	14·92 14·27 13·83 13·71 12·25	15·83 15·76 14·69 13·68 13·21	43·10 31·96 27·61 25·70 29·39	36·90 29·25 24·94 19·90 28·45	35·67 30·50 28·41 22·04 33·43	276 204 176 164 188	236 187 159 127 181	228 195 182 141 214
$1892-3 \\ 1893-4 \\ 1894-5 \\ 1895-6 \\ 1896-7$	23·11 28·24 27·76 22·98 34·91	24·08 29·55 28·94 24·37 37·24	11.58 13.36 15.50 9.84 21.88	12·35 14·11 16·95 10·75 23·86	12·10 14·07 16·31 10·35 22·80	12·50 16·19 13·44 14·53 15·36	11·73 15·44 11·99 13·62 13·38	11.98 15.48 12.63 14.02 14.44	22 · 61 40 · 94 37 · 12 23 · 18 36 · 62	20·40 31·53 33·18 22·77 35·77	23·72 34·52 34·36 22·78 41·40	144 262 238 148 234	130 202 212 145 229	152 221 220 146 265
1897-8 1898-9 1899-1900						13 · 56 12 · 71 14 · 69		13.04 12.22 14.00	18·20 33·23 37·00	13·95 28·65 33·85	15.01 30.91 37.68	116 213 236	89 183 216	90 197 241
RESULTS FOR MAXIMUM Maximum (1878-9) Minimum (1897-8)	40.17	1	1	26.03 6.66	94.38		15.02	3 HAR 16.67 13.04	59.36	46.52	60.94		297 89	389
AVERAGES FOR 6 for th	5, 5, 5, e 21st y	AND 5	Harv 97-8, 1	EST-Y che 22n	EARS (20 YE. 1898-	ARS, 18	877–8 the 23r	ro 189 d year,	6 - 7). 1899-	Also the 1900.	ne resu	ılts	
5 yrs., 1877-8 to '81-2 5 yrs., 1882-3 to '86-7 5 yrs., 1887-8 to '91-2 5 yrs., 1892-3 to '96-7	32·13 27·39 27·07 27·40	32·83 28·38 28·22 28·83	16.85	17·76 15·97 14·42 15·60	F			71	44·37 32·26 31·55 32·09	1.00	42·45 31·67 30·01	284 206 202 205	230 196 178 184	27: 20: 19: 20:
Mean, 20 years	28.50 18.21 23.25 30.04	29·57 19·51 24·69 31·02	15·00 5·95 11·99 16·33	15 · 94 6 · 66 12 · 48 16 · 93	14·90 6·47 12·48 17·02	14·57 13·56 12·70 14·69	13·63 12·85 12·21 14·09	14·67 13·04 12·21 14·00	$ \begin{array}{r} 35 \cdot 07 \\ 18 \cdot 20 \\ 33 \cdot 23 \\ 37 \cdot 00 \end{array} $		15·01 30·91 37·68	224 116 214 236	197 89 183 216	21 9 19 24
AVERAGES FOR							ST-YEA	RS (20	YEAR:	s, 1877 17 · 30	7-8 TO	1896	-7).	111
Sept. 1 to Dec. 31	11·19 7·37 9·94	11·57 7·76 10·24	$\begin{vmatrix} 7 \cdot 47 \\ 4 \cdot 74 \\ 2 \cdot 79 \end{vmatrix}$	$ \begin{vmatrix} 7 \cdot 72 \\ 5 \cdot 32 \\ 2 \cdot 90 \end{vmatrix} $	7·23 5·01 2·66	$\begin{vmatrix} 4 \cdot 10 \\ 3 \cdot 02 \\ 7 \cdot 45 \end{vmatrix}$	3·85 2·44 7·34	$ \begin{array}{r} 4 \cdot 34 \\ 2 \cdot 75 \\ 7 \cdot 58 \end{array} $	7·49 7·31	7 · 94 5 · 59	9:57 6:35	48 47	51 36	64
Total Harvest-year	28.50	29.57	15.00	15.94	14.90	14.57	13.63	14.67	35.07	30.83	33.87	224	197	21

⁽¹⁾ Calculated on the Rainfall shown by the $\frac{1}{1000}$ th acre gauge.

⁽²⁾ Commercial—reckoning 5 per cent. impurity.

NUMERICAL SUMMARY OF RESULTS	RELATING TO RAINFALL	AND DRAINAGE AT	ROTHAMSTED—continued.
------------------------------	----------------------	-----------------	-----------------------

HARVEST-YEARS. September 1 to August 31.		Rat	NFALL,		Drainage.			Difference (1), evaporated (or retained			Loss of Nitrogen per Aure in Drainage.					
			1					by soil).			Reckoned as Nitrogen.			Reckoned as Nitrate of Soda. (2)		
		5-inch Funnel Gauge.		Soil 20 ins. deep.	Soil 40 ins, deep.	Soil 60 ins. deep.	Soil 20 ins, deep.	Soil 40 ins. deep.	Soil 60 ins. deep.	Soil 20 ins, deep.	Soil 40 ins. deep.	Soil 60 ins. deep.	Soil 20 ins. deep.	Soil 40 Ins. deep.	Soil 60 in deep	
		AVERAG	ES FOR	EACH	Монтн	. 20 H	ARVEST	-Years	s, 1877	-8 то	1896–7					
September		inches.		inches.	inches.	inches.	inches.	inches.	inches.	lbs.	1bs.	lbs.	lbs.	1bs.	lbs.	
) a & a la	***	0.00	2.63	1.04	1:02	0.95	1.59	1 61	1.68	3.91	2.73	2.69	25.0			
T	5000 34	0.05	3.38	2.07	2:09	1.93	1.31	1 29	1.45	6.38	5.00	5.04	40:7			
	WW 54	$\frac{3}{2} \cdot 32$	3.14	2.41	2:51	2:37	0.73	0.63	0.77	6.30	5.66	5.86	40.3			
	148 14	1 01	2.04	1.95	2:10	1.98	0.47	0.32	0:44	3.68	3.91	4.36	23.5	25.0		
la la mara a mara	333 3	1.07	1.95	1.67	1.88	1.82	0.37	0.16	0.22	2.57	2.81	3.40	16.4			
AT 1	1881 18	1 00	1.88		1:71	1.58	0.40	0.24	0.37	2:46	2 51	3.00	15.7	16.0		
11	100 B	1.00	1.89	$\begin{vmatrix} 1.00 \\ 0.52 \end{vmatrix}$	1.15	1.08	0.88	0.73	0.80	1.48	1.67	2.00	9.4	10.6		
π [*]	1000	0.11	2.17	0.58		0.53	1.37	1 31	1.36	0.98	0.95	1.17	6.3	6.1	7	
une		0.00	2 33	0.61	0.65	0.57	1.59 1.72	1 52 1 68	1.60	1.13	1.04	1.19	7.2	6·6 7·0		
uly		0.79	2.80	0.43	0.63	0.69			1.72	1.24	1.09	1.25	7.9			
ugust		0.04	2.94	0.87	0.86	0.69	$2.07 \\ 2.07$	2.06	2 11	2.07	1.53	1.69	13.3	9.8		
Total		00. 50	29.57	15.00	15.94	14.90			2.15	2.87	1.93	2.22	18.4			
10000		. 28.50	120 01	110 00			14.57	Describer-	14.67	35.07	30.83	33 · 87	224 · 1	197.0	210	
eptember	-	. 0.52	0.60	1	HAR	vest-Ye	,		0.00		1					
Van Land	35. 3	0.75	2.89	1.22	1.16	1.06	$\begin{vmatrix} 0.60 \\ 1.67 \end{vmatrix}$	0.60	0.60	5.23	3.65	3.45	33.4	23.3	22	
Tourombon		0.00	2.44	1.87	1.81	1.72	0.57	1.73	0.72	8.67	5.73	5.70	55.4			
ecember		2.82	3.01	2.37	2.44	2.47	0.64	0.57	0.54	9.31	6.14	6.44	59.5			
anuary		2.79	2.96	2.46	$\frac{2}{2} \cdot 71$	2.75	0.50	0.25	0.21	5.13	6.20	6.65	32.8			
ebruary		2 33	2.44	2.08	2.13	2.12	0.36	0.31	0.32	2.26	3.41	4.17	14.5	0.00		
Iarch		0.80	0.87	(0.004)		0.04	0.87	0.83	0.83	0.01	0.06	0.07	(0.04)	0.4		
pril		2.61	2.73	0.67	0.76	0.82	2.06	1.97	1.91	0.85	1.22	1,53	5.4			
Гау		2.64	2.81	1.12	1.18	1.22	1.69	1.63	1.59	1.39	1.79	2.32	8.9			
une		1.48	1.58	0.04	0.04	0.07	1:54	1.54	1.51	0.09	0.07	0.13	0.6	0.4		
July		1.18	1.27	0.16	0.21	0.21	1.11	1.06	1.06	0.29	0.38	0.45	1.9	2.4		
August	***	1.01	1.09			(0.001)		1.09	1.09	-37	ä	(0.002)		944	(0.0	
Total		23.25	24 · 69	11.99	12.48	12.48	12.70	12.21	12:21	33 · 23	28.65	30.91	212-4	182.9	197	
			•	La	ST HA	RVEST-Y	EAR, 18	399-196	00.							
eptember	199 %	. 2.36	2.46	0.26	0.19	0.16	2.20	2 · 27	2.30	1.40	0.79	0.63	8.9		4	
October		3.60	3.75	2.67	2.68	2:56	1.08	1.07	1.19	10.22	7.16	8.46	65.3			
Vovember	34 %	3 69	3.76	3.46	3:45	3 44	0.30	0.31	0.32	10.58	8.82	9.49	67.6			
December		1:36	1.41	0.89	0.86	0.83	0.52	0.55	0.58	2.32	1.92	2.04	14.8			
anuary		3 54	3.67	3.29	3:54	3.46	0.38	0.13	0.21	5.20	6.48	7.20	33.3	41.4		
ebruary		4 82	4.91	4.22	4 39	4 71	0.69	0.52	0.20	3.24	5.06	6.61	20.7	32.3		
Iarch		0.95	10.96	0.13	0 27	0.28	0.83	0.69	0.68	0.15	0.34	0.44	1.0	2.1	2	
pril		1.33	1.33	0.31	0.40	0.41	1.02	0.93	0.92	0.47	0.50	0.43	3.0	3.2	2	
Iay		$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	1.08	(0.001)	0.01	0.02	1.08	1.07	1.06	0.07	0.01	0.04	.:.4	0.1	0	
une		1.07		0.04	0.05	0.08	2.59	2.58	2.55	0.07	0.08	0.14	0.4	0.5		
uly Lugust		3.70	3.93	1:06	1.08	0·03 1·04	1 13 2 87	1·12 2·85	1·10 2·89	3.35	0·04 2·65	0·04 2·16	21:4	0·2 16·9		
Total			31.02	16.33	16.93					37.00						
				Curi	RENT H	IARVEST-	YEAR,	1900-1	.901.							
eptember	(#S#) S#	. 0.73		(0.001)		0.01	0.84	0.83	0.83	1	0.01	0.02	F	0.1	0	
october		$2 \cdot 45$	2.60	0.74	0.57	0.53	1.86	2.03	2.07	3.40	1.56	2.18	21.7			
lovember		. 2.45	2.61	2.16	2.40	2.41	0.45	0.21	0.20	10.04	6.89	7.09	64.1			
December		. 3.52	3.65	2.99	3.21	3.19	0.66	0.44	0.46	9.88	7.85	7.72	63.1	50.2		
anuary	386 36	1.02	1.18	0.67	0.88	0.85	0.21	0.30	0.33	1.56	2.04	2.00	10.0			
ebruary	*** · **	1.19	1.26	0.60	0.90	0.77	0.66	0.36	0.49	1.18	1.87	1.70	7.5	11.9		
Iarch		2.39	2.57	1.50	1.84	1.75	1.07	0.73	0.82	2.54	3.24	3.44	16.2			
pril	300 3	2.42		1.31	1.49	1.47	1.20	1.02	1.04	2.01	2.37	2.72	12.9			
ſау	900 B	1.77	1.81	0.40	0.49	0.47	1.41	1.32	1.34	0.63	0.76	0.87	4:0	4.9	5	
une	(** 9			1 2	T A		1		to C	13						
uly		•		1	127		1 1 '	11 15	100	8 2	UD a	P15				
						1		1 1 1				T I				
lugust	388 G					7	11 14	the P		31						

⁽¹⁾ Calculated on the Rainfall shown by the $\frac{1}{1000}$ th acre gauge.

⁽²⁾ Commercial—reckening 5 per cent. impurity.