Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1901




Full Table of Content

Barley; Hoos Field

Rothamsted Research

Rothamsted Research (1902) *Barley; Hoos Field ;* Yields Of The Field Experiments 1901, pp 24 - 27 - **DOI:** https://doi.org/10.23637/ERADOC-1-229



(25)

RESULTS OF EXPERIMENTS MADE IN HOOS FIELD ON THE GROWTH OF

BARLEY,

for 50 years in succession on the same land—without manure, with Farmyard manure, and with various artificial manures.

The results show, that on the growth of Barley year after year on ordinary arable land, the produce by mineral manures alone is higher than that without manure; that nitrogenous manures alone give more produce than mineral manures alone; and that mixtures of both mineral and nitrogenous manures give much more than either used alone—indeed, generally twice, or more than twice, as much as mineral manures alone. Of mineral constituents, whether used alone or in mixture with nitrogenous manures, phosphates were much more effective than mixtures of salts of potash, soda, and magnesia. The averages show that, under all conditions of manuring (excepting with farmyard manure), the produce was less over the later than over the earlier periods of the experiments—a result partly due to the seasons. But the average produce for 48 years of continuous growth of Barley has, in all cases where nitrogenous and mineral manures (containing phosphates) were used together, been much higher than the average produce of the crop grown in ordinary rotation in the United Kingdom; and very much higher than the average in most other countries when so grown.

Barley is appropriately sown in a lighter soil than Wheat; and whilst Wheat is usually sown in the autumn, Barley is as a rule sown in the spring; and hence it relies in a much greater degree on the stores of the surface soil. Accordingly, it is more susceptible to exhaustion of the surface-soil in nitrogenous, and especially in mineral supplies; and hence, in the common practice of agriculture, it more generally requires the direct application of mineral manures, especially phosphatic manures, than does Wheat when grown under equal soil conditions. The exhaustion induced by both crops is, however, characteristically that of available nitrogen; and when, under the ordinary conditions of manuring and cropping, artificial manure is still required, nitrogenous manures are as a rule requisite for both crops; and for the spring sown Barley more generally than for Wheat, phosphatic manures also. It is not recommended that Barley should in practice be grown year after year on the same land by artificial manures as in these experiments; but, in addition to the lighter soils on which it is more appropriately grown in ordinary rotation, it may be grown, both in full quantity per acre and of good quality, after Wheat, or other grain crop, on the heavier soils, when the land is clean enough for a second cereal crop.

For details of the manuring and produce of the different plots, see pages 26 and 27.

26)

HOOS

EXPERIMENTS ON THE GROWTH OF BARLEY YEAR AFTER YEAR ON THE

Previous Cropping-1847, Swedish Turnips, with Dung and Superphosphate of Lime, the Roots carted off; 1848, Barley (with clover); 1849, Clover; 1850, Wheat; 1851, Barley manured with Amm.-salts. First Experimental Barley Crop in 1852. Barley every year since. The crop of the present year, 1901, is, therefore, the 50th Barley crop in succession. Unless stated to the contrary in the Table, or in the foot-notes, the same Manure has been applied year after year to the same Plot. Description of

(Area under experiment,

PLOTS.	1 acre = (about) 0.404 Hectare or 1.585 Prussian Morgen, 1 bushel = (about) 0.364 Hectolitre or 0.662 Prussian Scheffel, 1 lb. (pound avoir.) = (about) 0.453 Kilogramme or 0.907 Zollverein Pfund, 1 cwt. (hundredweight) = (about) 50.8 Kilogrammes or 1.016 Centner, 1 bushel per acre = (about) 0.9 Hectolitre per Hectare or 0.418 Pr. Scheffel per Pr. Morgen, 1 lb. per acre = (about) 1.12 Kilogramme per Hectare or 0.572 Zollv. Pfd, per Pr. Morgen, 1 cwt. per acre = (about) 125.6 Kilogrammes per Hectare or 0.641 Centner per Pr. Morgen. Manures, per acre, per annum. [In 1898 and since, 400 lbs. Basic Slag used throughout instead of Superphosphate.]
1 O. 2 O. 3 O. 4 O.	Unmanured continuously 3½ cwts. Superphosphate of Lime (1) 200 lbs. (2) Sulphate Potash, 100 lbs. (3) Sulphate Soda, 100 lbs. Sulphate Magnesia 200 lbs. (2) Sulphate Potash, 100 lbs. (3) Sulphate Soda, 100 lbs. Sulphate Magnesia, 3½ cwts. Superphosphate
1 A. 2 A. 3 A. 4 A.	200 lbs. Ammonium-salts (4) 200 lbs. Ammonium-salts, and 3½ cwts. Superphosphate 200 lbs. Ammonium-salts, 200 lbs. (2) Sulph. Potash, 100 lbs. (3) Sulph. Soda, 100 lbs. Sulph. Magnesia (200 lbs. Ammonium-salts, 200 lbs. (2) Sulph. Potash, 100 lbs. (3) Sulph. Soda, 100 lbs. Sulph. Magnesia, 3½ cwts. Superphosphate
$^{(5)} \begin{cases} 1 & AA \\ 2 & AA \\ 3 & AA \\ 4 & AA \end{cases}$	275 lbs. Nitrate Soda 275 lbs. Nitrate Soda, and 3½ cwts. Superphosphate 275 lbs. Nitrate Soda, 200 lbs. (2) Sulph. Potash, 100 lbs. (3) Sulph. Soda, 100 lbs. Sulph. Magnesia. (275 lbs. Nitrate Soda, 200 lbs. (2) Sulph. Potash, 100 lbs. (3) Sulph. Soda, 100 lbs. Sulph. Magnesia, 3½ cwts. Superphosphate
(f) {1 AAS. 2 AAS. 3 AAS. 4 AAS.	275 lbs. Nitrate Soda, 400 lbs. Silicate Soda (6)
$(7) \begin{cases} 1 & \text{C.} \\ 2 & \text{C.} \\ 3 & \text{C.} \\ 4 & \text{C.} \end{cases}$	1000 lbs. Rape-cake 1000 lbs. Rape-cake, and 3½ cwts. Superphosphate 1000 lbs. Rape-cake, 200 lbs. (2) Sulph. Potash, 100 lbs. (3) Sulph. Soda, 100 lbs. Sulph. Magnesia 1000 lbs. Rape-cake, 200 lbs. (2) Sulph. Potash, 100 lbs. (3) Sulph. Soda, 100 lbs. Sulph. Magnesia, 3½ cwts. Superphosphate
1 N. 2 N.	275 lbs. Nitrate Soda
5 O. 5 A. M.	200 lbs. (2) Sulphate Potash, $3\frac{1}{2}$ cwts. Superphosphate (10)
$6igl\{ 1 \ 2$	Unmanured continuously
$7{1 \choose 2}$	Farmyard Manure 14 tons, 20 yrs., 1852-71; unmanured since

the same amount of Nitrogen as 200 lbs. "Ammonium-salts."

^{(1) &}quot;Superphosphate of Lime," 1852 to 1887 inclusive, made from 200 lbs. Bone-ash, 150 lbs. Sulphuric acid, sp. gr. 1.7 (and water); 1888-1897, made from high percentage mineral phosphates, and containing 37 per cent., or more, of soluble phosphate. In 1898, and since, 400 lbs. Basic Slag used throughout instead of Superphosphate.

(2) 300 lbs. per annum for the first six years, 1852-7.

(3) 200 lbs. per annum for the first six years, 1852-7.

(4) The "Ammonium-salts" (excepting in 1887), equal parts Sulphate and Muriate of Ammonia of Commerce. In 1887 Sulphate Ammonia only, 225 lbs. per acre, equal in Nitrogen to the "Ammonium-salts" of previous years. In 1901, the south half of Plots 1 A, 2 A, 3 A, and 4 A, received instead of "Ammonium-salts" as above, Bicarbonate of Ammonia containing an amount of nitrogen equivalent to that of the "Ammonium-salts" as above, Bicarbonate of Ammonia containing an amount of nitrogen equivalent to that of the "Ammonium-salts" as above, Bicarbonate of Ammonia containing an amount of Nitrogen as 200 lbs. Ammonium-salts per annum; next 10 years, 1858-67, 200 lbs. Ammonium-salts per annum; 1868, and since, 275 lbs. Nitrate of Soda per annum. 275 lbs. Nitrate of Soda is reckoned to contain the same amount of Nitrogen as 200 lbs. "Ammonium-salts."

(27)

FIELD.

SAME LAND, WITHOUT MANURE, AND WITH DIFFERENT DESCRIPTIONS OF MANURE.

Barley—29 years, 1852–1880, Chevalier; 10 years, 1881–1890, Archer's Stiff Straw; 7 years, 1891–1897, Carter's Paris Prize; 1898 and since, Archer's Stiff Straw. In the spring of 1894 permanent division paths were laid out between plot and plot. Below is given, besides the usual averages, the produce for both 1899 and 1900.

For Plan of the Plots, and brief summary of results and conclusions, see pp. 24-25.

about 41 acres.)

Plots.		re'i					Produ	CE PER A	CRE.							
	Dressed Grain. Total Straw.															
	Quantity.					Weight per Bushel.					Total Staw.					PLOTS.
	Averages.			48th 4	49th	Averages.				49th	Averages.		3.	48th Year,	49th Year,	×
	24 Yrs. 1852-76.	24 Yrs. 1876–99.	48 Yrs. 1852-99.	Year, 1899.	Year, 1900.	24 Yrs. 1852-75.	24 Yrs. 1876-99.	48 Yrs. 1852-99.		Year, 1900.	24 Yrs. 1852-75,	24 Yrs. 1876-99.	48 Yrs. 1852-99.	1899.	1900.	
1 O. 2 O. 3 O. 4 O.	Bush. 187 244 218 258	Bush. $12\frac{1}{2}$ $16\frac{5}{8}$ $12\frac{1}{8}$ 16	Bush. 15\frac{5}{8} 20\frac{1}{2} 16\frac{5}{8} 21	Bush. 8 $11\frac{1}{2}$ $7\frac{1}{8}$ $11\frac{1}{2}$	Bush. $8\frac{1}{4}$ $10\frac{1}{4}$ $7\frac{1}{4}$ $12\frac{1}{2}$	$\begin{array}{c} \text{lbs.} \\ 52\frac{1}{2} \\ 53\frac{1}{4} \\ 53\frac{1}{6} \\ 53\frac{1}{2} \end{array}$	1bs. $51\frac{3}{4}$ $53\frac{1}{8}$ $52\frac{1}{8}$ $52\frac{7}{8}$	1bs. 52\frac{1}{53\frac{1}{4}} 52\frac{1}{52\frac{1}{4}} 52\frac{1}{4}	1bs. $51\frac{3}{8}$ $54\frac{1}{4}$ $53\frac{1}{8}$ $54\frac{5}{8}$	$ \begin{array}{r} $	Cwts. 11 123 1138 133	Cwts. 67/888666666666666666666666666666666666	Cwts. 9 108 91 108	Cwts. 51 62 55 78 78	Cwts. 5\frac{3}{6}\frac{1}{2}\$ 5 8\frac{1}{8}	1 O. 2 O. 3 O. 4 O.
1 A. 2 A. 3 A.	$31\frac{5}{8}$ $46\frac{3}{8}$ $34\frac{3}{8}$	$22\frac{3}{4}$ $35\frac{1}{4}$ $25\frac{3}{4}$	27¼ 40¼ 30½	$18\frac{1}{2}$ $28\frac{1}{4}$ $23\frac{1}{4}$	$16\frac{1}{8}$ $14\frac{1}{8}$ $20\frac{3}{4}$	$52\frac{1}{4}$ $53\frac{1}{2}$ $52\frac{7}{8}$	$51\frac{7}{8}$ 52 $52\frac{5}{8}$	52½ 52¾ 52¾	52 1 53 54	51 49 § 53	177 265 20	$12\frac{3}{4}$ $19\frac{1}{2}$ $14\frac{7}{8}$	15 <u>1</u> 23 17 <u>1</u>	16 13½	9 93 103 103	1 A. 2 A. 3 A.
4 A.	451	401	423	293	241	541	54	541	54 8	535	28	227	258	175	14	4 A.
1 AA. 2 AA. 3 AA.	36 48 ⁵ / ₈ 36 ³ / ₈	$ \begin{array}{r} 26\frac{1}{8} \\ 40\frac{3}{4} \\ 28 \end{array} $	31½ 44½ 32½	29 43½ 30¼	$ \begin{array}{r} 21\frac{1}{8} \\ 27\frac{1}{8} \\ 25\frac{3}{4} \end{array} $	52 1 53 8 52 1	$52\frac{1}{8}$ $53\frac{1}{2}$ $52\frac{7}{8}$	52½ 53¾ 52¾	53 - 55 8 54 8	51 3 52 3 52 3	$ \begin{array}{r} 21\frac{1}{8} \\ 29\frac{1}{2} \\ 23 \end{array} $	$15\frac{5}{8} \\ 23\frac{3}{4} \\ 17\frac{3}{8}$	18§ 26§ 20§	287	$14\frac{1}{8}$ $17\frac{1}{2}$ $15\frac{1}{2}$	1 AA. 2 AA. 3 AA.
4 AA.	485	393	444	$40\frac{1}{2}$	313	53 <u>5</u>	54	533	56 1	53½	311	243	27 ह	263	193	4 AA.
1 AAS. 2 AAS.	37 <u>8</u> 47 <u>1</u>	33½ 43½	(34§ 44§	35 ³ / ₄	$30\frac{7}{8}$ $26\frac{3}{8}$	54 <u>4</u> 55 8	53½ 54	533 548	54 3 56	528 534	$21\frac{3}{8}$ $28\frac{1}{2}$	$19\frac{1}{2}$ $25\frac{3}{8}$	(201 263		18½ 15½	1 AAS 2 AAS
3 AAS.	42	$35\frac{1}{8}$	(12) 37 8	35½	293	543	54	(12) 541	555	538	$24\frac{7}{8}$	207	(12){22 1	_	16 1	3 AAS
4 AAS.	487	431/2	45§	$40\frac{1}{2}$	36½	554	543	547	56 8	53 8	305	27½	288	26 §	211/8	4 AAS
1 C. 2 C. 3 C.	$44\frac{1}{2}$ $46\frac{1}{4}$ $42\frac{3}{4}$	35 8 38 8 34 8	397 428 388	$ \begin{array}{r} 30\frac{3}{4} \\ 33\frac{1}{2} \\ 27\frac{7}{8} \end{array} $	$25\frac{1}{4}$ $23\frac{1}{4}$ 20	53½ 53½ 53½	54 54 8 54 8	533 544 54	548 548 548	52½ 52½ 52½	25½ 27§ 26	197 213 194	223 248 223	18	$12\frac{7}{8}$ $13\frac{1}{8}$ $11\frac{7}{8}$	1 C. 2 C. 3 C.
4 C.	463	37	417	31	$22\frac{1}{2}$	533	541	54	545	523	288	215	25	177	12	4 C.
1 N. 2 N.	37 41	29 7 34 1	$(^{13})$ ${338 \atop 37\frac{1}{2}}$	33 37½	27 § 28 §	52 <u>3</u> 52 3	52 7 53 1	$\binom{13}{53\frac{1}{8}}$	548 54½	51 7 507 508	$22\frac{1}{2}$ $25\frac{1}{4}$	$17\frac{1}{2}$ $20\frac{1}{2}$	$\binom{13}{228}$	$\frac{19}{22\frac{7}{8}}$	14 8 16	1 N. 2 N.
5 O. 5 A. M.	$21\frac{3}{8}$ $43\frac{5}{8}$ $19\frac{7}{8}$	$ \begin{array}{c c} 14\frac{1}{8} \\ 32 \\ 18\frac{1}{2} \end{array} $	$\binom{13}{378}$ $\binom{14}{378}$ $\binom{14}{194}$	$9\frac{3}{4}$ $27\frac{1}{4}$ $\binom{15}{1}$	$7\frac{1}{8}$ $20\frac{1}{8}$ $(^{15})$	53½ 54 53¾	53½ 53½ 53¼	$\binom{13}{53}$ $\binom{53}{53}$ $\binom{14}{53}$	52 55§ (15)	50¾ 53¼ (15)	$11\frac{3}{8}$ $27\frac{1}{4}$ $11\frac{5}{8}$	83 20½ 97 8	$\binom{13}{23\frac{3}{4}}$ $\binom{14}{10\frac{3}{4}}$	$6\frac{1}{2}$ $17\frac{1}{2}$ $(^{15})$	$5rac{58}{8}$ $15rac{1}{8}$ $(^{15})$	5 O. 5 A. M.
$6{1 \choose 2}$	$\frac{20\frac{5}{8}}{21}$	$\begin{array}{ c c c }\hline 13\frac{3}{4} \\ 14\frac{7}{8} \\ \end{array}$	171 172	$\frac{6\frac{1}{2}}{7\frac{5}{8}}$	$10\frac{1}{4}$ $11\frac{1}{8}$	52 5 52 3 52 3	52½ 52¾	528 528	52 52	51½ 52½	$11\frac{5}{8}$ $11\frac{5}{8}$	$\frac{7\frac{3}{4}}{7\frac{7}{8}}$	9§ 9§	5 8 5 1	$7\frac{1}{2}$ $6\frac{5}{8}$	${1 \choose 2} 6$
$7{1 \choose 2}$	48g 48g 48g	$27\frac{7}{8}$ $48\frac{3}{8}$	(16) 368 485 485	$\frac{12\frac{3}{8}}{42}$	157 317	54 8 54 8	54 54§	(16) 54½ 54¾	55 <u>1</u> 57	52 5 54 1	$28\frac{1}{4}$ $28\frac{5}{8}$	$15\frac{3}{4}$ $30\frac{5}{8}$	(16) 21 29§	8 28	10 1 18 1	${1 \choose 2} 7$

⁽⁶⁾ The application of Silicates did not commence until 1864; in 1864-5-6 and 7, 200 lbs. Silicate of Soda and 200 lbs. Silicate of Lime were applied per acre; but in 1868, and since, 400 lbs. Silicate of Soda, and no Silicate of Lime. These plots ("AAS") comprise, respectively, one half of the original "AA" plots, and, excepting the addition of the Silicates, have been, and are, in other respects, manured in the same way as the "AA" plots.

(7) 2000 lbs. Rape-cake per annum for the first six years, and 1000 lbs. only, each year since.

(8) 300 lbs. Sulphate of Potash, and 3½ cwts. Superphosphate of Lime, without Nitrate of Soda, the first year (1852); Nitrate alone each year since.

(9) 550 lbs. Nitrate of Soda for 1853-4-5-6, and 7; and 275 lbs. only, each year since.

(10) Ammonium-salts also the first year, but not since.

(11) By mistake 400 lbs. in 1880.

(12) Averages of 12, 24, and 36 years, 1864-99.

(13) Averages of 23, 24, and 47 years, 1853-99.

(14) Averages of 17, 17, and 34 years, 1858-78, and 1880-92. The produce of 1879 was not weighed, owing to the foulness of the plot, from the wet season.

(15) Not recorded.

(16) Averages of 20, 28, and 48 years, 1852-99.