

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1948

[Full Table of Content](#)



Yields of the Field Experiments 1948 - Results

Rothamsted Research

Rothamsted Research (1949) *Yields of the Field Experiments 1948 - Results* ; Yields Of The Field Experiments 1948, pp 1 - 98

14474

Rothamsted Experimental Station
Harpenden
LAWES AGRICULTURAL TRUST

RESULTS
OF THE
FIELD
EXPERIMENTS
1948



LIBRARY

STATS SECTION



Rothamsted Experimental Station
Harpenden

Lawes Agricultural Trust

RESULTS
of the
FIELD
EXPERIMENTS
1948

The summaries given in this report are similar to those contained in the appendices to the Annual Reports of the Station before the war. With one or two special exceptions only experiments conducted at Rothamsted and Woburn are included. The design and supervision of these experiments are the responsibility of the Field Plots Committee (present members: E.M.Crowther (Chairman) H.V.Garner (Secretary), H.H.Mann, J.R.Moffatt, D.J.R.Watson, F.Yates). The results of series of experiments conducted on commercial farms (such as the factory series of sugar beet fertilizer experiments) will be published elsewhere.

Reports covering the war years are being prepared.

Price: 5/-

Index

Classicals

Broadbalk	Wheat	A/1
Hoosfield	Barley	A/2
Hoosfield	Wheat after fallow	A/3
Barnfield	Mangolds	A/4
Park Grass	Hay	A/5
Agdell	Rotation	see A/3

Long Term

2-course	Rothamsted	Ba/1
3-course	Rothamsted	Ba/2
4-course	Rothamsted	Ba/3
6-course	Rothamsted and Woburn	Ba/4
6-course	Deep cultivation	Bb/1
Green Manuring	Woburn	Bc/1
Ley and Arable	Woburn	Bd/1
Market Garden	1st crops	Be/1
Market Garden	2nd crops	Be/2
Highfield Grazing		Bf/1

Short Term

Wheat	Eyespot	Ca/1
Wheat	Residual organics	Ca/2
Wheat	Wireworm 1	Ca/3
Wheat	Wireworm 2	Ca/4
Wheat	Inoculation	Ca/5
Wheat	Varieties	Ca/6
Spring Sown Cereals		Cb/1
Beans	Cultivations and varieties	Cc/1
Potatoes	Organics	Cd/1
Potatoes	Cultivation	Cd/2
Potatoes	Time of planting	Cd/3
Sugar Beet	Fertilizer placement	Ce/1
Linseed	Rothamsted: Time and date of sowing	Cf/1
Linseed	Rothamsted: Rates and application of manures	Cf/2
Linseed	Woburn: Time and date of sowing	Cf/3
Linseed	Woburn: Rates and application of manures	Cf/4
Peas		Cg/1

Outside Experiments

Hay	Lady Manners School: Artificial and compost	Da/1
Hay	Lady Manners School: Effects of fertilizers	Da/2

Miscellaneous Data

Chemical Analyses of Manures	Z/1
Meteorological Readings	Z/2

48/A/1.1

WHEAT - BROADBALK 1948

The 105th year

Treatments as listed in 1938 Report p.p. 115-6, except that since the year 1940-41 rape cake has been replaced by castor bean meal, at the same rate of application.

Cultivations, etc.:

Cropped sections. Dung applied: Sept. 24. Ploughed: Sept 22-
Sept. 24. Artificial's applied: Oct. 13-15. Rolled and
springtine harrowed: Oct. 22. Tooth harrowed: Oct. 23.
Seed drilled: Oct. 24-25. Harrowed in: Oct. 25. Harrowed:
Apr. 12. Ring rolled: Apr. 20. Nitrogeous fertilizers
applied: Apr. 27-28. Second dressing of nitrate of soda
applied to plot 16: May 7. Wild oats hand pulled: various
days June 15 - July 30. Harvested: Aug. 19-20. Variety:
Squareheads Master (13/4).
Fallow section. Ploughed: Sept. 22-24. Rolled and springtine
harrowed: Oct. 22. Tooth harrowed: Oct. 23 and 25.
Springtine harrowed: Apr. 12. Thistles cut: Apr. 20.
Ploughed: May 25-27. Ring rolled: June 9. Thistles cut:
July 1. Ploughed: July 14-15. Rolled and springtine
harrowed: Aug. 6.

Section Years after Fallow	Total Grain: cwt. per acre					Total straw: % cwt. per acre				
	II	I	III	IV	Mean	II	I	III	IV	Mean
	1	2	3	4		1	2	3	4	
Plot 2A	32.4	23.5	20.0	23.1	24.8	62.0	47.5	29.5	50.0	47.2
2B	36.0	30.4	23.8	25.3	28.9	68.9	58.5	47.8	51.8	56.8
3	17.3	17.2	10.9	9.8	13.8	31.3	27.5	19.5	19.1	24.5
5	20.9	14.9	9.8	10.4	14.0	38.2	28.5	21.8	22.4	27.7
6	21.6	20.5	14.3	14.5	17.7	42.0	37.3	26.8	31.9	34.5
7	26.1	27.9	20.2	20.4	23.6	47.8	48.3	41.7	44.0	45.4
8	28.4	31.8	25.6	27.5	28.3	55.2	67.5	52.2	52.6	56.9
9	22.2	22.3	15.4	16.6	19.1	40.1	40.8	31.2	29.6	35.4
10	14.2	24.2	17.6	12.8	17.2	32.2	40.8	28.1	24.9	31.5
11	16.3	21.8	17.8	14.7	17.6	28.8	38.1	30.7	26.8	31.1
12	20.2	24.4	19.7	19.7	21.0	33.2	38.6	32.6	34.0	34.6
13	26.2	26.4	18.7	17.7	22.2	43.8	46.2	33.5	35.1	39.6
14	19.4	28.3	20.2	20.6	22.1	34.7	44.7	33.8	37.7	37.7
15	21.6	18.4	16.8	15.0	18.0	41.6	41.5	31.1	33.1	36.8
16	28.1	26.6	22.6	23.1	25.1	49.5	39.9	46.6	46.3	45.6
17	20.5	22.5	17.6	16.6	19.3	42.6	39.2	33.0	34.6	37.4
18	17.3	12.9	8.8	8.2	11.8	30.9	20.7	17.8	17.5	21.7
19	19.6	26.5	18.3	17.8	20.6	38.8	43.6	34.1	36.8	38.3
20	14.0	25.1	-	-	19.6	29.4	45.1	-	-	37.2

48/A/1.2

* Incl des straw, cavings and chaff

48/A/2

BARLEY - HOOSFIELD 1948

Treatments as listed in 1938 Report, p. 117, except that since 1940-41 rape cake has been replaced by castor bean meal at the same rate of application.

Cultivations, etc.: Ploughed (shallow): Sept. 18-20. Springtime harrowed: Sept 30. Dung applied: Nov 24. Ploughed: Nov 22-28. Springtime harrowed: Mar 1-5. Artificials applied: Mar 10-11. Thistles cleared: Mar 27. Harrowed: Mar 30. Seed drilled: Mar. 31 Harrowed in: Apr 6. Rolled: Apr 10. Wild oats hand pulled: July 5-15 and July 30 - Aug 10, and scythed: Aug 9. Harvested: Aug 30.

Note: On plots 50, 5A and 3C, sections badly infected with wild oats were scythed out and discarded. On all other plots the full area was harvested.

Plot	Total Grain cwt./acre	Total Straw [*] cwt./acre
1 O	7.6	10.9
2 O	10.3	11.9
3 O	8.9	11.5
4 O	14.2	18.3
5 O	16.3	21.2
1 A	10.6	16.5
2 A	16.0	21.1
3 A	12.2	20.2
4 A	21.6	27.5
5 A	19.3	23.2
1 AA	13.3	25.9
2 AA	20.5	27.1
3 AA	15.3	23.2
4 AA	22.4	27.1
1 AAS	17.5	27.8
2 AAS	21.8	31.5
3 AAS	22.4	30.0
4 AAS	24.6	30.1
1 C	15.2	21.6
2 C	17.7	23.2
3 C	17.3	26.4
4 C	23.3	18.9
7 - 1	12.6	15.6
7 - 2	28.2	40.2
6 - 1	6.6	10.1
6 - 2	8.7	11.4
1 N	12.8	22.8
2 N	16.9	27.4

* Includes straw, cavings and chaff.

48/A/3

WHEAT AFTER FALLOW - HOOSFIELD 1948

Without manure 1851 and since.

For details of treatments see 1938 Report p.109.

Cultivations, etc.:

Cropped sections: Ploughed: Sept.15-16. Rolled, springtime harrowed and harrowed: Oct.24. Seed drilled and harrowed in: Oct.25. Ring rolled: Mar.31. Weeds hand pulled: July 29. Harvested: Aug.23.

Fallow sections: Ploughed: Sept.15-16. Rolled, springtime harrowed and harrowed: Sept.24. Springtime harrowed: Mar.10. Thistles cut: Apr.22. Ploughed: May 21. Ring rolled: June 9.

No. of years of Fallow Section	Produce: cwt. per acre			Mean
	1 A2	1 A3	3 A4	
Total Grain	9.3	11.1	10.6	10.3
Total Straw	16.6	19.6	17.5	17.9

Note "CROPS IN ROTATION - AGDELL FIELD 1948"

Manures were applied as usual, but turnips were sown on plots 1 and 2 only. The crop was badly infected with "finger and toe", and was discarded.

MANGOLDS - BARNFIELD 1948.

48/A/4.1

Treatments as listed in 1938 Report, p.110, except that on cross dressings AC and C, rape cake has been replaced by castor bean meal at the same rate of application.

Note. Both the Mangold and Sugar Beet crops were almost destroyed by flea beetle, and only Mangolds were resown.

Cultivations, etc.: Dung applied: Nov.25. Ploughed various days: Nov. 26-Dec. 16. Cultivated: Mar. 25. Artificials applied: Apr.20 - 23. Thistles cut: Apr.24. Harrowed: Apr. 26. Rolled: Apr. 27. Seed drilled: Apr. 28. Dusted with flea beetle dust: May 14, 19 and 22. Plant almost destroyed by flea beetle by May 29, so decision made to resow mangolds only. Thistles cut: May 29. Springtine harrowed: June 8. Harrowed, rolled and seed redrilled June 9. Top dressings applied: June 10. Harrowed in seed and artificials: June 11. Dusted: June 19-22. Dung plot hoed: July 8. Hoed: July 12-15. Singled: July 12-21. Hoed: July 26-27, Aug. 16-25, Sept. 9-11, Sept 24-27 and Oct. 11. Lifted: Nov. 3-18. Variety: Yellow Globe.

Note: For leaf weights, two random rows were chosen from each plot. The weight of each of these samples was recorded separately.

		Cross Dressings				48/A/4.2
Strip	O	N	A	AC	C	
Roots: tons per acre						
1	9.52	18.19	15.49	18.35	12.61	
2	10.84	18.90	18.31	22.52	17.12	
4	2.19	(a) 12.15	14.45	19.80	10.01	
		(b) 12.88				
5	1.54	9.27	6.34	8.38	3.90	
6	1.85	10.78	13.32	16.84	6.91	
7	1.47	8.28	11.12	12.83	5.57	
8	1.35	6.53	5.58	7.44	4.78	
9	11.73					
Leaves: tons per acre						
1	3.34	4.82	4.16	5.75	6.02	
2	4.94	6.29	5.82	6.63	6.43	
4	1.96	(a) 5.43	5.65	7.68	5.99	
		(b) 6.14				
5	1.10	4.21	2.94	3.84	2.76	
6	1.20	4.31	5.02	5.31	4.33	
7	0.91	4.62	5.46	7.68	4.65	
8	1.13	3.72	3.06	3.38	3.16	
9	5.19					

HAY - THE PARK GRASS PLOTS, 1948.

48/A/5

For details of treatments and notes, see 1935 Report p.151 and 1938 Report p.111.

Cultivations etc.: Chain harrowed: Dec.10. Limed: Jan.3-7
 Minerals applied: Jan.17-20. Ring rolled: Mar. 24.
 Nitrogenous manures applied: 1st dressing Apr. 5; 2nd dressing May 4. 1st cut: June 16-18. 2nd cut: Oct.7-9

Yield of Hay: cwt. per acre.

Plot	Not limed			Limed		
	1st Crop	2nd Crop***	Total	1st Crop	2nd Crop***	Total
1	9.1	10.0	19.1	18.0	5.4	23.4
2	15.8	9.2	25.0	16.6	6.6	23.2
3	13.9	9.0	22.9	14.6	7.0	21.6
4-1	19.1	7.0	26.1	18.3	11.9	30.2
4-2	20.2	6.2	26.4	31.0	6.9	37.9
5-1	9.4	4.2	13.6	-	-	-
5-2	16.0	12.1	28.1	-	-	-
6	30.0	14.4	44.4	-	-	-
7	31.5	17.9	49.4	32.4	17.4	49.8
8	25.5	10.6	36.1	12.2	6.9	19.1
9	34.6	11.2	45.8	41.0	7.6	48.6
10	18.7	11.6	30.3	32.5	7.0	39.5
11-1	46.2	23.9	70.1	51.5	10.0	61.5
11-2	42.3	10.4	52.7	63.3	9.1	72.4
12	13.7	9.5	23.2	-	-	-
13	28.4	12.4	40.8	31.2	7.6	38.8
14	55.1	15.0	70.1	52.5*	11.2*	63.7*
				47.2**	4.8**	52.0**
15	21.6	22.2	43.8	19.4	7.8	27.2
16	39.7	15.8	55.5	42.9	8.1	51.0
17	22.4	8.8	31.2	24.1	8.5	32.6
18	3.5	8.1	11.6	35.1+	5.8+	40.9+
	-	-	-	29.2++	8.0++	37.2++
19	26.0	10.5	36.5	26.6+	6.2+	32.8+
	-	-	-	32.7++	0.4++	33.1++
20	39.7	12.0	51.7	41.6+	11.8+	53.4+
	-	-	-	40.4++	8.9++	49.3++

* Sun

** Shade

*** The figures for the second crop are estimated hay yields calculated from the dry matter.

+ Heavy liming

++ Light liming

48/Ba/1.1

TWO COURSE ROTATION EXPERIMENT

Cumulative Effects of Agricultural Salt

Rothamsted 1948

(For full details see 1942 Report)

Treatments:

All combinations of:

- (1) Agricultural salt: None, $2\frac{1}{2}$, 5 and $7\frac{1}{2}$ cwt per acre applied to sugar beet or at half these rates to barley.
- (2) Muriate of potash: None, the equivalent of half the single dressing of salt (approximately 1 cwt. K_2O per acre), the equivalent of the single dressings of salt (approximately 2 cwt. K_2O per acre), applied to sugar beet at sowing.
- (3) Time of application of salt: In seed bed at sowing, before ploughing in winter.
- (4) Salt applied to sugar beet only, salt repeated at half rate on barley.

Note: treatment (3) applies to both sugar beet and barley crops; the barley receives no potash treatment. For ease of analysis K_2O of the muriate of potash dressings is made equivalent to Cl of agricultural salt dressings.

Basal dressings, applied to all plots at sowing:

Barley: 0.3 cwt. N per acre as sulphate of ammonia
Sugar beet: 0.8 cwt. N per acre as sulphate of ammonia
0.6 cwt. P_2O_5 per acre as superphosphate.

Cultivations, etc:

Sugar Beet. Series I. Long Hoos VII

Agricultural salt applied: Oct. 21. Ploughed: Nov. 10-21. Cultivated: Mar. 16. Rolled and springtime harrowed, ring rolled: Mar. 27. Sulphate of ammonia applied: Mar. 30. Superphosphate, agricultural salt and muriate of potash applied: Apr. 12. Harrowed: Apr. 14. Seed drilled: Apr. 15. Seed harrowed in: Apr. 16. Rolled: Apr. 22. Dusted with Flea beetle dust: May 14. Plots 6, 12, 18, 24, 30, 36, 42, 48 damaged by herbicide spray from adjacent barley; May 14. Bare patches resown: late May. Hoed: May 20-21, 25. Weeds hand pulled: June 9-11. Singled: June 10-11. Hoed: June 21 - July 22. Lifted: Nov. 23-30. Variety: Klein E. Previous crop: Barley.

48/Ba/1.2

Barley. Series II. Long Hoos V.
Agricultural salt applied: Dec 2. Ploughed: Dec 10-12.
Springtime harrowed: Mar 1. Harrowed, seed drilled,
sulphate of ammonia applied: Mar 8. Applied agricultural
salt, harrowed in, ring rolled: Mar 9. Sprayed with
"Denocate" to clear charlock, (plots 66, 65, 64, 63, 62,
61 damaged by an overdose of spray): May 14. Hand weeded:
June 30. Harvested: Aug. 14. Variety: Plumage Archer.
Previous crop: Sugar Beet.

Standard errors per plot:

Sugar beet, total sugar,	4.40 cwt. per acre or	10.9%
tops,	1.13 tons per acre or	7.8%
Barley, grain,	2.20 cwt. per acre or	7.1%

All standard errors estimated from 22 d.f.

48/Ba/1.3

Series I: Sugar Beet

Salt in 1948 (cwt. per acre)	Muriate of potash K ₂ O cwt. per acre			Salt applied		Mean
	0.0	1.0	2.0	Winter	In seed bed	
Total Sugar: cwt. per acre						
		(±2.20)		(±1.68)		(±1.27)
0	33.4	34.2	35.7	41.6	39.6	34.4
2.5	43.7	37.9	40.2	41.6	39.6	40.6
5.0	41.8	44.6	42.0	42.2	43.5	42.8
7.5	41.9	44.6	46.1	43.4	45.1	44.2
Mean	40.2	40.3 (±1.10)	41.0	42.4	42.7 (±1.02)	40.5
Sugar percentage						
0	16.24	16.34	16.57			16.38
2.5	16.49	16.46	16.78	16.57	16.58	16.57
5.0	16.75	16.90	16.42	16.58	16.80	16.69
7.5	16.59	16.62	16.84	16.78	16.60	16.69
Mean	16.51	16.58	16.65	16.64	16.66	16.58
Roots (washed): tons per acre						
0	10.26	10.44	10.74			10.48
2.5	13.27	11.50	11.97	12.56	11.93	12.25
5.0	12.49	13.20	12.74	12.69	12.93	12.81
7.5	12.63	13.42	13.71	12.92	13.59	13.25
Mean	12.16	12.14	12.29	12.73	12.81	12.20
Tops: tons per acre						
		(±0.566)		(±0.433)		(±0.327)
0	14.21	14.80	13.90			14.30
2.5	14.57	14.27	14.68	13.05	15.97	14.51
5.0	13.28	13.75	15.36	14.86	13.40	14.13
7.5	14.90	17.08	13.56	13.79	16.57	15.18
Mean	14.24	14.98 (±0.283)	14.37	13.90	15.31 (±0.261)	14.53

48/Bn/1.4

Series I: Sugar Beet (contd.)

Salt in 1948 (cwt. per acre)	Muriate of potash K ₂ O cwt. per acre			Salt applied		Mean
	0.0	1.0	2.0	Winter	In seed bed	
	Plant No.: thousands per acre					
0	22.3	22.2	21.4			22.0
2.5	22.0	21.9	22.3	21.9	22.3	22.1
5.0	21.9	22.6	21.6	21.7	22.3	22.0
7.5	22.1	22.4	21.5	22.2	21.8	22.0
Mean	22.1	22.3	21.7	21.9	22.1	22.0
	Noxious Nitrogen: m.g. per 100 grams.					
0	30.2	31.5	31.2			31.0
2.5	28.5	30.2	28.8	28.7	29.7	29.2
5.0	29.8	28.6	28.0	29.3	28.3	28.8
7.5	31.0	29.5	30.8	31.2	29.7	30.4
Mean	29.9	30.0	29.7	29.7	29.2	29.8

48/Ba/1.5

Series II: Barley

Salt in 1947 (cwt. per acre)	Muriate of potash K ₂ O cwt. per acre applied in 1947			Salt applied In seed bed		Salt in 1948 None Half Rate		Mean
	0.0	1.0	2.0	Winter				
	Grain: cwt. per acre			cwt. per acre				
	(±1.10)			(±0.84)				(±0.63)
0	31.3	33.6	30.6					31.8
2.5	32.6	30.8	31.2	31.6	31.5	31.2	31.9	31.5
5.0	30.2	31.1	29.8	30.8	30.0	29.1	31.6	30.4
7.5	29.6	30.0	29.6	30.5	29.0	29.1	30.3	29.7
Mean	30.9	31.4	30.3	31.0	30.2	29.8	31.3	30.9
	(±0.55)			(±0.51)				
	Straw: cwt. per acre			cwt. per acre				
0	35.9	37.0	35.3					36.1
2.5	36.8	33.2	34.7	34.8	34.9	35.6	34.1	34.9
5.0	32.7	33.4	32.0	31.9	33.5	32.1	33.3	32.7
7.5	32.4	32.2	31.5	31.4	32.7	30.7	33.4	32.0
Mean	34.4	34.0	33.4	32.7	33.7	32.8	33.6	33.9

48/Ba/2.1

THREE COURSE ROTATION EXPERIMENT

Long Hoos VI, 1948

Effect of ploughing in straw

Treatments as given in 1933 Report, pp. 118-9, except that no comparisons of winter green manuring crops are now made, and that commencing in 1942 a yearly dressing of $2\frac{1}{2}$ cwt. per acre magnesium sulphate is applied to one of the replicate plots of each treatment, in each crop block.

Cultivations, etc.

Barley, Series I.

Applied Adco and straw with accompanying artificials, ploughed: Nov 10-21. Springtine harrowed: Mar 1. Artificials applied: Mar 3. Harrowed, seed drilled and harrowed in: Mar 4. Ring rolled: Mar 9. Weeded: June 8. Harvested: Aug 18. Variety: Plumage Archer. Previous crop: Potatoes.

Potatoes, Series II.

Applied Adco and straw with accompanying artificials, ploughed: Nov 10-21. Cultivated: Mar 16. Harrowed and rolled: Mar 22. Ridged: Mar 24. Artificials applied, potatoes planted and covered in: Mar 25. Ridges rolled down: Mar 27. Harrowed: Apr 30. Weeded: June 11. Grubbed: June 9, 16. Earthed up: June 17. Sprayed with "Perinox": Aug 9. Sprayed to kill haulm: Sept 13. Lifted: Sept 29. Variety: Majestic. Previous crop: Sugar Beet.

Sugar Beet, Series III.

Applied Adco and straw with accompanying artificials, ploughed: Nov 10-21. Cultivated: Mar 16. Harrowed and rolled: Mar 22. Artificials applied: Apr 13. Harrowed: Apr 14. Seed drilled: Apr 15. Harrowed in: Apr 16. Rolled: Apr 22. Dusted with Flea beetle dust: May 14. Plots 54, 60, 66 and 72 were damaged by herbicide spray from 2 Course Rotation Experiment, May 14, and bare patches (including the whole of plots 66 and 72) were redrilled May 20. Hoed: May 20. Late sown plants dusted with Flea beetle dust: June 9. Weeded: June 10-11. Singled: June 12. Hoed: June 21, July 20. Lifted: Nov 12, 15. Variety: Klein E. Previous crop: Barley.

48/Ba/2.2

Note: Considerable damage to sugar beet tops was caused by sheep on plots 49, 50, 51, 55, 56, 57, 61, 62, 67, 68.

Standard errors per plot:

Sugar beet,	roots (washed),	± 0.573 tons per acre or 5.9%
	tops,	± 1.348 tons per acre or 18.8%
	total sugar,	± 2.69 cwt. per acre or 7.7%
	sugar percent:	± 0.437
Barley,	plant number,	± 1.66 thousands per acre or 7.6%
	grain,	± 3.47 cwt. per acre or 10.4%
Potatoes	straw,	± 4.75 cwt. per acre or 12.6%
	total tubers,	± 0.661 tons per acre or 6.4%
	percentage ware	± 0.581

All standard errors are based on 8 d.f.

48/Ba/2.3

Summary of Results

	Treatments applied 1946/7					Treatments applied 1947/8				
	Arts	Adco	St 1	St 2	Mean	Arts	Adco	St 1	St 2	Mean
<u>Series I</u>										
Barley										
Grain cwt./acre	34.0	32.1 (±2.00)	33.3	33.3	33.1 (±1.00)	36.8	32.3 (±2.00)	31.0	35.2	33.9 (±1.00)
Straw cwt./acre	36.8	35.9 (±2.74)	36.8	36.5	36.5 (±1.37)	43.1	36.5 (±2.74)	36.9	40.0	39.1 (±1.37)
<u>Series II</u>										
Potatoes tons/acre	8.47	10.12 (±0.382)	9.78	10.41	9.70 (±0.191)	10.40	10.40 (±0.382)	12.35	11.12	11.07 (±0.191)
Percentage Ware	92.0	93.5 (±0.335)	93.6	93.6	93.2 (±0.168)	94.0	93.7 (±0.335)	95.3	94.4	94.4 (±0.168)
<u>Series III</u>										
Sugar Beet Roots (washed) tons/acre	9.30	9.12 (±0.331)	8.95	9.63	9.25 (±0.166)	10.79	9.04 (±0.331)	10.21	10.69	10.18 (±0.166)
Tops tons/acre	6.41	6.24 (±0.778)	7.09	7.43	6.79 (±0.389)	9.49	6.14 (±0.778)	7.28	7.40	7.58 (±0.389)
Sugar percentage	17.32	17.89 (±0.252)	17.13	17.59	17.48 (±0.126)	18.18	18.18 (±0.252)	18.12	17.88	18.09 (±0.126)
Total sugar cwt./acre	32.2	32.6 (±1.55)	30.6	33.9	32.3 (±0.78)	39.3	32.9 (±1.55)	37.0	38.3	36.9 (±0.78)
Plant number thous./acre	22.2	22.6 (±0.96)	20.0	22.6	21.9 (±0.48)	21.8	22.0 (±0.96)	22.3	22.2	22.1 (±0.48)

48/Ba/2.4

Responses to Magnesium Sulphate

	Treatments applied 1946/7					Treatments applied 1947/8				
	Arts	Adco	St.1	St.2	Mean	Arts	Adco	St.1	St.2	Mean
<u>Series I</u>										
Barley Grain cwt./acre	-4.3	-0.6 (±4.25)	2.8	-5.0	-1.8	0.2	-8.0 (±4.25)	-5.9	1.9	-3.0
Straw cwt./acre	-4.0	-4.2 (±5.82)	6.1	-3.4	-1.4	-0.2	-5.4 (±5.82)	-4.9	0.4	-2.5
<u>Series II</u>										
Potatoes tons/acre	-0.38	0.22 (±0.810)	0.07	-0.07	-0.04	1.12	0.44 (±0.810)	0.82	-1.14	0.31
Percentage Ware	-0.4	0.3 (±0.71)	-0.6	-0.6	-0.3	0.2	0.3 (±0.71)	0.6	0.0	0.3
<u>Series III</u>										
Sugar Beet Roots (washed) tons/acre	-0.34	-1.02 (±0.702)	-1.80	-0.50	-0.92	1.06	1.28 (±0.702)	-1.21	0.01	+0.28
Tops tons/acre	-1.77	-0.92 (±1.652)	0.56	2.06	-0.02	1.38	1.21 (±1.652)	-2.44	0.52	0.17
Total sugar cwt/acre	0.7	-2.7 (±3.30)	-5.6	-2.7	-2.6	7.3	4.8 (±3.30)	-5.0	1.1	2.0
Sugar percentage	1.02	0.56 (±0.535)	0.36	-0.46	0.37	1.54	0.07 (±0.535)	-0.30	0.48	0.45
Plant number thous./acre	0.6	-2.2 (±2.03)	-7.1	-3.8	-3.1	0.6	2.6 (±2.03)	-4.6	-1.2	-0.6

48/Ba/3.1

FOUR COURSE ROTATION EXPERIMENT

Hoosfield, 1948

Residual values of organic and phosphatic fertilizers.

The scheme of the experiment is as given in the 1932 Report, pp.127-8, with the following alterations:-

1. From 1935 onwards, clover ryegrass ley has been replaced by ryegrass alone, sown in autumn after ploughing barley stubble, with fertilizers applied as on wheat.
2. From 1935 onwards, lime was applied every year at the rate of 10 cwt. per acre to the potato break after the crop had been lifted.
3. Each plot of the potato break has been split from 1942 onwards, a random half of each plot receiving an additional 2 cwt. per acre sulphate of ammonia.
4. Majestic potato seed has been used since 1942 in place of Ally.

Manures applied 1947-8

RESULTS OF FIELD EXPERIMENTS - 1948

ERRATUM

48/Ba/3.1, Table of manures applied 1947-8 should read:-

Manures applied 1947-8

Treatment	Organic fertilizers (cwt. per acre)				Additional artificial fertilizers (cwt. per acre)		
	Organic matter	N	P ₂ O ₅	K ₂ O	N as Sulph. of amm.	P ₂ O ₅ as Super	K ₂ O as mur. of potash
Dung	50 (as FYM)	1.404	0.709	2.230	0.396	0.491	0.770
Adco	50 (as Adco)	0.980	0.510	0.419	0.820	0.690	2.581
Straw	77 (as Straw)	1.392	0.104	0.627	0.408	1.096	2.373
Super			None		0.36	1.2	0.6
Rock phosphate			None		0.36	1.2*	0.6

* As mineral phosphate

48/Ba/3.2

Cultivations, etc.

Potatoes, Series I.

Ploughed: Sept. 8. Dung, Adco with accompanying artificials applied; straw with first dressing of artificials applied; ploughed in: Nov. 20, 21. Second dressing of artificials applied to straw plot: Jan. 22. Ploughed, harrowed, rolled: Mar. 16-23. Ridged: Mar. 30. Artificials, including third dressing to straw plot applied in the ridges; Apr. 2. Potatoes planted and covered in: Apr. 7. Ridges rolled: Apr. 10. Ridges harrowed: May 10. Grubbed: June 9-16. Weeded: June 11-13. Earthed up: June 26. Hoed: June 30. Weeded: July 29. Sprayed with "Perinox": Aug. 9. Sprayed to kill haulm: Sept. 13. Lifted: Sept. 27, 28. Variety: Majestic. Previous crop: Wheat.

Barley, Series II

Applied dung, Adco and supplementary artificials: Nov. 12. Straw with first dressing of artificials applied; ploughed: Nov. 12, 13. Ground chalk (22 cwt. per acre) applied: Nov. 13. Second dressing of artificials applied to straw plots: Jan. 22. Springtine harrowed: Mar. 1. Artificials applied, including third dressing to straw plots: Mar. 4. Harrowed, seed drilled and harrowed in: Mar. 5. Ring rolled: Mar. 10. Hand weeded to clear thistles: May 28. Hoed: June 4. Harvested: Aug. 19. Variety: Plumage Archer. Previous crop: Potatoes. Note: Ground chalk was used in place of ground lime.

Wheat, Series III

Applied dung and Adco with supplementary artificials, applied straw with first dressing of artificials, ploughed: Sept. 15-24. Flat rolled and harrowed: Sept. 25. Artificials applied excluding special dressing straw plots and sulphate of ammonia: Sept. 27. Harrowed: Oct. 2. Springtine harrowed, harrowed, seed (Squareheads Master 13/4) drilled and harrowed in: Oct. 3. Resown with Squareheads Master (13/4) as plant had failed, second dressing of artificials applied to straw

48/Ba/3.3

plot, harrowed in: Nov.29. Plant failed over winter, decided to resow, springtime harrowed, harrowed: Mar.8. Redrilled with Atlo, applied sulphate of ammonia and third dressing of artificials to straw plots, harrowed in: Mar.9. Ring rolled: Mar.10. Weeded to clear thistles: May 28. Hoed: June 4. Harvested: Aug.28. Variety: Atle. Previous crop: Ryegrass.

Ryegrass, Series IV.

Dung, Adco with accompanying artificials applied: Sept.24. Straw with first dressing of artificials applied, ploughed: Sept.23-25. Harrowed and rolled: Sept.26. Autumn artificials applied not including sulphate of ammonia: Sept.27. Harrowed twice, ring rolled: Nov.10. Seed sown and harrowed in: Nov.13. Second dressing of artificials applied to straw plots: Nov.29. Applied sulphate of ammonia, third dressing of artificials applied to straw plot: Mar.17. Rolled: Mar.20. Hand weeded to clear thistles: June 12. Cut: June 15. Variety: Western Worths. Previous crop: Barley.

FOUR COURSE ROTATION
Summary of Results 1948

Manure	Year of Cycle	Wheat		Barley		Ryegrass cwt per acre Dry matter	Total tubers Additional N		tons per acre Mean to N		Percentage Ware Additional N		Resp. to N
		Grain	Straw	Grain	Straw		Without	With	Without	With	Mean	Resp. to N	
Manure as F.Y.M.	I	16.3	37.6	32.1	46.3	6.5	5.94	8.70	7.32	2.76	97.1	97.6	0.5
	II	9.7	29.5	27.1	28.1	5.7	7.89	10.09	8.99	2.20	97.2	97.7	0.5
	III	9.9	28.0	22.9	27.4	5.6	7.75	9.09	8.42	1.33	97.3	97.7	0.4
	IV	8.6	25.7	24.0	31.7	4.9	8.05	9.61	8.63	1.56	97.9	98.0	0.1
	V	10.4	29.2	21.4	29.0	4.4	7.62	9.42	8.52	1.80	97.2	98.4	1.2
Manure as A.D.C.	I	20.2	47.8	35.7	26.6	16.7	9.09	10.72	9.90	1.63	98.1	98.4	0.3
	II	10.7	32.5	24.4	30.4	3.7	6.37	9.69	8.03	3.32	97.9	98.8	0.9
	III	8.9	23.0	25.1	30.4	3.5	6.04	9.71	7.88	3.67	97.4	98.2	0.8
	IV	9.0	21.2	23.6	30.6	4.4	7.94	9.46	8.70	1.52	98.1	98.2	0.1
	V	9.7	24.6	22.6	31.9	2.8	5.85	9.28	7.56	3.43	97.0	98.3	1.3
Manure as Straw	I	21.6	47.8	36.5	47.1	19.2	10.19	12.30	11.24	2.11	97.6	98.1	0.5
	II	11.0	33.0	28.3	33.6	6.6	6.54	8.55	7.54	2.01	96.1	97.3	1.2
	III	13.1	32.0	23.9	25.0	6.5	6.79	10.56	8.68	3.77	97.2	98.2	1.0
	IV	10.6	29.0	26.4	32.0	4.8	7.39	9.67	8.53	2.28	97.2	98.4	1.2
	V	9.7	25.8	24.7	28.7	2.9	8.29	10.65	9.47	2.36	97.5	97.6	0.1
Super phos- phate	I	15.1	32.4	32.1	34.2	9.3	10.69	11.65	11.17	0.96	97.5	97.2	-0.3
	II	14.1	32.0	27.6	32.5	5.5	9.20	9.81	9.50	0.61	98.1	98.8	0.7
	III	12.7	30.4	30.2	33.0	9.4	7.69	11.38	9.64	3.49	98.5	98.8	0.3
	IV	11.6	23.8	27.5	32.1	6.1	7.59	8.39	7.99	0.80	97.7	98.0	0.3
	V	11.7	28.4	29.0	31.8	2.9	7.32	8.90	8.11	1.58	97.6	98.7	1.1
Rock phos- phate	I	16.6	33.5	26.8	33.8	4.6	9.25	9.95	9.60	0.70	97.3	98.1	0.8
	II	12.1	27.1	26.2	31.2	4.0	7.03	7.82	7.42	0.79	97.3	97.8	0.5
	III	11.9	27.8	28.4	31.9	4.0	6.13	7.80	6.96	1.67	96.6	98.0	1.4
	IV	13.5	26.2	27.5	35.2	2.4	6.31	5.82	6.06	-0.49	97.2	95.9	-1.3
	V	10.3	21.8	29.0	45.6	4.6	6.74	6.22	6.48	-0.52	97.7	97.5	-0.2

48/Ba/3.4

48/Ba/4.1

SIX COURSE ROTATION EXPERIMENT

1947-8

Seasonal effects of N, P₂O₅ and K₂O

Rotation and treatments as given in 1932 Report, p. 131, with the exceptions that since 1934 the forage crop has been replaced by rye, harvested as a mature crop, and that the green manure crops are now omitted. Since 1934 lime has been applied at the rate of 10 cwt. per acre at two stages in the rotation: immediately after the removal of the potato crop and before sowing barley.

Cultivations, etc.:

Rothamsted, Long Hoos IV

Sugar beet. Series 3.

Ploughed: Sept 11, 12. Harrowed: Mar 22. Ploughed, harrowed and ring rolled: Apr 15, 16. Artificials applied, seed drilled and harrowed in: Apr 16. Ring rolled: Apr 22. Applied Agroxide ($\frac{1}{2}$ cwt. per acre) against wireworm: Apr 29. Dusted with Flea beetle dust: May 8, 14, 15. Applied Agroxide ($1\frac{1}{2}$ cwt. per acre): May 10. Hoed: June 8 - July 22. Singled: June 9. Lifted: Nov 10. Variety: Klein E. Previous crop: Rye.

Barley. Series 4.

Sugar beet tops remaining from 1947 crop ploughed in: Nov 7. Ground chalk applied (22 cwt. per acre); Nov 14. Springtine harrowed: Mar 1. Artificials applied: Mar 4. Harrowed, seed drilled and harrowed in: Mar 5. Ring rolled: Mar 9. Clover seed undersown: Apr 15. Harrowed and ring rolled: Apr 16. Harvested: Aug 18. Variety: Plumage Archer. Previous crop: Sugar beet.

Clover. Series 1.

Seed undersown in barley, harrowed in: Apr 12. Ring rolled: Apr 14. Phosphate and potash fertilizer applied: Dec 10. Rolled: Mar 10. Sulphate of ammonia applied: Mar. 16. Self sown barley cut: Apr 26. Cut: July 6. Variety: Late flowering Montgomery Red. Previous crop: Barley.

Wheat. Series 5.

Ploughed: June 30. Twice harrowed: Oct 1. Autumn artificials applied, seed drilled and harrowed in: Oct 2. Seed resown (owing to plant failure) and harrowed in: Nov 12. Rolled: Mar 16. Sulphate of ammonia applied: Apr 15. Harvested: Aug 18. Variety: Yeoman. Previous crop: Clover.

48/Ba/4.2

Potatoes. Series 6.

Ploughed: Sept. 9, 10. Harrowed: Mar. 10. Ploughed and harrowed: Mar. 10, 11. Harrowed and rolled: Mar. 22. Ridged: Mar. 23. Artificials applied, potatoes planted and covered in: Mar. 24. Ridges rolled down: Mar. 27. Harrowed: Mar. 30. Hoed: June 4-10. Grubbed: June 7, 18. Earthed up: June 19. Sprayed with "Perinox": Aug. 9. Sprayed to kill haulm: Sept. 13. Lifted: Sept. 29. Variety: Majestic. Previous crop: Wheat.

Rye. Series 2.

Ploughed: Sept. 30 - Oct. 1. Harrowed twice, ring rolled, harrowed: Oct. 1. Autumn artificials applied, seed drilled and harrowed in: Oct. 3. Ground chalk (22 cwt. per acre) applied: Nov. 14. Ring rolled: Mar. 31. Sulphate of ammonia applied: Apr. 16. Cut: July 30. Previous crop: Potatoes.

Woburn, Stackyard Field, Series B.

Sugar beet. Series 4.

Ploughed: Sept. 16, Oct. 1, Feb. 25-28. Springtime harrowed: Mar. 17. Harrowed: Mar. 19. Rolled: Mar. 27. Seed drilled: Apr. 2. Artificials applied and harrowed in: Apr. 5. Rolled: Apr. 6. Hoed: May 19. Singled, and weeded to clear thistles, hoed: May 25 - June 1. Hoed: June 5 - July 21. Lifted: Oct. 4. Variety: Klein E. Previous crop: Rye.

Barley. Series 6.

Sugar beet tops from 1947 crop ploughed in: Oct. 29-30. Ground chalk applied (96.5% CaCO₃, at rate of 1½ tons per acre of CaCO₃). Springtime harrowed: Mar. 1. Artificials applied, harrowed: Mar. 4. Seed drilled and harrowed in: Mar. 10. Rolled, Clover seed undersown: Mar. 12. Harrowed: Mar. 13. Harvested: Aug. 14. Variety: Plumage Archer. Previous crop: Sugar beet.

Clover. Series 3.

Clover seed (late flowering Montgomery Red) undersown in barley, Apr. 17, 1947. Failed owing to drought. Ploughed: Sept. 26, 27, Feb. 26-28. Harrowed, all artificials including sulphate of ammonia applied: Mar. 4. Rolled, seed sown by hand and harrowed in: May 12. Harrowed and rolled: Mar. 13. Cut: July 1. Variety: Early red Trifolium. Previous crop: Barley.

48/Ba/4.3

Wheat. Series 1.

Ploughed: July 9. Cultivated: Aug. 21, Sept. 1. Rolled three times, cultivated three times, harrowed: Oct. 9. Autumn artificials applied: Oct. 14. Seed drilled and harrowed in: Oct. 14-15. Sowing destroyed by birds. Harrowed, seed drilled and harrowed in: Nov. 14. Harrowed: Apr. 14. Sulphate of ammonia applied: Apr. 22. Rolled: Apr. 26. Harvested: Aug. 28. Variety: Squareheads Master (13/4). Previous crop: Clover.

Potatoes. Series 2.

Ploughed: Sept. 30, Oct. 6, Feb. 27-28. Springtine harrowed: Mar. 17. Harrowed: Mar. 19. Rolled: Mar. 27. Ridged: Apr. 6-7. Artificials applied, potatoes planted and covered in: Apr. 9. Ridges harrowed down: May 8. Grubbed: May 28. June 8. Hoed, earthed up: June 21. Sprayed with "Perinox": Early Aug. Sprayed to kill haulm: Sept. 22. Lifted: Sept. 30. Variety: Majestic. Previous crop: Wheat.

Rye. Series 5.

Ploughed: Oct. 6, 7. Harrowed: Oct. 9. Ground chalk spread (14 cwt. per acre): Oct. 13. Autumn artificials applied: Nov. 14. Seed drilled and harrowed in: Nov. 15. Harrowed: Apr. 14. Sulphate of ammonia applied: Apr. 22. Cut: July 30. Previous crop: Potatoes.

48/Ba/4.4

Mean yields per acre and increments in yield per cwt. of N, P₂O₅ and K₂O

	Rothamsted		Woburn		Rothamsted		Woburn		
	Response	S.E.	Response	S.E.	Response	S.E.	Response	S.E.	
Sugar Beet: Roots (washed) tons/acre					Clover Hay: Dry Matter cwt/acre				
Yield	11.51		8.52		36.3		32.6		
N	0.84	1.263	<u>9.11</u>	1.237	-1.2	5.15	-1.3	6.54	
P	-1.23	1.263	<u>-0.85</u>	1.237	-8.1	5.15	-1.9	6.54	
K	0.19	0.758	-1.35	0.742	6.3	3.09	1.1	3.92	
Tops: tons/acre					Wheat grain: cwt/acre				
Yield	10.72		6.92		26.6		20.8		
N	3.45	2.191	<u>13.31</u>	1.510	7.5	2.15	<u>23.1</u>	5.79	
P	-3.17	2.191	<u>-2.02</u>	1.510	-0.3	2.15	<u>-13.2</u>	5.79	
K	0.70	1.315	<u>-2.28</u>	0.906	3.3	1.29	<u>-11.8</u>	3.47	
Sugar Percentage					Wheat straw: cwt/acre				
Yield	17.87		17.57		47.8		40.2		
N	0.09	0.696	0.76	0.727	18.5		36.9		
P	-0.43	0.696	0.47	0.727	-1.5		-13.3		
K	-0.39	0.418	0.56	0.436	7.4		-19.9		
Total Sugar: cwt/acre					Potatoes: total tubers tons/acre				
Yield	41.2		30.0		9.52		8.94		
N	3.1	4.38	<u>32.9</u>	4.02	6.09	1.541	7.62	1.326	
P	-5.5	4.38	<u>-2.1</u>	4.02	-1.08	1.541	<u>0.55</u>	1.326	
K	-0.3	2.63	<u>-3.8</u>	2.41	<u>2.41</u>	0.924	-0.57	0.795	
Plant Number: thous/acre					Potatoes: Percentage Ware				
Yield	20.9				97.1		93.2		
N	-1.3	2.77			1.7	1.36	-0.7	2.30	
P	2.9	2.77			-2.5	1.36	-4.8	2.30	
K	0.8	1.66			1.8	0.82	<u>-3.5</u>	1.38	

Note: The underlined values are significant at 5% level.

48/Ba/4.5

	Rothamsted Response S.E.	Woburn Response S.E.	Rothamsted Response S.E.	Woburn Response S.E.
Barley Grain: cwt/acre			Rye Grain: cwt/acre	
Yield	<u>36.2</u>	<u>27.5</u>	<u>25.8</u>	<u>21.2</u>
N	<u>10.7</u> 3.71	<u>17.6</u> 5.33	<u>18.4</u> 3.50	<u>15.0</u> 4.39
P	<u>-2.3</u> 3.71	<u>3.5</u> 5.33	<u>2.6</u> 3.50	<u>-5.3</u> 4.39
K	<u>-1.2</u> 2.23	<u>1.7</u> 3.20	<u>2.3</u> 2.10	<u>7.8</u> 2.64
Barley Straw: cwt/acre			Rye Straw: cwt/acre	
Yield	<u>39.6</u>	<u>26.4</u>	<u>47.3</u>	<u>42.8</u>
N	<u>24.7</u>	<u>20.9</u>	<u>14.7</u>	<u>4.5</u>
P	<u>-1.6</u>	<u>-6.5</u>	<u>0.7</u>	<u>-3.8</u>
K	<u>-3.7</u>	<u>3.7</u>	<u>4.6</u>	<u>10.6</u>

Note: The underlined values are significant at 5% level.

48/Bb/1

DEEP CULTIVATION ROTATION EXPERIMENT

Long Hoos I and III, 1948

Objects:

- (1) To compare deep ploughing (about 12 inches) with shallow (about 6 inches)
- (2) To compare the effects of ploughing in the mineral fertilizers at different depths with applying them in the seedbed.
- (3) To compare the effects of ploughing in dung at different depths.

Rotation: Sugar beet, barley, clover-ryegrass ley, wheat, potatoes, spring oats. (Ley ploughed in (not deeper than 6 inches) soon after first cut. Sugar beet tops carted off).

Treatments:

Whole plots. All combinations of:

1. Shallow ploughing (6 inches) v. deep ploughing (12 inches). (This is done before potatoes and sugar beet, and after ley before wheat, at the same time in autumn for all three).
2. Farmyard manure: None v. 20 tons per acre for potatoes and 10 tons per acre for sugar beet, applied before ploughing.
3. Superphosphate: None v. 0.8 cwt. P_2O_5 per acre for potatoes and 0.6 cwt. P_2O_5 per acre for sugar beet.
4. Muriate of potash: None v. 1.0 cwt. K_2O per acre for potatoes and 0.6 cwt. K_2O per acre for sugar beet.

Treatments 2, 3 and 4 are applied only to potatoes and sugar beet.

Half plots (potatoes and sugar beet):

Phosphate and potash (where given) ploughed in v. applied in the seedbed.

Basal manuring: Applied in the ridges for potatoes, as a top dressing for wheat, and in the seed bed for other crops.

Sulphate of ammonia, for potatoes 0.6 cwt. N per acre, for sugar beet 0.8 cwt. N per acre, for spring oats 0.2 cwt. N per acre, for wheat 0.5 cwt. N per acre, for barley 0.3 cwt. N per acre.

Basic slag, for barley 0.6 cwt. P_2O_5 per acre.

Area of each whole plot: 0.03125 acre (before rejecting edge rows).

Cultivations, etc.:

Ley. Series 1-KS.

Seeds undersown in barley: Apr 12, 1947. Rolled: Mar 10, 1948. Cut: June 14, 15. Seeds mixture per acre: 18 lb. ryegrass (Aberystwyth S2.4), 8 lb. late flowering red clover (Cert. N.Z. Mother seed), 2 lb. Alsike clover. Previous crop: Barley.

48/Bb/1.2

Potatoes. Series 2 - KP.

Artificially applied: Sept 29. Dung applied (before deep ploughing), ploughed deep (about 12 inches): Sept 30. Dung applied (before shallow ploughing), ploughed shallow (about 6 inches): Oct 1,2. Cultivated: Mar 11, 20. Rolled and springtime harrowed: Mar 22. Harrowed twice, ring rolled: Mar 23. Ridged: Mar 31. Spring artificials applied, potatoes planted and covered in: Apr 7. Rolled down ridges: Apr 12. Ridges harrowed: May 10. Grubbed: June 9, 18. Earthed up: June 19, 21. Hand weeded: June 10-11, July 28. Sprayed with "Perinox": Aug 9. Sprayed to kill haulm: Sept 13. Lifted: Sept 30. Variety: Majestic (Scotch seed A). Previous crop: Wheat.

Spring oats. Series 3 - KO.

Ploughed: Oct 15-20. Springtime harrowed: Mar 1. Sulphate of ammonia applied; seed drilled and harrowed in: Mar 5. Ring rolled: Mar 9. Weeded: May 28 - June 7. Harvested: Aug 16. Variety: Star. Previous crop: Potatoes.

Sugar beet. Series 4 - KS.

Artificially applied (before ploughing): Sept 29. Dung applied (before deep ploughing), ploughed deep (about 12 inches): Sept 30. Dung applied (before shallow ploughing); ploughed shallow (about 6 inches): Oct 1,2. Cultivated: Mar 11, 20. Rolled and springtime harrowed: Mar 22. Harrowed twice, ring rolled: Mar 23. Spring artificials applied: Apr 13. Harrowed: Apr 14. Seed drilled: Apr 15. Harrowed in: Apr 16. Ring rolled: Apr 22. Dusted with Flea beetle dust: May 8, 14, 15. Thistles hand weeded: June 5. Hoed: June 7 - July 21. Singled: June 8. Hand weeded for thistles: July 28. Lifted: Dec 2-4. Variety: Klein E. Previous crop: Oats.

Barley. Series 5 - KB.

Ploughed: Nov 22-25. Springtime harrowed: Mar 1. Artificially applied, seed drilled and harrowed in: Mar 5. Ring rolled: Mar 9. Seeds mixture undersown: Apr 15. Harrowed and ring rolled: Apr 16. Weeded: May 28 - June 7. Harvested: Aug 16. Variety: Plumage Archer. Previous crop: Sugar beet.

Wheat. Series 6 - KW.

Ploughed: June 17-18, 1947. Ring rolled: June 19. Ploughed deep (about 12 inches): Sept 1. Ploughed shallow (about 6 inches) Sept 2. Cultivated (deep ploughed plots twice, shallow ploughed plots once), ring rolled deep ploughed plots, all plots cultivated: Oct 2. Rolled twice and springtime harrowed: Nov 1. Rolled and springtime harrowed twice: Nov 3. Seed drilled and harrowed in: Nov 4. Rolled: Mar 16. Sulphate of ammonia applied: May 5. Weeded: May 28-June 7. Harvested: Aug 16. Variety: Yeoman. Previous crop: Seeds

Standard errors per plot:

Ley. 1.58 cwt. per acre or 2.4% (4 d.f.)

Potatoes. Ware yield, whole plot, 2.058 tons per acre or 13.9% (4 d.f.)
sub-plot, 2.506 tons per acre or 16.9% (7 d.f.)

Spring oats. Grain, 4.46 cwt. per acre or 14.5% (4 d.f.)

48/Bb/1.3

Sugar beet. Total sugar, whole plot, 3.19 cwt. per acre or 6.0% (4 d.f.)
sub-plot, 5.87 cwt. per acre or 11.1% (7 d.f.)
Tops, whole plot, 0.651 tons per acre or 3.5% (4 d.f.)
sub-plot, 1.849 tons per acre or 9.9% (7 d.f.)
Barley. Grain, 0.846 cwt. per acre or 2.1% (4 d.f.)
Wheat. Grain, 2.69 cwt. per acre or 7.1% (4 d.f.)

48/Bb/1.4

Series I, Ley
Residual Effects

	Responses to treatments								
	Mean	Ploughing		Dung		Super		Potash	
		Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Hay: Mean yield 66.5 cwt. per acre (± 1.12 , Means ± 0.79)									
Ploughing deep									
-shallow	-1.1	-	-	-1.6	-0.6	-1.7	-0.5	-1.1	-1.1
Dung	3.3	2.8	3.8	-	-	0.9	5.7	2.2	4.4
Superphosphate	1.0	0.4	1.6	-1.4	3.4	-	-	4.2	-2.2
Potash	5.8	5.8	5.8	4.7	6.9	9.0	2.6	-	-

Series 2, Potatoes

	Responses to treatments								
	Mean	Ploughing		Dung		Super		Potash	
		Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Ware tubers: Mean yield 14.85 tons per acre (± 1.46 , Means ± 1.03)									
Ploughing deep									
-shallow	0.23	-	-	0.63	-0.17	0.98	-0.52	-1.62	2.08
Dung	4.64	5.04	4.24	-	-	4.23	5.05	6.29	2.99
Superphosphate	1.57	2.32	0.82	1.16	1.98	-	-	0.56	2.58
Potash	2.37	0.52	4.22	4.02	0.72	1.36	3.38	-	-

Percentage ware: Mean 98.7

Ploughing deep									
-shallow	0.3	-	-	0.2	0.4	0.4	0.2	0.2	0.4
Dung	0.4	0.3	0.5	-	-	0.3	0.5	0.6	0.2
Superphosphate	0.0	0.1	-0.1	-0.1	0.1	-	-	0.1	-0.1
Potash	0.5	0.4	0.6	0.7	0.3	0.6	0.4	-	-

48/Bb/1.5

Series 2, Potatoes

	Superphosphate			Potash			Mean
	None	Ploughed in	In ridges	None	Ploughed in	In ridges	
Ware tubers: tons per acre							
	(a)	(b) and (c)		(a)	(b) and (c)		
Shallow	13.57	14.77	17.02	14.47	14.58	15.41	14.74
Deep	14.55	14.51	16.24	12.85	15.82	18.34	14.97
No Dung	11.95	11.40	14.82	10.52	13.39	15.70	12.53
Dung	16.18	17.88	18.44	16.81	17.02	18.05	17.17
Mean	14.06	14.64	16.63	13.66	15.20	16.87	14.84
Percentage ware							
Shallow	98.5	98.4	98.8	98.3	98.6	99.0	98.5
Deep	98.8	98.6	98.8	98.5	99.0	99.2	98.8
No Dung	98.5	98.2	98.7	98.1	98.6	99.0	98.5
Dung	98.8	98.8	98.9	98.7	99.0	99.1	98.9
Mean	98.7	98.5	98.8	98.4	98.8	99.1	98.7

Standard errors (a) ± 1.03 (b) ± 1.25 (c) ± 1.36

Standard error (b) is for use in horizontal comparisons only; standard errors (a) and (c) for use in all other comparisons.

Series 3, Spring Oats

Residual Effects

	Mean	Responses to treatments							
		Ploughing		Dung		Super		Potash	
		Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.

Grain: Mean yield 30.8 cwt. per acre (± 3.16 , Means ± 2.23)

Ploughing deep - shallow	0.9	-	-	2.4	-0.6	-1.8	3.6	-0.1	1.9
Dung	-0.2	1.3	-1.7	-	-	-0.2	-0.2	-2.1	1.7
Superphosphate	-1.0	-3.7	1.7	-1.0	-1.0	-	-	0.8	-2.8
Potash	-1.3	-2.3	-0.3	-3.2	0.6	-0.5	-3.1	-	-

Straw: Mean yield 49.8 cwt. per acre

Ploughing deep - shallow	2.5	-	-	4.1	0.9	0.4	4.6	1.8	3.2
Dung	1.8	3.4	0.2	-	-	3.0	0.6	1.0	2.6
Superphosphate	-3.0	-5.1	-0.9	-1.8	-4.2	-	-	-1.4	-4.6
Potash	-1.9	-2.6	-1.2	-2.7	-1.1	-0.3	-3.5	-	-

48/Bb/1.6

Series 4: Sugar Beet

	Mean	Responses to treatments							
		Ploughing		Dung		Super		Potash	
		Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Total Sugar: Mean yield 52.9 cwt. per acre (± 2.25 , Means ± 1.59)									
Ploughing deep									
-shallow	-4.5	-	-	-0.8	-8.2	-4.9	-4.1	-1.3	-7.7
Dung	2.6	6.3	-1.1	-	-	3.7	1.5	5.5	-0.3
Superphosphate	-0.4	-0.8	0.0	0.7	-1.5	-	-	-0.3	-0.5
Potash	0.6	3.8	-2.6	3.5	-2.3	0.7	0.5	-	-

Roots (washed): Mean yield 15.75 tons per acre									
	Mean	Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Ploughing deep									
-shallow	-1.13	-	-	-0.15	-2.11	-1.22	-1.04	-0.35	-1.91
Dung	0.91	1.89	-0.07	-	-	1.09	0.73	1.48	0.34
Superphosphate	-0.06	-0.15	0.03	0.12	-0.24	-	-	-0.24	0.12
Potash	0.25	1.03	-0.53	0.32	-0.32	0.07	0.43	-	-

Sugar Percentage: Mean 16.79

	Mean	Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Ploughing deep									
-shallow	-0.19	-	-	-0.06	-0.32	-0.22	-0.16	0.01	-0.39
Dung	-0.15	-0.02	-0.28	-	-	0.00	-0.30	0.14	-0.44
Superphosphate	-0.02	-0.05	0.01	0.13	-0.17	-	-	0.19	-0.23
Potash	-0.08	0.12	-0.28	0.21	-0.37	0.13	-0.29	-	-

Tops: Mean yield 13.62 tons per acre (± 0.461 , Means ± 0.326)

	Mean	Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Ploughing deep									
-shallow	2.60	-	-	2.54	2.66	1.52	3.68	2.13	3.07
Dung	1.65	1.59	1.71	-	-	0.94	2.36	0.64	2.66
Superphosphate	-0.01	-1.09	1.07	-0.72	0.70	-	-	-0.96	0.94
Potash	1.44	0.97	1.91	0.43	2.45	0.49	2.39	-	-

Plant Number: Mean 18.3 thousands per acre

	Mean	Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Ploughing deep									
-shallow	-0.7	-	-	-0.4	-1.0	-0.5	-0.9	-0.3	-1.1
Dung	0.4	-0.7	0.1	-	-	0.6	0.2	0.4	0.4
Superphosphate	-0.3	-0.1	-0.5	-0.1	-0.5	-	-	-0.1	-0.5
Potash	-0.1	0.3	-0.5	-0.1	-0.1	0.1	-0.3	-	-

Noxious Nitrogen: Mean 43.9

	Mean	Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Ploughing deep									
-shallow	4.9	-	-	5.8	4.0	4.1	5.7	4.9	4.9
Dung	1.8	2.7	0.9	-	-	2.4	1.2	-0.1	3.7
Superphosphate	-0.9	-1.7	-0.1	-0.3	-1.5	-	-	0.6	-2.4
Potash	2.4	2.4	2.4	0.5	4.3	3.9	0.9	-	-

48/Bb/1.7

Series 4. Sugar Beet

	Superphosphate			Potash			Mean
	None	Ploughed in	In seed bed	None	Ploughed in	In seed bed	
Total Sugar: cwt per acre							
	(a)	(b) and (c)		(a)	(b) and (c)		
Shallow	55.5	57.0	52.5	53.2	60.1	54.0	55.1
Deep	50.6	53.0	48.4	52.0	54.3	44.3	50.7
No dung	51.2	53.4	50.6	49.9	57.8	48.9	51.6
Dung	54.9	56.6	50.3	55.3	56.6	49.4	54.2
Mean	53.1	55.0	50.5	52.6	57.2	49.2	52.9

Roots (washed): tons per acre							
Shallow	16.40	16.83	15.66	15.80	17.75	15.93	16.32
Deep	15.17	16.02	14.39	15.45	16.55	13.30	15.19
No dung	15.23	15.69	15.04	14.89	17.01	14.41	15.30
Dung	16.33	17.17	15.01	16.37	17.29	14.82	16.21
Mean	15.78	16.43	15.02	15.63	17.15	14.61	15.75

Sugar Percentage							
Shallow	16.91	16.93	16.78	16.83	16.94	16.95	16.88
Deep	16.69	16.59	16.82	16.84	16.44	16.68	16.70
No dung	16.80	17.04	16.84	16.76	16.98	16.96	16.87
Dung	16.80	16.49	16.78	16.90	16.49	16.67	16.72
Mean	16.80	16.76	16.80	16.83	16.69	16.81	16.79

Standard errors (b) are for use in horizontal comparisons only, (a) and (c) for use in all other comparisons

Standard errors	(a)	(b)	(c)
Total sugar	±1.59	±2.93	±2.62

48/Bb/1.8

Series 4. Sugar Beet

	Superphosphate			Potash			Mean
	None	Ploughed in	In seed bed	None	Ploughed in	In seed bed	
Tops: tons per acre							
	(a)	(b)	(c)	(a)	(b)	(c)	
Shallow	17.86	17.14	16.40	16.83	18.21	17.40	17.32
Deep	19.39	21.51	19.39	18.97	21.84	19.91	19.92
No dung	18.15	17.65	17.21	17.58	18.62	17.39	17.79
Dung	19.10	21.00	18.58	16.22	21.43	19.92	19.44
Mean	18.62	19.33	17.90	17.90	20.02	18.65	18.62

Plant Number: thous. per acre							
Shallow	18.7	18.3	18.9	18.5	18.9	18.7	18.7
Deep	18.2	17.4	18.0	16.2	17.7	17.7	17.9
No dung	18.1	18.0	18.1	18.1	18.1	18.0	18.1
Dung	18.7	17.7	18.8	18.5	18.5	18.4	18.5
Mean	18.4	17.9	18.4	18.3	18.3	18.2	18.3

Noxious Nitrogen							
Shallow	42.2	40.2	41.0	40.2	44.0	41.2	41.4
Deep	46.4	46.2	46.2	45.1	48.0	47.0	46.3
No dung	43.1	41.8	44.0	42.8	43.5	43.0	43.0
Dung	45.5	44.6	43.2	42.6	48.5	45.2	44.8
Mean	44.3	43.2	43.6	42.7	46.0	44.1	43.9

Standard errors (b) are for use in horizontal comparisons only, (a) and (c) for use in all other comparisons.

Standard errors:	(a)	(b)	(c)
Tops	±0.326	±0.925	±0.730

48/Bb/1.9

Series 5. Barley

Residual effects

Responses to treatments.

	Mean	Ploughing		Dung		Super		Potash	
		Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Grain: Mean yield 40.9 cwt. per acre (± 0.60 , Means ± 0.42)									
Ploughing deep									
-shallow	0.2	-	-	1.4	-1.0	0.0	0.4	0.4	0.0
Dung	1.6	2.8	0.4	-	-	3.1	0.1	2.4	0.8
Superphosphate	1.3	1.1	1.5	2.8	-0.2	-	-	2.4	0.2
Potash	-1.0	-0.6	-1.2	-0.2	-1.8	0.1	-2.1	-	-

Straw: Mean yield 44.8 cwt. per acre

Ploughing deep									
-shallow	0.1	-	-	0.0	0.2	1.0	-0.8	-0.5	0.7
Dung	2.3	2.2	2.4	-	-	3.6	1.0	4.9	-0.3
Superphosphate	0.2	1.1	-0.7	1.5	-1.1	-	-	-1.0	1.4
Potash	0.2	-0.4	0.8	2.3	-2.4	-1.0	1.4	-	-

Series 6. Wheat

Residual effects

Responses to treatments

	Mean	Ploughing		Dung		Super		Potash	
		Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Grain: Mean yield 40.7 cwt. per acre (± 2.04 , Means ± 1.45)									
Ploughing deep									
-shallow	-3.5	-	-	-4.5	-2.5	-2.5	-4.5	-2.3	-4.7
Dung	2.2	1.2	3.2	-	-	2.3	2.1	4.5	-0.1
Superphosphate	0.7	1.7	-0.3	0.8	0.6	-	-	1.0	0.4
Potash	0.9	2.1	-0.3	3.2	-1.4	1.2	0.6	-	-

Straw: Mean yield 60.6 cwt. per acre

Ploughing deep									
-shallow	-3.4	-	-	-3.2	-3.6	-1.2	-5.6	-1.6	-5.2
Dung	2.7	2.9	2.5	-	-	3.3	2.1	5.3	-0.9
Superphosphate	0.2	2.4	-2.0	0.8	-0.4	-	-	-0.6	1.0
Potash	0.2	2.0	-1.6	3.8	-3.4	-0.6	1.0	-	-

48/Bc/1.1

GREEN MANURING EXPERIMENT

LB and LU - Woburn Stackyard, Series A, 1948

Treatments as given in 1936 Report, p. 203, with the exceptions that from 1946 onwards lupins replace tares, and rape replaces mustard as green manuring crops, while kale is replaced by winter cabbages as a testing crop. From 1944 onwards a top dressing of sulphate of ammonia has been applied to half the plots under barley, and in 1946 this top dressing was applied to the green manuring crops. The experiment is now in half replicate, according to the identity $I \equiv (R + C - M - F - T)DSNA$, A representing the top dressing of sulphate of ammonia.

Cultivations, etc.:

Upper Half. Cabbages

Red Clover and Italian Ryegrass undersown in barley: Apr 25, 1947. Failed owing to drought. Whole area ploughed : Nov 24 - Dec 8. Springtime harrowed: Feb 18-19, Mar 9. Harrowed: Mar 17. Rolloled: Mar 22. Rape sown, ryegrass and clover resown, sulphate of ammonia applied, harrowed in: Mar 24. Lupins drilled: Apr 2, 6. Rape plots dusted with Flea beetle dust: Apr 27. Lupin plots hoed: May 11. All plots hoed: May 20-24. Lupin plots hoed: June 8. Fallow plots cultivated: June 25. Dung and straw applied on appropriate plots, green manure rolled: June 28. Whole area ploughed: June 30. Harrowed: July 1. Basal manures and sulphate of ammonia applied, harrowed: July 2. Rolloled: July 3. Cabbages transplanted: July 5-8. Gaps filled in: July 14-15, July 31 - Aug 7. Hoed: Aug 16-20, Sept. 6-7, Oct 8. Harvested: Jan 24, Feb 11, 23, Mar 3. Variety: January King. Previous Crop: Barley.

Lower Half. Barley

Ploughed: Mar 3-9. Harrowed: Mar 9. Lime at 3 cwt. CaO per acre applied: Mar 10. Harrowed: Mar 11. Sulphate of ammonia applied to appropriate plots, seed drilled: Mar 12. Harrowed in: Mar 17. Rolloled: Mar 22. Dorset Mail Clover and Italian Ryegrass undersown on appropriate plots: Apr 16. Harrowed in: Apr 23. Whole area rolled: Apr 26. Hand weeded: June 15. Harvested: Aug 14. Variety: Plumage Archer. Previous Crop: Cabbages.

Standard errors per plot:

Cabbages:	total yield,	1.175 tons per acre or 15.7% (9 d.f.)
	plant number,	0.27 thousands per acre or 1.5% (9 d.f.)
Barley:	grain,	2.78 cwt. per acre or 12.4% (9 d.f.)
	straw,	3.26 cwt. per acre or 11.9% (9 d.f.)

48/Bc/1.2

Upper Half - Cabbages

Green manure crop	None	Lupins	Clover	Rape	Ryegrass	Mean
	Total Weight: tons per acre (± 0.588)					(± 0.263)
No Dung	6.02	7.61	7.41	4.76	5.98	6.36
Dung	8.12	9.37	9.67	7.89	8.18	8.65
No Straw	6.47	8.83	8.37	6.94	7.31	7.58
Straw	7.68	8.15	8.72	5.70	6.86	7.42
Sulph. amm.						
2 cwt./acre	6.47	6.56	7.80	5.21	6.44	6.50
4 cwt./acre	7.68	10.42	9.29	7.43	7.73	8.51
Sulph. amm. to barley						
absent	6.20	8.44	9.26	6.16	7.24	7.46
Present	7.95	8.54	7.82	6.48	6.93	7.54
Mean (± 0.415)	7.07	8.49	8.54	6.32	7.08	7.50
	Total number: thousands per acre (± 0.136)					(± 0.061)
No Dung	17.4	17.6	17.6	17.3	17.6	17.5
Dung	17.5	17.7	18.0	17.7	17.7	17.7
No Straw	17.2	17.7	17.7	17.4	17.7	17.5
Straw	17.6	17.6	17.9	17.6	17.5	17.6
Sulph. amm.						
2 cwt./acre	17.4	17.7	17.8	17.5	17.7	17.6
4 cwt./acre	17.4	17.7	17.8	17.6	17.6	17.6
Sulph. amm. to barley						
absent	17.5	17.7	17.9	17.5	17.6	17.6
Present	17.4	17.6	17.8	17.5	17.6	17.6
Mean (± 0.096)	17.4	17.7	17.8	17.5	17.6	17.6

48/Bc/1.3

Upper Half - Cabbages

Differential Responses

	Mean	Dung		Straw		Sulph. amm.		Sulph. amm. to barley	
		Abs.	Pres.	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Total Weight: tons per acre (± 0.531 Means ± 0.372)									
Dung	2.29	-	-	2.30	2.27	2.33	2.24	2.36	2.21
Straw	-0.16	-0.14	-0.17	-	-	-0.01	-0.30	-0.28	-0.03
Sulph. amm.	2.01	2.05	1.96	2.15	1.86	-	-	2.11	1.90
Sulph. amm. to barley	0.08	0.15	0.00	-0.04	0.20	0.18	-0.02	-	-

Total number: thousands per acre (± 0.12) Means (± 0.09)

Dung	0.2	-	-	0.2	0.1	0.2	0.1	0.1	0.2
Straw	0.1	0.1	0.0	-	-	0.1	0.0	0.1	0.0
Sulph. amm.	0.0	0.0	0.0	0.0	0.0	-	-	0.0	0.0
Sulph. amm. to barley	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-	-

48/Bc/1.4

Lower Half - Barley

Green Manure Crop	Lupins	Rape	Clover	Ryegrass	Fallow	Mean
Grain: cwt. per acre (± 1.39)						(± 0.62)
No Dung to Cabbages	21.7	20.7	23.3	16.9	22.9	21.1
Dung	25.6	21.9	26.6	22.1	22.6	23.8
No Straw to Cabbages	22.6	21.6	25.0	19.8	21.7	22.1
Straw	24.7	21.0	24.9	19.2	24.1	22.8
Sulph. amm. to Cabbages						
2 cwt./acre	23.0	22.1	24.9	17.3	21.5	21.6
4 cwt./acre	24.3	20.6	25.0	21.6	24.3	23.2
Sulph. amm. to Barley						
Absent	19.9	18.2	20.9	15.0	18.9	18.6
Present	27.4	24.5	29.0	23.9	26.8	26.3
Mean (± 0.98)	23.7	21.3	24.9	19.5	22.9	22.4
Straw: cwt. per acre (± 1.63)						(± 0.73)
No Dung to Cabbages	26.8	24.5	26.5	19.0	26.8	24.7
Dung	31.5	28.8	32.7	26.2	31.2	30.1
No Straw to Cabbages	28.2	26.4	28.0	23.0	28.1	26.7
Straw	30.1	26.9	31.2	22.2	30.0	28.1
Sulph. amm. to Cabbages						
2 cwt./acre	27.7	24.4	29.1	21.3	27.7	26.0
4 cwt./acre	30.6	28.9	30.1	23.8	30.3	28.7
Sulph. amm. to Barley						
Absent	25.6	22.8	25.0	19.3	24.4	23.4
Present	32.7	30.4	34.2	25.6	33.6	31.3
Mean (± 1.15)	29.2	26.6	29.6	22.6	29.0	27.4

48/Bc/1.5

Lower Half - Barley

	Mean	Dung		Differential Responses				Sulph. amm. to barley	
		Abs.	Pres.	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Grain: cwt. per acre (± 1.26 Means ± 0.88)									
Dung to Cabbages	2.7	-	-	1.4	3.9	4.3	1.0	3.3	2.0
Straw to Cabbages	0.6	-0.6	1.8	-	-	-0.6	1.8	1.3	-0.1
Sulph. amm. to Cabbages	1.4	3.0	-0.2	0.1	2.6	-	-	1.8	0.9
Sulph. amm. to Barley	7.8	8.4	7.1	8.5	7.0	8.2	7.3	-	-
Straw: cwt. per acre (± 1.47 Means ± 1.03)									
Dung to Cabbages	5.4	-	-	5.3	5.4	7.1	3.6	5.5	5.2
Straw to Cabbages	1.3	1.2	1.3	-	-	1.0	1.5	2.6	0.0
Sulph. amm. to Cabbages	2.7	4.4	0.9	2.4	2.9	-	-	2.6	2.7
Sulph. amm. to Barley	7.9	8.0	7.7	9.2	6.5	7.8	7.9	-	-

48/Bd/1.1

LEY AND ARABLE ROTATIONS

Woburn - Stackyard Series D, 1948

Details as given in 1938 Report, pp. 135-137, except that owing to the unsatisfactory crops obtained on kale plots in the years 1938-44 sugar beet has been substituted for kale from 1945 onwards.

Cultivations, etc.:

Block I. Ley. Second year. Grazed by sheep: May 3-21, June 13-17, July 9-13, Aug 17-25, Sept 7-11, Nov 4-10.

Lucerne. Second year. Harrowed three times: Nov 1. Hoed: Nov 3. First cut: July 9. Second cut: Sept 6. Third cut: Nov 12.

Wheat. Ploughed: Oct 7-8. Harrowed: Oct 9. Seed (Squarheads Master) drilled, harrowed: Oct 14, 15. Failure due to birds. Harrowed, seed redrilled: Oct 21. Plant failure due to damage by birds. Harrowed, sulphate of ammonia applied, seed (Atle) drilled, harrowed: Mar 10. Rolled: Mar 13. Harrowed: Apr 23. Rolled: Apr 26. Harvested: Sept 4. Variety: Atle. Previous crop: Potatoes.

Block II. Ley. Third year. Grazed by sheep: May 14-20, June 17-21, July 13-17, Aug 25-Sept 2, Sept 11-15, Nov 10-14.

Lucerne. Third year. Hoed: June 14-18. First cut: July 9. Second cut: Sept 6. Third cut (small crop only): Nov 12.

May. Grass and clover mixture under sown in wheat: Apr 17, 1947. Total failure due to drought. All wheat plots ploughed: Sept 30. Springtime harrowed:

Mar 1. Sulphate of ammonia applied, harrowed: Mar 10. Harrowed twice: Mar 18. Rolled: Mar 22. Seed sown:

Mar 23. Harrowed and rolled: Mar 25. Rolled: Mar 27. First cut: June 9. Second cut (small): Sept 6. Seeds

mixture: Giant Italian Ryegrass: (22 lb per acre) and Trifolium (27 lb per acre). Previous crop: Wheat.

Sugar beet. Ploughed: Sept 30, Mar 18. Harrowed: Mar 19. Rolled: Mar 27. Seed drilled: Apr 2. Nitrate

of soda applied: Apr 5. Rolled: Apr 6. Hoed: Apr 19, June 1-3. Singled: June 1-3. Hoed: June 8-July 2.

Lifted: Oct 5, 6. Variety: Klein E. Previous crop: Wheat.

Block III. Ley. First year. Ploughed: Sept 27-29. Springtime harrowed, harrowed: Mar 10. Rolled, harrowed, rolled: Mar 20. Artificials applied: Mar 23. Seed sown and harrowed in: Mar 25. Topped, toppings left on ground: July 9. Grazed by sheep: June 9-13, July 1-9,

48/Ba/1.2

July 17-22, Aug 9-17, Sept 2-7, Sept 15-16 (Plot 44 only),
Nov 29-Dec 4. Seeds mixture: S.23 Perennial Ryegrass
(21 lb per acre), S.143 Cocksfoot (12 lb per acre), late
flowering Montgomery Red Clover (6 lb per acre), S.100
White Clover (3 lb per acre). Previous crop: Barley.

Lucerne. First year. Ploughed: Sept 27-29.
Springtime harrowed, harrowed: Mar 10. Rolled, harrowed,
rolled: Mar 20. Artificials applied: Mar 23. Rolled,
seed sown: Mar 24. Failed, probably due to Flea beetle
attack. Harrowed and rolled: Mar 25. Cultivated:
May 15. Seed drilled: May 18. Dusted with Flea beetle
dust: June 11. Cleaned: June 13, 23. Hoed: June 22-
Sept 22. First cut: Sept 6. Second cut: Nov 12.

Variety: Provence. Previous crop: Barley.
Potatoes. Ploughed: Sept 27-29. Springtime
harrowed: Mar 10. Rolled, harrowed, rolled: Mar 20.
Rolled: Mar 27. Ridged: Apr 6-7. Artificials applied,
potatoes planted and covered in: Apr 9. Ridges harrowed:
May 8. Grubbed: May 28. Earthed up: June 21. Sprayed
with "Perinox": Aug 11. Sprayed to kill haulm: Sept 20.
Lifted: Sept 30. Variety: Majestic. Previous crop:
Barley.

Block IV. Potatoes. Ploughed: Oct 30-Nov 4. Springtime
harrowed: Mar 1, 18. Harrowed: Mar 20. Rolled: Mar 27.
Ridged: Apr 6-7. Dung applied: Apr 8. Potatoes planted,
artificials applied, covered in: Apr 9. Ridges harrowed
down: May 8. Grubbed: May 28. Earthed up: June 21.
Sprayed with "Perinox": Aug 11. Lifted: Oct 1, 2.
Variety: Majestic. Previous crop: Ley, lucerne, hay,
sugar beet.

Block V. Barley. Ploughed: Oct 9-10. Springtime harrowed:
Feb 27. Lime applied (14 cwt per acre, 82% Cao),
harrowed, sulphate of ammonia applied: Mar 4. Harrowed,
seed drilled: Mar 10. Rolled: Mar 13. Harvested:
Aug 16. Variety: Plumage Archer. Previous crop:
Potatoes.

Standard errors per plot:

Block V.

Barley, grain: whole plot, 1.61 cwt. per acre or 5.3%
sub-plot, 1.49 cwt. per acre or 4.9%
straw: whole plot, 2.68 cwt. per acre or 6.5%
sub-plot, 1.50 cwt. per acre or 3.6%

Block IV

Potatoes, total produce: whole plot, 0.626 tons per acre or
sub-plot, 0.903 tons per acre or
percentage ware: whole plot, 0.380
sub-plot, 0.435

All standard errors are estimated from 4 d.f.

48/Ba/1.3

Note: For the ley plots of blocks I, II, III, only the mean number of sheep days of grazing per acre and the mean number of sheep carried per acre for the year are given for each block in the summary tables. Each plot was grazed separately, but the number of grazing days is not sufficiently accurately determined for each plot to justify treatment comparisons.

48/Bd/1.4

Block I

<u>Ley.</u>	2nd Year.	Sheep days of grazing per acre	No. of sheep carried per acre for the year
Mean	1986		5.4

Lucerne. 2nd Year.

Yield of Lucerne Hay (85% dry matter): tons per acre

	1st crop	2nd crop	3rd crop	Total
No dung	2.10	1.74	0.47	4.32
Dung in 1945	2.01	1.77	0.52	4.30
Mean	2.06	1.76	0.49	4.30
Increase	-0.09	0.03	0.05	-0.02
<u>Previous Rotation</u>				
Lucerne	1.94	1.64	0.48	4.04
Arable with sugar beet	2.18	1.88	0.51	4.56

Wheat

	Grain: cwt. per acre	Straw: cwt. per acre
No dung	14.7	30.2
Dung in 1945	14.1	31.4
Mean	14.4	30.8
Increase	-0.6	1.2
<u>Previous Rotation</u>		
Ley	18.6	37.2
Lucerne	14.2	29.8
Arable with hay	12.7	27.0
Arable with sugar beet	12.1	29.0

48/Ba/1.5

Block II

Ley. 3rd Year.

	Sheep days of grazing per acre	No. of sheep carried per acre for the year
Mean	1551	4.3

Lucerne 3rd year.

Yield of Lucerne Hay (85% Dry Matter): tons per acre

	1st Crop	2nd Crop	3rd Crop	Total
No Dung	1.38	1.41	0.04	2.84
Dung in 1944	1.84	1.68	0.08	3.61
Mean	1.62	1.54	0.06	3.22
Increase	0.46	0.27	0.04	0.77
Previous Rotation				
Lucerne	1.40	1.30	0.06	2.75
Arable with hay	1.83	1.80	0.08	3.70

<u>Hay</u>	1st Crop	2nd Crop	Total
No Dung	1.04	0.20	1.24
Dung in 1944	1.34	0.24	1.58
Mean	1.19	0.22	1.41
Increase	0.30	0.04	0.34
Previous Rotation			
Lucerne	1.38	0.16	1.54
Arable with hay	1.00	0.28	1.28

Sugar Beet

	Clean Beet tons per acre	Tops tons per acre	Total Sugar cwt per acre	Sugar %
No Dung	10.40	10.54	35.8	17.21
Dung in 1944	9.64	9.00	34.7	18.02
Mean	10.02	9.78	35.2	17.62
Increase	-0.76	-1.54	-1.1	0.81
Previous Rotation				
Ley	10.41	9.54	36.6	17.62
Arable with sugar beet	9.63	10.01	33.8	17.62

48/Ba/1.6

Block III

Ley. 1st Year.

	Sheep days of grazing per acre	No. of sheep carried per acre for the year
Mean	1986	5.4

Potatoes

	Total tubers tons per acre	Percentage Ware
No Dung	14.90	96.7
Dung in 1946	17.13	97.4
Mean	16.01	97.0
Increase	2.23	0.7
Previous Rotation		
Ley	17.08	97.35
Lucerne	15.35	96.45
Arable with hay	15.22	97.30
Arable with sugar beet	16.42	97.00

Lucerne 1st Year.

Yield of Lucerne Hay (85% dry matter): Tons per acre

No Dung	0.32
Dung in 1946	0.66
Mean	0.49
Increase	0.34
Previous Rotation	
Lucerne	0.46
Arable with hay	0.53

48/Ed/1.7

Block IV

Potatoes

Previous Crop Rotation

	Ley	Lucerne	Arable with hay	Arable with sugar beet	Mean
Total tubers: tons per acre					
No Dung	15.86	16.37	14.19	13.63	15.01
Dung in 1948 (± 0.632) ⁽¹⁾	19.02	21.35	19.00	17.44	19.20
Mean (± 0.443)	17.44	18.86	16.59	15.54	17.11
Increase (± 0.903)	3.16	4.98	4.81	3.81	4.19 (± 0.452)
Percentage Ware					
No Dung	97.8	97.4	96.6	96.5	97.1
Dung in 1948 (± 0.35) ⁽¹⁾	97.8	97.7	97.1	97.2	97.5
Mean (± 0.27)	97.8	97.6	96.8	96.9	97.3
Increase (± 0.44)	0.0	0.3	0.5	0.7	0.4 (± 0.22)

Block V

Barley

	Ley	Lucerne	Arable with hay	Arable with sugar beet	Mean
Grain: cwt per acre					
No Dung	30.6	31.6	29.2	28.1	29.9
Dung in 1947 (± 1.36) ⁽¹⁾	32.5	28.3	33.1	31.3	31.3
Mean (± 1.14)	31.5	30.0	31.1	29.7	30.6
Increase (± 1.49)	1.9	-3.3	3.9	3.2	1.4 (± 0.75)

Standard error (1) for comparisons other than vertical ones.

Block V

48/Da/1.8

Barley

	Ley	Lucerne	Arable with hay	Arable with sugar beet	Mean
Straw: cwt. per acre					
No Dung (± 2.03) (1)	42.3	41.3	36.2	34.6	38.6
Dung in 1947	46.3	42.3	44.1	41.2	43.5
Mean (± 1.89)	44.3	41.8	40.2	37.9	41.0
Increase (± 1.50)	4.0	1.0	7.9	6.6	4.9 (± 0.75)

Standard error (1) for comparisons other than vertical ones

48/Be/1.1

WOBURN MARKET GARDEN EXPERIMENT

Globe Beet and Peas. First crops of 7th year.

The use of heavy dressings of organic manures for making a market garden soil, and the effect of sulphate of ammonia.

JRB and JPE - Lansome, 1948.

System of replication: 2 series, one of each crop, each consisting of 4 randomized blocks of 10 plots each, certain interactions being confounded with block differences.

Area of each plot: $1/80$ acre.

Treatments:

Organic manures: Dung, sewage sludge compost, sewage sludge (West Middlesex), and vegetable compost, each at 15 and 30 tons per acre.

Sulphate of ammonia: None, 0.2 cwt. N per acre on organic manure plots. None, 0.2, 0.4, 0.6 cwt. N per acre on plots without organic manure.

Basal manuring: Superphosphate, 0.4 cwt. P_2O_5 per acre.
Muriate of potash, 0.5 cwt K_2O per acre.

Cultivations, etc.:

Series A. Globe Beet.

Organic manures applied: Mar 31. Ploughed: Apr 1.
Harrowed: Apr 2. Ground lime applied to all plots receiving sulphate of ammonia (plots 1,7 at 33 cwt per acre, plots 24, 27 at 22 cwt per acre, other plots at 11 cwt per acre). Harrowed in: Apr 5. Rolled, basal manures and sulphate of ammonia dressings applied, seed drilled and harrowed in: Apr 6. Ring rolled: Apr 10. Dusted with Flea beetle dust: Apr 27. Hoed: May 6-10. Hoed and hand weeded: May 13-28. Harvested: July 6,12-13. Variety: Crimson Globe. Previous crop: Leeks.

Series B. Peas.

Organic manures applied: Jan 16-19. Ploughed: Jan 19-20. Springtime harrowed: Feb 17-19. Harrowed: Mar 15. Rolled, basal manures and sulphate of ammonia applied, peas drilled: Mar 16. Ring rolled: Mar 23. Hoed: Apr 29, May 10. Harvested: July 14-26. Variety: Kelvedon Wonder. Previous crop: Winter cabbage*.

Note: The winter cabbage crop failed owing to drought.

Standard errors per plot:

Peas, marketable produce, 11.88 cwt. per acre or 18.3%
Globe beet, weight of marketable produce, 1.272 tons per acre or 24.0%
total plant number, 4.01 thousands per acre or 12.3%
weight of bulbs, 0.719 cwt per acre or 26.4%

48/3e/1.2

Summary of Results

Organic manures	Level of manuring (tons/acre)	Sulphate of ammonia, cwt. N per acre				Mean
		None	0.2	0.4	0.6	

Green peas, marketable produce: cwt. per acre
(± 8.40 , means ± 5.94)

None		66.1	61.8	45.0	71.1	63.9*
Dung	15	96.1	75.0			85.5
Dung	30	79.3	41.3			61.8
Composted	15	75.7	85.0			80.4
sewage sludge	30	80.4	73.9			77.1
Sewage	15	47.1	46.8			47.0
sludge	30	35.7	38.2			37.0
Vegetable	15	85.4	84.3			84.8
compost	30	52.1	57.1			54.6

Globe beet, total produce: tons per acre
(± 0.900 , means ± 0.636)

None		1.30	2.92	3.86	2.37	2.11*
Dung	15	4.50	6.02			5.26
Dung	30	9.34	7.58			8.46
Composted	15	3.07	3.68			3.38
sewage sludge	30	4.72	6.19			5.45
Sewage	15	6.37	5.94			6.15
sludge	30	6.74	6.35			6.54
Vegetable	15	4.32	5.02			4.67
compost	30	6.85	8.78			7.82

Globe beet, weight of bulbs: tons per acre
(± 0.508 , means ± 0.360)

None		0.47	1.37	2.04	1.20	0.92*
Dung	15	2.04	3.20			2.62
Dung	30	5.01	4.12			4.56
Composted	15	1.39	1.77			1.58
sewage sludge	30	2.46	3.26			2.86
Sewage	15	3.25	3.08			3.17
sludge	30	3.50	3.30			3.40
Vegetable	15	2.10	2.62			2.36
compost	30	3.53	4.74			4.14

* These means are for 0.0 and 0.2 cwt. N per acre only

48/Be/1.3

Organic manures	Level of manuring (tons/acre)	Sulphate of ammonia, cwt. N per acre				Mean
		None	0.2	0.4	0.6	
Globe beet, total plant number: thous. per acre (± 2.84 , means ± 2.01)						
None		27.1	32.4	33.4	25.2	29.8*
Dung	15	32.8	31.6			32.2
Dung	30	42.8	29.4			36.1
Composted	15	29.6	35.0			32.4
sewage sludge	30	33.5	34.2			33.9
Sewage	15	37.6	29.9			33.8
sludge	30	28.8	27.2			28.0
Vegetable	15	37.0	39.0			38.0
compost	30	31.8	34.9			33.4

*These means are for 0.0 and 0.2 cwt. N per acre only

48/Be/2.1

WOBURN MARKET GARDEN EXPERIMENT

Leeks and Winter Cabbage. 2nd crops for 7th year

The use of heavy dressings of organic manures for making garden soil, and the effect of sulphate of ammonia.

JLE and JU - Lancaster 1948 - 9

System of replication: 2 series, one for each crop, each consisting of 4 randomized blocks of 10 plots each, certain interactions being confounded with block differences.

Area of each plot: $\frac{1}{80}$ acre.

Treatments:

Organic manures (applied to previous crops): Dung, sewage, sludge compost, sewage sludge (West Middlesex), and vegetable compost, each at 15 and 30 tons per acre.

Sulphate of ammonia: None, 0.4 cwt. N per acre on organic manure plots. None, 0.4, 0.8, 1.2 cwt. N per acre on plots without organic manure.

Basal manuring: None.

Cultivations, etc:

Series A. Winter Cabbage.

Ploughed and harrowed: July 15, 16. Rolled, sulphate of ammonia applied, cabbages planted: July 19, 20. Cabbages replanted where necessary: Aug 3 - 7. Hoed: Aug 24 - Sept 29. Cut: Jan 11, Feb 8, 15, 22. Variety: January King. Previous crop: Globe beet.

Series B. Leeks.

Ploughed: July 26, 27. Harrowed, rolled: July 27. Sulphate of ammonia applied, plots 48, 51, 72, 78 receiving only half dressings: July 30. Leeks planted: Aug 3 - 5. Hoed: Aug 24 - Sept 29. Second dressing of sulphate of ammonia to plots 48, 51, 72, 78: Sept 27. Harvested: Mar 16 - 29. Variety: Musselburgh. Previous crop: Peas.

Standard errors per plot:

Winter cabbages, marketable weight, \$0.828 tons per acre or 11.8%

total number, 0.372 thousands per acre or 2.1%

Leeks, total weight 14.40 cwt. per acre or 12.0%

total number, 0.879 thousands per acre or 2.1%

48/Be/2.2

Summary of Results

Organic manures	Level of manuring tons per acre	Sulphate of Ammonia cwt N per acre				Mean
		None	0.4	0.8	1.2	
Cabbages, Marketable Weight: tons per acre						
(± 0.586 Means ± 0.414)						
None		2.41	5.98	7.59	7.49	4.20*
Dung	15	3.36	6.66			5.01
Dung	30	8.14	7.86			8.00
Composted	15	4.98	6.91			5.94
sewage sludge	30	5.88	8.05			6.96
Sewage	15	8.34	9.66			9.00
sludge	30	9.77	9.88			9.82
Vegetable	15	5.68	6.99			6.33
compost	30	6.95	8.26			7.61
Cabbages, Total Number: thous. per acre						
(± 0.263 Means ± 0.166)						
None		17.6	17.9	18.0	18.2	17.8*
Dung	15	16.7	18.3			17.5
Dung	30	18.3	18.0			18.1
Composted	15	17.8	18.2			18.0
sewage sludge	30	18.1	17.8			17.9
Sewage	15	17.8	17.8			17.8
sludge	30	17.4	18.0			17.7
Vegetable	15	17.9	17.5			17.7
compost	30	18.0	18.3			18.2

* These means are for 0.0 and 0.4 cwt N. per acre only.

48/Be/2.3

Organic Manures	Level of manuring tons per acre	Sulphate of Ammonia cwt. N per acre			Mean
		None	0.4	0.8	

Leeks, Total Weight: cwt per acre

(± 10.2 Means ± 7.20)

None		66.3	113.7	107.3	116.2	90.0*
Dung	15	114.4	113.6			114.0
Dung	30	139.6	143.7			141.6
Composted	15	103.8	124.1			114.0
sewage sludge	30	120.5	113.2			116.8
Sewage	15	127.6	125.5			126.5
sludge	30	131.8	163.5			147.6
Vegetable	15	109.0	106.2			107.6
compost	30	124.4	126.8			125.6

Leeks, Total Number: thous. per acre

(± 0.622 Means ± 0.440)

None		40.9	42.6	38.7	41.1	41.8*
Dung	15	41.8	41.9			41.9
Dung	30	42.8	40.6			41.7
Composted	15	41.5	41.9			41.7
sewage sludge	30	41.1	40.6			40.8
Sewage Sludge	15	40.9	39.9			40.4
" "	30	41.5	42.4			41.9
Vegetable compost	15	43.5	40.5			42.0
" "	30	42.3	42.1			42.2

* These means are for 0.0 and 0.4 cwt N per acre only.

48/Bf/1

GRAZING EXPERIMENT

Highfield - 1948

The effects of cake feeding and equivalent manures on the nutritive value of the herbage.

For details see 1937 Report, pp. 24-26

Area of each plot: Plots 4, 5, 6: 4.93 acres

Cultivations: All plots topped to cut off thistles: June 17-18

Grazing periods: Cattle: May 4 - June 30, and Nov. 4 - Dec. 3
 Sheep: June 8 - Sept. 22

Block II

Plot	5	4	6
	No Manure	Cake 1938, 40, 43, 46	Equivalent Manures
	Live weight increases: lb per acre		
Cattle	631	480	566
Sheep	116	134	128
	Grazing days per acre		
Cattle	144	104	160
Sheep	327	289	336
	Starch equivalent: lb per acre		
Cattle	2285	1720	2236
Sheep	889	911	922

48/Ca/1.1

WHEAT

Control of "Eyespot"

The effects of sulphate of ammonia, of rate and depth of sowing and of spraying, on the sixth successive Wheat crop.

RW - Little Knott 1948

System of replication: 3 x 3 x 3 design in 6 blocks of 9 plots each, certain three-factor interactions and the effect of spraying being confounded with block differences.

Area of each plot 0.0151 acre.

Treatments:

Sulphate of ammonia: None, 0.4, 0.8 cwt N per acre applied as a top-dressing in March (N_0, N_1, N_2).
 Depth of sowing: Shallow ($\frac{1}{2}$ " $_0, 1, 2$), normal ($1\frac{1}{2}$ " $_0, 1, 2$), deep (3" $_0, 1, 2$).
 Rate of sowing: $1\frac{1}{2}, 2\frac{1}{2}$ or $3\frac{1}{2}$ bushels per acre (R_0, R_1, R_2).
 Spraying: 3 Blocks sprayed with sulphuric acid ($12\frac{1}{2}\%$ by volume B.O.V. at 100 gallons per acre) before germination and again in spring.

Basal Manuring: 3 cwt per acre superphosphate and 1 cwt per acre muriate of potash drilled across the plots.

Cultivations etc:

Ploughed: Sept 8 - 13. Basal Manures applied: Oct 8. Cultivated: Oct 17. Ring-rolled: Oct 18. Springtined, rolled and springtined and ring-rolled: Oct 21. Seed drilled: Oct 22. Harrowed in: Oct 23. Sprayed with B.O.V.: Nov 1 and again Mar 8. Sulphate of Ammonia applied: Mar 15. Ring-rolled: Apr 28. Sprayed against weeds: May 14. Hand pulled rye: May 21. Hand pulled wild oats and other weeds: June 15 - 16. Hand pulled wild oats: various days: June 20 - 30. Harvested: Aug 21. Variety: Squareheads Master ($13/4$).
 Previous crop: Wheat.

Standard errors per plot:

Grain: 2.02 cwt per acre or 10.4%	(24 d.f.)
Straw: 5.19 cwt per acre or 7.9%	(24 d.f.)
% Eyespot, (transformed values).	
Sprayed blocks - 2.57	6 d.f.)
Unsprayed - 6.48	6 d.f.)
% Take All (transformed values) - 8.68	24 d.f.)

Note: The analysis of diseased wheat has been carried out on percentages transformed to degrees and all tests of significance should be applied to the transformed values.

Grain: cwt per acre (± 1.17 Means ± 0.67)

48/Ca/1.2

	Unsprayed				Sprayed				Effect of Spraying (± 0.95)(1)
	R ₀	R ₁	R ₂	Mean	R ₀	R ₁	R ₂	Mean	
D ₀	15.0	17.8	16.0	16.3	25.8	22.9	23.4	24.0	7.7
D ₁	16.4	15.6	13.5	15.2	23.5	21.2	22.6	22.4	7.2
D ₂	17.2	17.2	12.9	15.8	24.9	22.8	20.9	20.9	7.1
	N ₀	N ₁	N ₂		N ₀	N ₁	N ₂		
D ₀	17.0	14.2	17.7		20.4	25.6	26.1		
D ₁	14.4	15.8	15.4		17.0	23.7	26.7		
D ₂	15.2	15.7	16.5		15.7	25.1	27.9		
R ₀	17.7	13.6	17.3	16.2	18.9	26.2	29.1	24.7	8.5
R ₁	17.0	16.1	17.6	16.9	19.1	23.5	24.3	22.3	5.4
R ₂	11.7	16.1	14.7	14.2	15.0	24.7	27.2	22.3	8.1
Mean	15.5	15.2	16.5	15.8	17.7	24.8	26.9	23.1	
Effect of spraying (± 0.95)(1)					2.2	9.6	10.4		

Straw: cwt per acre (± 3.13 Means ± 1.81)

	Unsprayed				Sprayed				Effect of Spraying (± 2.45)(1)
	R ₀	R ₁	R ₂	Mean	R ₀	R ₁	R ₂	Mean	
D ₀	55.3	62.6	55.4	57.7	81.1	74.9	78.2	78.1	20.4
D ₁	59.3	58.3	52.1	56.6	75.0	69.9	72.7	72.5	15.9
D ₂	56.5	63.7	48.6	56.3	76.2	72.2	70.0	72.8	16.5
	N ₀	N ₁	N ₂		N ₀	N ₁	N ₂		
D ₀	55.4	54.6	63.3		71.0	80.3	83.0		
D ₁	49.4	56.8	63.5		57.8	74.5	85.3		
D ₂	51.9	55.8	61.1		54.5	78.5	85.4		
R ₀	56.7	51.7	62.7	57.0	64.3	79.2	88.8	77.4	20.4
R ₁	57.0	59.3	68.3	61.5	61.1	76.2	79.7	72.3	10.8
R ₂	42.9	56.2	56.9	52.0	57.9	77.9	85.1	73.6	21.6
Mean	52.2	55.7	62.6	56.9	61.1	77.8	84.5	74.5	
Effect of spraying (± 2.45)(1)					8.9	22.1	21.9		

(1) Standard error for comparison between main effects only.

Percentage Straws showing Severe Eyespot at Harvest
(means from transformed values)

48/0a/1.0

	Unsprayed				Sprayed				Effect of Spraying
	R ₀	R ₁	R ₂	Mean	R ₀	R ₁	R ₂	Mean	
D ₀	51.6	57.5	42.5	50.5	10.3	15.0	9.7	11.4	-39.1
D ₁	62.3	38.6	66.5	56.0	11.4	12.1	11.3	11.7	-44.3
D ₂	35.3	58.0	52.2	48.4	10.9	15.0	9.1	12.2	-36.2
	N ₀	N ₁	N ₂		N ₀	N ₁	N ₂		
D ₀	37.2	62.8	51.8		12.3	12.8	9.7		
D ₁	35.7	58.8	72.3		6.4	13.3	16.2		
D ₂	37.5	55.4	52.8		10.3	16.1	10.7		
R ₀	41.3	56.0	51.8	49.7	9.1	13.8	11.8	11.4	-38.3
R ₁	36.8	52.8	64.3	51.4	14.4	14.8	13.0	14.0	-37.4
R ₂	32.3	68.0	61.0	53.8	5.8	13.7	11.4	10.1	-43.7
Mean	36.8	59.0	59.0	51.6	9.4	14.0	12.1	11.8	
	Effect of Spraying				-27.4	-45.0	-46.9		

(Transformed values)

	Unsprayed (± 3.74 Means ± 2.16)				Sprayed (± 1.48 Means ± 0.86)				Effect of Spraying (± 2.32) ⁽¹⁾
	R ₀	R ₁	R ₂	Mean	R ₀	R ₁	R ₂	Mean	
D ₀	45.9	49.3	40.7	45.3	18.7	22.8	18.1	19.8	-25.5
D ₁	52.1	38.4	54.6	48.4	19.8	20.4	19.7	20.0	-28.4
D ₂	36.5	49.6	46.2	44.1	21.0	22.8	17.6	20.5	-23.6
	N ₀	N ₁	N ₂		N ₀	N ₁	N ₂		
D ₀	37.6	52.4	46.0		20.5	20.9	18.1		
D ₁	36.7	50.1	58.3		14.6	21.4	23.8		
D ₂	37.8	48.1	46.5		18.7	23.7	19.1		
R ₀	40.0	48.4	46.0	44.8	17.6	21.8	20.1	19.8	-25.0
R ₁	37.4	46.6	53.3	45.8	22.3	22.6	21.1	22.0	-23.8
R ₂	34.6	55.6	51.4	47.2	13.9	21.7	19.8	18.5	-28.7
Mean	37.4	50.2	50.2	45.9	17.9	22.0	20.4	20.1	
	Effect of Spraying (± 2.32) ⁽¹⁾				-19.5	-28.2	-29.8		

(1) Standard error for comparison between main effects only.

Percentage Straws showing "take-all" at Harvest 48/Ca/1.4
(means from transformed values)

Unsprayed					Sprayed				Effect of Spraying
	R ₀	R ₁	R ₂	Mean	R ₀	R ₁	R ₂	Mean	
D ₀	17.9	23.3	19.8	20.3	28.9	28.1	27.0	28.0	7.7
D ₁	6.6	23.3	30.8	19.0	29.8	18.7	36.3	28.0	9.0
D ₂	14.5	20.6	16.6	17.1	20.6	34.7	41.3	32.0	14.9
	N ₀	N ₁	N ₂		N ₀	N ₁	N ₂		
D ₀	35.5	17.1	10.8		37.5	26.5	20.6		
D ₁	37.3	9.0	14.4		50.2	19.1	17.6		
D ₂	39.2	15.0	4.4		42.2	37.7	17.4		
R ₀	23.2	20.4	1.3	12.5	41.2	24.0	15.8	26.3	13.8
R ₁	34.2	17.0	17.4	22.4	30.2	27.2	23.5	26.8	4.4
R ₂	55.8	5.5	14.4	22.0	59.0	31.5	16.6	34.7	12.7
Mean	37.3	13.6	9.3	18.9	43.2	27.5	18.6	29.2	
Effect of Spraying					5.9	13.9	9.3		

(Transformed values)
(± 5.01 Means ± 2.89)

Unsprayed					Sprayed				Effect of Spraying (± 4.09)(1)
	R ₀	R ₁	R ₂	Mean	R ₀	R ₁	R ₂	Mean	
D ₀	25.0	28.9	26.4	26.3	32.5	32.0	31.3	31.9	5.1
D ₁	14.9	28.9	33.7	25.8	33.1	25.6	37.1	31.9	6.1
D ₂	22.4	27.0	24.1	24.5	27.0	36.1	40.0	34.4	9.9
	N ₀	N ₁	N ₂		N ₀	N ₁	N ₂		
D ₀	36.6	24.5	19.2		37.8	31.0	27.0		
D ₁	37.7	17.5	22.3		45.1	25.9	24.8		
D ₂	38.7	22.8	12.0		40.5	37.9	24.7		
R ₀	28.8	26.9	6.5	20.7	39.9	29.3	23.4	30.9	10.2
R ₁	35.8	24.4	24.7	28.3	33.3	31.4	29.0	31.2	2.9
R ₂	48.3	13.5	22.3	28.0	50.2	34.1	24.1	36.1	8.1
Mean	37.7	21.6	17.8	25.7	41.1	31.6	25.5	32.7	
Effect of Spraying					3.4	10.0	7.7		

(1) Standard error for comparison between main effects only.

48/Ca/2.1

WHEAT

Residual effects of various dungs, of additional straw to dungs, of rotted bracken and of straw with sulphate of ammonia added.

RW - Great Harpenden II 1948

System of replication: The design was intended to be a 5 x 5 lattice square in 3 replicates, but owing to the interchange of a pair of treatments in two of the three blocks, the lattice design was not attained, and it was impracticable to treat the experiment as other than 3 randomized blocks of 25 plots.

Area of each plot: 0.0171 acre.

Treatments: Applied in 1947 to potatoes. Of the 25 plots in each replicate, 3 received no organic manures, and the remaining 22 were treated with the following organic manures, applied at two rates: rotted bracken (B) straw with sulphate of ammonia added (A), and nine dungs:- from boxes:- stored (2 months) made with normal and heavy litter (W and X); from straw bale yards:- fresh, made with normal and heavy litter (Y and Z), stored (12 months), made with normal and heavy litter (R and S) and fresh (low ration and low ration plus sulphate of ammonia to straw) (T and V); from sunken yards:- stored (12 months) commercial dung (L).

Rates of application: The commercial dung (L) and rotted bracken (B) at 8 and 16 tons per acre, the straw with sulphate of ammonia (A) at 2 and 4 tons of chaffed straw per acre plus 0.3 and 0.6 N per acre as sulphate of ammonia, the stored normal dung from boxes (W) at the equivalent of 8 and 16 tons per acre weighed before storing, dung X, Y, Z, R and S, at weights produced by the same quantity of feeding stuffs as 8 and 16 tons of fresh normal dung from bullock boxes, and dungs T and V at the same rates as Z.

Dungs	Actual rates of application Tons per acre		Litter straw lbs/head/day
	Level 1	Level 2	
W	5.37	10.74	
X	4.79	9.58	9.0
Y	8.66	17.32	18.0
Z	10.78	21.55	9.2
R	6.79	13.57	17.1
S	7.76	15.51	9.8
TV	10.78	21.55	19.7
			16.1

48/Ca/2.2

Basal Manuring: 2 cwt. Sulphate of Ammonia per acre

Cultivations etc: Ploughed: Oct. 31 -- Nov. 4. Rolled and
springtime harrowed: Nov. 4 and again Nov. 5. Seed drilled:
Nov. 6. Harrowed in: Nov. 7. Ring rolled: Apr. 23.
Sulphate of ammonia drilled: early May. Cut and shocked:
Aug. 10-12. Reset shocks: Aug. 18. Raked and carted in:
Aug. 19-20. Threshed Sept. 6-10. Variety: Bersee.
Previous crop: Potatoes.

Standard error per plot: Grain 2.52 cwt. per acre or 5.75%
(50a.f.)

48/Oa/2.3

Organic Manure	Grain: cwt. per acre Level of organic		Straw: cwt. per acre Level of organic	Mean
	1	2		
	(±1.45)	(±1.03)		
None	42.5	41.9(1)	54.1	54.1
Stored (bullock boxes) normal litter	45.2	43.9	56.5	56.5
Stored (bullock boxes) heavy litter	44.9	43.2	57.5	57.5
Fresh (straw bala yards) normal litter	42.6	42.5	55.1	55.1
Fresh (straw bala yards) heavy litter	46.9	45.3	55.5	55.5
Stored (straw bala yards) normal litter	45.2	44.0	55.5	55.5
Stored (straw bala yards) heavy litter	44.1	43.3	55.2	55.2
Fresh (straw bala yards low feeding)	43.6	42.6	56.4	56.4
As above with Sulphate of Ammonia	44.6	44.7	57.0	57.0
Stored (Sunken yard)	44.8	45.1	59.0	59.0
Rotted Bracken	44.4	43.1	54.1	54.1
Straw with Sulphate of Ammonia	49.1	46.8	67.5	62.7
Mean	43.0	43.8	56.3	57.4

Standard error (1): ± 0.84

WHEAT

Wireworm Experiment (1)

The effect of various insecticides, and their methods of application.

RW - Little Hoos 1948

System of replication: 3 randomized blocks of 9 plots each

Area of each plot: 0.0289 acre

Treatments:

None

D.D. injected 400 lbs per acre.

Ethylene Dibromide 4.1% solution injected 15 gallons per acre.

D.D.T. dust combine drilled $\frac{5}{4}$ cwt per acre.

Gammexane; broadcast 2 cwt per acre, combine drilled $\frac{3}{4}$ cwt per acre, or applied as seed dressing.

Basal manuring: $2\frac{3}{4}$ cwt per acre sulphate of ammonia.

Cultivations etc.: Floughed: Sept 16-27. Harrowed and rolled: Oct 7.

DD and Ethylene Dibromide injected: Oct 10. Seed drilled, and remaining

treatments applied: Oct 29-30. Harrowed in: Oct 30. Harrowed: Mar 30.

Ring rolled: Apr 1. Sulphate of ammonia drilled: May 5. Hand weeded:

June 5, 7-8, 22-23, and various days June 28 - July 23. Harvested:

Aug 17. Variety: Bersee. Previous crop: Linseed.

Standard errors per plot:

Grain, 2.62 cwt per acre or 12.6% (18 d.f.)

Straw, 6.37 cwt per acre or 13.5% (18 d.f.)

	Un- treated	DD In- jected	Ethylene Dibromide Injected	DDT Dust Drilled	Gammexane Broad- cast	Drilled	Treated seed	Mean
Grain: cwt. per acre								
Mean Yield (± 1.51)	8.9 ⁽¹⁾	28.3	32.1	20.7	30.6	24.8	24.0	20.8
Increase (± 1.74)		19.4	23.2	11.8	21.7	15.9	15.1	
Straw: cwt. per acre								
Mean Yield (± 3.68)	22.8 ⁽²⁾	64.1	71.6	46.0	65.9	54.2	54.6	47.2
Increase (± 4.25)		41.3	48.8	23.2	43.1	31.4	31.8	

Standard errors (1) ± 0.87
 (2) ± 2.12

48/Ca/4.1

WHEAT

Wireworm Experiment (2)

The effect of treatment of seed with gammexane and of three strengths of gammexane dust.

RW - Little Hoos 1948

System of replication: 3 replicates of 3 incomplete randomized blocks of 3 plots each.

Area of each plot: 0.0289 acre

Treatments:

None.

Seed treated with gammexane dressing.

Gammexane dust, $\frac{1}{4}$, $\frac{1}{2}$ and 1 cwt per acre; combine drilled with seed (filler added where necessary to make total dressing of 1 cwt per acre)

Basal Manuring: $2\frac{3}{4}$ cwt per acre sulphate of ammonia.

Cultivations etc:

Ploughed: Sept 16-27. Harrowed and rolled: Oct 7. Seed and gammexane dust drilled: Oct 29-30. Harrowed in: Oct 30.

Harrowed: Mar 30. Ring rolled: Apr 1. Sulphate of ammonia drilled: May 5. Hand weeded: June 5, 7-8, 22-23, and various days June 23-July 23. Harvested: Aug 17.

Variety: Barsee. Previous crop: Linseed.

Standard errors per plot:

Grain: Block error, 1.61 cwt per acre or 7.15% (4d.f.)
Plot error, 2.28 cwt per acre or 10.1% (16d.f.)

Straw: Block error, 2.29 cwt per acre or 4.63% (4d.f.)
Plot error, 4.88 cwt per acre or 9.86% (16d.f.)

48/Ca/5.1

SPRING WHEAT

The effects of inoculation and wetting of seed, and of sulphate of ammonia.

RW - Long Hoos 1948

System of replication: 3 randomized blocks of 6 plots each.

Area of each plot: 0.0158 acre.

Treatments:

Seed: untreated, inoculated by wet method, wetted but not inoculated.

Sulphate of ammonia: None, 0.3 cwt. N per acre.

Note: The inoculum was a bacterial culture of French origin.

Basal manuring: 3 cwt. Superphosphate and 1 cwt. muriate of potash per acre.

Cultivations, etc.: Ploughed: Sept. 30 - Oct. 2. Rolled and springtine harrowed: Mar. 10. Sulphate of ammonia applied and basal manuring broadcast by machine: Mar. 12. Harrowed and seed drilled: Mar. 13. Harrowed in: Mar. 15. Rolled: Mar. 20. Weeds hand pulled: June 8. Harvested: Aug. 24. Variety: Bersee.

Standard errors per plot (10 d.f.): Grain, 3.35 cwt. per acre
or 17.1%
Straw, 3.00 cwt. per acre
or 7.20%

48/0a/5.2

	Seed Untreated	Seed inoculated (wet method)	Seed wetted not inoculated	Mean
Grain: cwt. per acre				
Mean (± 1.37)	19.8	18.5	20.5	19.6
Sulphate of Ammonia:				
Absent (± 1.93)	16.5	17.2	17.8	17.2
Present	23.2	19.8	23.1	22.0
Response (± 2.74)	6.7	2.6	5.3	4.8 (± 1.58)
Straw: cwt. per acre				
Mean (± 1.22)	41.3	41.3	42.3	41.6
Sulphate of Ammonia:				
Absent (± 1.73)	38.4	38.6	38.7	38.6
Present	44.2	44.1	45.8	44.7
Response (± 2.45)	5.8	5.5	7.1	6.1 (± 1.41)

48/Oa/6.1

SPRING-WHEAT VARIETY TRIAL

The comparison of eight varieties of spring wheat, and the effect on them of three levels of sulphate of ammonia.

RW - Little Knott 1948.

System of replication: 2 randomized blocks of 8 plots, plots split into 3 for the application of sulphate of ammonia.

Area of each sub-plot: 0.022 acre.

Treatments.

Whole plots: Varieties: Atle, Meteor, Fylgia, Extra Kolben II Bersee, Vilmorin 27, April Bearded, Vilmorin 29

Sub-plots: Sulphate of ammonia; none, 0.4, 0.8 cwt N per acre.

Basal dressing: None.

Cultivations, etc.: Ploughed: various days Dec. to Jan.
Springtime harrowed: Mar. 4, and again Mar. 11. Tooth harrowed: Mar. 15 and again Mar. 16. Sulphate of Ammonia applied: Mar. 16. Seed drilled: Mar. 18. Harrowed in: Mar. 19. Rolled: Mar. 23. Sprayed against weeds: May 19. Weeds pulled: June 7. Wild oats pulled: June 20-30. Harvested North block except Vilmorins by binder: Sept 7. Harvested Vilmorins North block: Sept. 22. Harvested South block by combine: Sept. 27. Previous crop: Kale

Standard Errors per plot: Grain

Whole plot: 1.99 cwt. per acre or 7.53%

Sub-plot: 1.88 cwt. per acre or 7.14%

Grain: cwt. per acre

Sulphate of ammonia	II						Mean (±0.47)
	Atle	Meteor	Fylgia	Kolben	Bersee	Vilmorin 27	
None	30.6	24.3	21.6	27.0	29.2	24.5	24.5
0.4 cwt. N per acre	31.4	25.2	24.6	27.7	30.0	24.0	25.9
0.8 cwt. N per acre	31.5	24.8	25.5	25.5	33.4	28.2	23.3
Mean	31.2	24.8	23.9	26.7 (±1.41)	30.8	25.6	24.5

Standard errors for body of table: ±1.33 for vertical comparisons only
±1.78 for all other comparisons

Straw: cwt. per acre[#]

Sulphate of ammonia	II						Mean
	Atle	Meteor	Fylgia	Kolben	Bersee	Vilmorin 27	
None	51.5	54.1	55.3	52.8	43.8	80.0	60.6
0.4 cwt. N per acre	55.6	53.4	52.5	56.8	53.5	72.2	61.1
0.8 cwt. N per acre	58.2	54.9	54.3	58.4	59.3	81.8	63.9
Mean	55.1	54.1	54.1	56.0	52.2	78.0	61.9

1 Block (North) only

48/Cb/1.1

SPRING SOWN CEREAL EXPERIMENT

Comparison of barley, oats and two varieties of wheat, and of the effects on them of four levels of sulphate of ammonia, of superphosphate, and of muriate of potash.

RV - Long Hoos III, 1948

System of replication: 4 randomized blocks of four plots each, each plot being split into 4, certain first-order interactions of artificials being confounded with whole plots.

Area of each sub-plot: 0.0191 acre.

Treatments:

Whole plots: Crops:- Oats (S.84), wheat (Atle and Bersce) and barley (Plumage Arthur)

Sub-plots: Sulphate of ammonia: None, 0.3, 0.6, 0.9, cwt N per acre
Superphosphate: None, 0.6 cwt. P_2O_5 per acre
Muriate of potash: None 0.6 cwt. K_2O per acre

Basal manuring: None.

Cultivations, etc.: Ploughed Sept.30 - Oct.2. Rolled and Springtime harrowed: Mar.10. Artificials applied: Mar.13.
Harrowed: Mar.15. All seed drilled and harrowed in: Mar.17.
Rolled: Mar.20. Hand weeded: June 7-8. All crops harvested: Aug. 23-24. Previous crop: Barley.

Standard errors: (grain):

per whole plot, 1.49 cwt. per acre or 6.9% (6.d.f.)
per sub-plot, 2.82 cwt. per acre or 13.0% (24 d.f.)

48/Cb/1.2

	Grain: cwt. per acre			Straw: cwt. per acre			
	Oats	Wheat (Atle)	Wheat (Bersee)	Oats	Wheat (Atle)	Wheat (Bersee)	Barley
Mean	19.0	21.6 (±0.75)	19.1	46.1	46.0	43.8	36.4
Sulphate of ammonia		(a) and (b)					
None	14.6	18.2	13.4	44.2	37.4	33.5	27.4
0.3 cwt. N per acre	19.6	20.8	20.0	48.6	44.0	41.2	35.2
0.6 cwt. N per acre	20.9	23.4	20.8	47.3	50.3	50.6	40.6
0.9 cwt. N per acre	20.9	24.1	22.2	44.4	52.5	49.9	42.6
Superphosphate		(±1.03) [‡]					
Absent	18.3	22.3	19.1	46.5	46.1	42.9	35.7
Present	19.7	20.9	19.1	45.8	46.0	44.7	37.1
Response	1.4	-1.4	0.0	-0.7	-0.1	1.8	1.4
Muriate of Potash		(±1.03) [‡]					
Absent	19.5	21.8	17.8	46.7	45.2	43.9	35.8
Present	18.5	21.4	20.4	45.6	46.9	43.7	37.1
Response	-1.0	-0.4	2.6	-1.1	1.7	-0.2	1.3

Standard errors: (a) ± 1.41 (for vertical comparisons only)
 (b) ± 1.43 (for all other comparisons)

[‡] Standard error for use in horizontal comparisons only

48/Cc/1.1

BEANS

The comparison of four varieties of beans and the effects on them of two methods of placement, two rates and two times of sowing.

RE - Great Field I 1948

System of replication: 4 randomized blocks of 8 plots each, the effect of time of sowing and certain high order interactions being confounded with block differences.

Area of each plot: 0.0286 acre.

Treatments:

Varieties: Giant, Essex, Lines and Wilts.

Methods of placement: Seed broadcast before ploughing, dropped in furrow during ploughing.

Rate of sowing: 2 cwt, 3 cwt seed per acre.

Time of sowing: Early, Oct 28th. Late, Nov 18th.

Basal dressings: Dung: 10 tons per acre

Superphosphate 2 cwt per acre drilled across the plots

Muriate of potash: 2 cwt per acre

Nitrochalk: 2 cwt per acre.

Cultivations etc: Superphosphate and Muriate of Potash applied: Oct 20.

Dung applied, ploughed in, beans placed in furrows, 1st sowing "Early":

Oct 27-29. Tractor rolled and disced each way after ploughing: Oct 29.

Chalk 45 cwt per acre applied: Nov 6. Dung spread, ploughed in beans

placed in furrows, 2nd sowing "Late": Nov 17-18. Harrowed "Early"

plots: Mar 16. Ring rolled all plots: Mar 30. Harrowed late plots:

Mar 31. Nitrochalk applied: May 4. Harvested: Aug 25. Previous

crop: Linseed.

Standard errors per plot :

Grain: 2.62 cwt per acre or 10.7% (11 d.f.)

Straw: 4.40 cwt per acre or 11.2% (11 d.f.)

4.8/Cc/1.2

Grain: cwt per acre

	Giant	Essex	Lincoln -shire	Wilts	Mean
(±0.92) Mean	23.8	26.5	24.4	23.2	24.5
(±1.31) Broadcast	23.2	25.3	24.4	24.1	24.3
(±1.31) Ploughed In	24.3	27.7	24.3	22.3	24.6(±0.65)
(±1.85) Ploughed In -Broadcast	1.1	2.4	-0.1	-1.8	0.3(±0.92)
Seed rate per acre					
(±1.31) 2 cwt	21.5	24.1	22.8	22.1	22.6(±0.65)
(±1.31) 3 cwt	26.1	28.9	26.0	24.3	26.3(±0.65)
(±1.85) 3 cwt - 2 cwt	4.6	4.8	3.2	2.2	3.7(±0.92)
(±1.85) ^{xx} Early-Late Sowing	4.1	8.0	7.2	6.0	

	Seed rate per acre		Difference	Early- Late Sowing
	2 cwt	3 cwt		
(±0.92) Broadcast	22.4	26.1	3.7(1)	7.1
(±0.92) Ploughed in	22.8	26.5	3.7(1)	5.5
(±1.31) Difference	0.4	0.4	0.0(2)	-1.6(2)
(±1.85) Early-Late Sowing	7.3	5.3	-2.0(2)	

Standard errors (1) ±1.31

(2) ±1.85

^{xx} Standard error for comparison between main effects only

48/Cc/1.3

Straw: cwt per acre

	Giant	Essex	Lincoln -shire	Wilts	Mean
(±1.56) Mean	39.6	38.5	38.7	40.3	39.3
(±2.20) Broadcast	39.3	36.8	39.8	40.2	39.0 (±1.10)
Ploughed In	39.9	40.3	37.7	40.4	
(±3.11) Ploughed In -Broadcast	0.6	3.5	-2.1	0.2	0.6 (±1.56)
Seed rate per acre					
(±2.20) 2 cwt	36.4	34.9	36.9	37.3	36.4 (±1.10)
3 cwt	42.8	42.1	40.6	43.3	
(±3.11) 3 cwt - 2 cwt	6.4	7.2	3.7	6.0	5.8 (±1.56)
(±3.11) [#] Early-Late Sowing	7.5	8.2	9.2	11.6	

	Seed rate per acre 2 cwt	3 cwt	Difference	Early- Late Sowing
(±1.56) Broadcast	36.7	41.3	4.6(1)	10.8
Ploughed In	36.0	43.1	7.1(1)	7.4
(±2.20) Ploughed In -Broadcast	-0.7	1.8	2.5(2)	-3.4(2)
Early-Late Sowing	8.9	9.3	0.4(2)	

Standard Errors (1) (±2.20)

(2) (±3.11)

[#] Standard error for comparison between main effects only

48/Ca/1.1

POTATOES.

Effects of various dungs, of additional straw to dung, of rotted bracken, and of sulphate of ammonia and muriate of potash.

RP - Sawyers II, 1948

System of replication: 5 x 5 lattice square in 3 replicates, plots split into two for application of nitrogen and potash, the interaction of these being confounded with whole plots.

Area of each sub-plot 0.0126 acre.

Treatments:

Of the 25 whole plots in each replicate, 3 received no organic manures, and the remaining 22 were treated with the following organic manures, applied at two rates: rotted bracken (B) and ten dungs: from bullock boxes:- fresh, made with normal and heavy litter (W and X), and stored (12 months under cover) made with normal and heavy litter (R and S): from straw bale yards:- fresh made with normal and heavy litter (Y and Z), stored (12 months in open) made with normal and heavy litter (A and K) and stored (12 months in open) low ration, and low ration plus sulphate of ammonia to straw (T and V).

Rates of application: The rotted bracken (B) and the fresh normal dung from boxes (W) at 8 and 16 tons per acre, dungs X, Y, Z, R, S, A and K at weights produced by the same quantity of feeding stuffs as 8 and 16 tons of fresh normal dung from boxes, and dungs T and V at the same rates as Z.

48/Ca/1.2

		Actual rates of application		Litter Straw
		Tons per acre		lbs/head/day
		Level 1	Level 2	
Dungs	W	8.00	16.00	10.6
	X	6.90	13.81	20.3
	Y	8.74	17.49	10.4
	Z	8.21	16.42	20.9
	R	2.65	5.31	9.1
	S	2.74	5.49	18.3
	A	3.04	6.09	9.3
	K	3.66	7.33	17.3
	T and V	3.66	7.33	16.2

Sulphate of ammonia: None, 0.6 cwt N per acre
 Muriate of potash: None, 1.0 cwt K₂O per acre

Basal Manuring: 3.75 cwt. Superphosphate per acre

Cultivations, etc: Ploughed: during Jan. Cultivated: Mar.16
 Harrowed: Mar. 25. Ridged: Apr.23-24. Superphosphate
 drilled, sulphate of ammonia and muriate of potash applied
 Apr.26. Organics applied: Apr.29-30. Potatoes planted
 and covered in: Apr. 30-May 1. Rolled down ridges: May 6
 Harrowed: May 25. Grubbed: June 15. Weeded and earthed
 up: July 2-3. Sprayed with "Perinox": Aug.10. Sprayed
 to kill off haulm: Sept.15. Lifted: Oct.1-4. Variety:
 Majestic (Scotch A). Previous crop: Barley.

Standard errors per plot:

Total tubers, per whole plot, 0.661 tons per acre or 5.28%
 (24 d.f.)
 per sub-plot, 0.983 tons per acre or 7.84%
 (29 d.f.)

48/Ca/2.1

POTATOES

Effects of intensive intertillage, of earthing up, of mulching with straw, of applying artificials before and after ridging, and of spraying haulm before lifting.

RF - Sawyers II, 1948

System of replication: 4 randomized blocks of 10 plots each.

Area of each plot (after rejecting edge rows): 0.0098 acres.

Treatments:

Cultivations: All four combinations of intensive and little intertillage between rows, with and without earthing up; also intensive cultivation until the crop was well through the ground, then mulching with 3 tons chaffed straw per acre between rows.

Application of fertilizers: Broadcast before ridging, applied in the bouts.

Spraying: Two of the four blocks were sprayed with sulphuric acid to kill off haulm before lifting.

Cultivations, etc.:

Ploughed: during Jan. Cultivated: Mar 16. Harrowed: Mar 25.
Artificials applied before ridging: Apr 10. Ridged, artificials applied, potatoes planted and covered in: Apr 12. Rolled ridges: Apr 21. Chain-harrowed: May 18. Grubbed "intensive intertillage" plots: June 14. Grubbed, all plots: June 19. Earthed up, appropriate plots: June 26. Chaffed straw applied: July 8. Sprayed to kill off haulm: Sept 14. Lifted: Oct 5, 6. Variety: Majestic (Scotch A).
Previous crop: Barley.

Standard errors per plot:

Total tubers,	1.35 tons per acre or 10.2%
Percentage ware,	0.780
Percentage greened,	5.02

All standard errors estimated from 22 d.f.

	Little Intertillage		Intensive Intertillage		Mean	Effect of Spraying
	Not earthen up	Earthen up	Not earthen up	Earthen up		
	Total tubers, tons per acre (± 0.673)					
Artificials applied before ridging after ridging	12.12 13.69	12.37 13.19	12.15 13.72	11.72 13.21	(± 0.301) 12.63 13.76	(± 0.602) -0.38 -0.06
Mean (± 0.476)	12.91	13.03	12.93	12.46	13.19	
Effect of Spraying (± 0.952)	-0.26	0.60	-0.64	-0.57	-0.23	
	Percentage Warts (± 0.276)					
Artificials applied before ridging after ridging	97.92 97.70	97.75 98.50	97.42 97.92	98.90 98.80	98.08 98.82	(± 0.349) 0.27 0.54
Mean (± 0.276)	97.81	98.12	97.68	98.85	98.45	
Effect of Spraying (± 0.552)	0.32	0.40	-0.10	0.15	1.25	

48/Ca/2.2

48/Ca/3.1

POTATOES

The effects of four times of planting, of dung, sulphate of ammonia, superphosphate and muriate of potash.

RP - Sawyers II, 1948

System of replication: 4 randomized blocks of 16 plots each, certain high order interactions being confounded with block differences.

Area of each plot: 0.0133 acre

Treatments:

Time of planting: 10th April, 24th April, 8th May, 22nd May.

Dung: None, 15 tons F.Y.M. per acre

Sulphate of ammonia: None, 0.6 cwt. N per acre

Superphosphate: None, 0.6 cwt. P_2O_5 per acre

Muriate of Potash: None, 1.0 cwt. K_2O per acre.

Cultivations: Whole experiment; Ploughed: during Jan. Cultivated: Mar 16
Tooth harrowed: Mar 25. Sprayed with Perenox: Aug 10. Sprayed with
20% B.O .V. to kill off haulm: Sept 15. Lifted: Oct 4-5.

1st planting; Bouted: Apr 8. Artificially applied: Apr 9.

Dung applied, potatoes planted and covered in: Apr 10. Rolled ridges:
Apr 12. Chain harrowed: May 18. Grubbed: June 14 and again June 19.
Earthed up: June 21.

2nd planting; Ridged, dung and artificially applied, potatoes
planted and covered in: Apr 24. Rolled ridges: May 6. Chain harrowed:
May 18. Grubbed: June 14 and again June 19. Earthed up: June 26.

3rd planting; Thistles cut, ridged, dung and artificially
applied, potatoes planted and covered in: May 7. Rolled ridges: May 8.
Chain harrowed: June 5. Weeded: July 5. Earthed up: July 7.

4th planting; Thistles cut: May 7. Dung and artificially
applied, potatoes planted and covered in: May 21. Rolled ridges: May 22.
Chain harrowed: June 5. Weeded: July 5. Earthed up: July 29.

Variety: Majestic (Scotch A), Previous crop: Barley.

Standard error per plot: Total tubers, ± 1.98 tons per acre or 21.0% (34 d.f.)

Total tubers: tons per acre

48/Ca/3.2

	Time of Planting				Mean
	10th April	24th April	8th May	22nd May	
Mean (± 0.496)	12.15	9.22	9.42	6.91	9.42
No Dung (± 0.701)	10.75	8.35	8.61	7.10	8.71
Dung	13.55	10.08	10.23	6.71	10.14
Response to Dung (± 0.992)	2.80	1.73	1.62	-0.39	1.43 ⁽¹⁾
No Nitrogen (± 0.701)	11.96	9.53	8.92	6.54	9.24
Nitrogen	12.34	8.91	9.92	7.28	9.61
Response to Nitrogen (± 0.992)	0.38	-0.62	1.00	0.74	0.37 ⁽¹⁾
No Superphosphate (± 0.701)	10.98	8.63	9.46	7.36	9.11
Superphosphate	13.32	9.81	9.38	6.46	9.74
Response to Superphosphate (± 0.992)	2.34	1.18	-0.03	-0.90	0.63 ⁽¹⁾
No Potash (± 0.701)	11.22	9.19	9.17	6.83	9.10
Potash	13.08	9.25	9.68	6.99	9.75
Response to Potash (± 0.992)	1.86	0.06	0.51	0.16	0.65 ⁽¹⁾
Standard Error (1) ± 0.496					

48/08/3.3

Responses to Treatments ± 0.701

Response to	Dung		Sulphate of Ammonia		Super-Phosphate		Muriate of Potash	
	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Dung	-	-	1.14	1.74	1.69	1.19	1.97	0.91
Sulphate of ammonia	0.07	0.67	-	-	0.41	0.33	0.60	0.14
Super-phosphate	0.89	0.39	0.68	0.60	-	-	0.58	0.70
Muriate of potash	1.18	0.12	0.88	0.42	0.59	0.71	-	-

48/Ce/1.1

SUGAR BEET

The effects of two methods of broadcasting and of two other methods of placement of two levels of a compound fertilizer.

RS - Long Hoos. 1948

System of replication: 3 randomized blocks of 11 plots each

Area of each plot (after rejecting edge rows): 0.00906 acre

Treatments:

Levels of fertilizer: None, 5, 10 cwt per acre of National Compound No 2 (9% N, 7% P₂O₅, 4½% K₂O)

Methods of placement: Broadcast before all cultivations, broadcast on seedbed and harrowed in at once, placed in band 2" below and 1" to side of seed, placed in band 2" below and 3" to side of seed.

Basal Manuring: None

Cultivations etc: Ploughed: Sept. 30 - Oct 2. Springtine harrowed then rolled and springtined: Mar 10. Ground chalk applied 2 tons per acre: Mar 13. Fertilizer broadcast before cultivation on appropriate plots: Mar 19. Cultivated: Mar 20. Rolled and springtine harrowed: Mar 22. Harrowed and ring rolled, fertilizer broadcast on seedbed, seed drilled, and fertilizers placed: Mar 30. Dusted against flea beetle: May 7, 14 and 15. Hoed: May 27 - 28. Singled: June 4 - 5. Hoed: June 12 - 15, 16, 23 - 24, July 15 - 16 and 19 - 20. Lifted: Dec 8. Variety: Klein E. Previous crop: Barley.

Standard errors per plot:

Clean beet:	±0.950 tons per acre or 7.33%
Sugar percentage:	±0.352
Total sugar:	±3.27 cwt per acre or 7.4%
Tops:	±1.70 tons per acre or 12.2%
Plant number:	±1.02 thousand per acre or 4.4%

All standard errors estimated from 22 d.f.

48/Ce/1.2

Fertilizer (cwt/acre)	Broad- cast before cultivations	Broad- cast on seed bed	Band 2" below and 1" to side	Band 2" below and 3" to side	Mean
Clean Beet: tons per acre (± 0.549)					
None					11.49 (± 0.317)
5	13.49	13.77	13.24	13.19	13.42 (± 0.274)
10	13.88	13.37	13.98	13.23	13.61
Mean (± 0.388)	13.68	13.57	13.61	13.21	12.97
Sugar Percentage: (± 0.203)					
None					17.14 (± 0.117)
5	17.09	17.11	17.11	16.98	17.07 (± 0.102)
10	16.77	16.79	16.79	17.26	16.90
Mean (± 0.144)	16.93	16.95	16.95	17.12	17.03
Total Sugar: cwt per acre (± 1.89)					
None					39.4 (± 1.09)
5	46.1	47.1	45.3	44.8	45.8 (± 0.95)
10	46.6	44.9	47.0	45.7	46.0
Mean (± 1.34)	46.3	46.0	46.2	45.2	44.1
Tops: tons per acre (± 0.983)					
None					10.44 (± 0.567)
5	13.64	12.36	13.52	15.01	13.63 (± 0.491)
10	16.42	15.96	18.89	16.55	16.95
Mean (± 0.695)	15.03	14.16	16.21	15.78	13.97
Plant Number: thous. per acre (± 0.589)					
None					23.2 (± 0.340)
5	22.7	22.7	23.1	23.3	23.0 (± 0.295)
10	23.2	23.1	23.4	22.6	23.1
Mean (± 0.417)	23.0	22.9	23.3	22.9	23.1

48/Cf/1.1

LINSEED

The effects of time of sowing, seed rate, sulphate of ammonia, superphosphate and muriate of potash.

R/JL - Bones Close 1948

System of replication: 4 randomised blocks of 8 plots each; certain high order interactions confounded with block differences.

Area of each plot: 0.0212 acre.

Treatments:

Time of sowing: 24th March (Early), 20th April (Late).
Seed rate: 60 and 90 lbs per acre.
Sulphate of ammonia: None, 0.45 cwt N per acre.
Superphosphate: None, 0.60 cwt P_2O_5 per acre.
Muriate of Potash: None, 0.45 cwt K_2O per acre.

Basal Manuring: None.

Cultivations etc: Ploughed: Nov 10-18. Springtime harrowed: Mar 3-4. Rolled and springtime harrowed: Mar 12. Harrowed: Mar 23. "Early" seed drilled, and artificials applied to "Early" plots: Mar 24. Harrowed in: Mar 25. Ring rolled: Mar 27. "Late" seed drilled, and artificials applied to "Late" plots: Apr 20. Harrowed and light ring rolled: Apr 21. Dusted against flea beetle: May 18. Thistles hand hoed: May 25. Agroxone used against weeds: May 27. Hoed: June 4-7 and 8-9. Harvested: Aug 28. Variety: Royal
Previous crop: Barley.

Note: The late sown crop was practically a failure, and was discarded.

Standard error per plot: ± 0.979 per acre or 12.6% (3 d.f.)

Grain: cwt per acre

Mean yield 7.78

Responses to treatments (± 0.693 . Means ± 0.490)

Responses to:	Sulphate of Ammonia		Superphosphate		Muriate of Potash		Seed Rate lb per acre
	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.	
Sulphate of Ammonia	1.61	-	1.23	1.99	-	-	60
Superphosphate	-1.08	-1.46	-	-	-0.75	-1.41	1.25
Muriate of Potash	1.90	-	2.23	1.57	-	-	1.97
Seed Rate 90-60	1.28	0.92	0.61	1.95	2.56	0.00	-1.75
							3.18
							0.62

Note:

Owing to the discarding of the late sown crop, the interaction of Sulphate of Ammonia and Muriate of Potash was confounded with block differences.

48/cf/1.2

48/Cf/2

LINSEED

The effects of rates and methods of application of a compound fertilizer.

R/JL - Bones Close 1948

System of replication: 4 randomized blocks of 6 plots each.

Area of each plot: 0.0191 acre.

Treatments:

None

$4\frac{1}{2}$ and 9 cwt compound fertilizer per acre, broadcast.

$2\frac{1}{4}$ and $4\frac{1}{2}$ cwt compound fertilizer per acre, drilled.

The compound fertilizer was made up of:

	lb
Sulphate of Ammonia	32
Superphosphate	57
Muriate of Potash	<u>11</u>
Total	100

= 6.7% N, 9.0% P₂O₅, 6.7% K₂O.

Basal Manuring: None.

Cultivations etc: Ploughed: Nov 10-18. Springtime harrowed: Mar 3-4.
 Rolled and springtime harrowed: Mar 12. Tooth-harrowed: Mar 23. Seed
 drilled, and compound drilled and broadcast: Mar 24. Harrowed in:
 Mar 25. Ring rolled Mar 27. Dusted against flea beetle: May 18.
 Thistles hand hoed: May 25. Agroxone applied against weeds: May 27.
 Hoed: June 4-7 and again 8-9. Harvested: Aug 28. Variety: Royal.
 Previous crop: Barley.

Standard error per plot: 1.42 cwt per acre or 14.4% (16 d.f.)

Grain: cwt per acre

Level	Compound			Mean
	None	Broadcast	Drilled	
1		11.2	9.6	10.4
2		8.2	7.9	8.1
Mean	11.2	9.7	8.8	9.9
Difference		-3.0	-1.7	-2.3

(±0.71) (±0.50)

(±1.01) (±0.71)

48/Cf/3.1

LINSEED

The effects of time of sowing, seed rate, sulphate of ammonia, superphosphate and muriate of potash.

Broad Mead I Woburn 1948

System of replication: 4 randomized blocks of 8 plots each; certain high order interactions confounded with block differences.

Area of each plot: 0.0212 acre.

Treatments:

Time of sowing: 22nd March (Early), 16th April (Late)

Seed rate: 60 and 90 lbs per acre.

Sulphate of ammonia: None, 0.45 cwt N per acre.

Superphosphate: None, 0.60 cwt P_2O_5 per acre.

Muriate of potash: None, 0.45 cwt K_2O per acre

Basal Manuring: None.

Cultivations etc: Ploughed: Feb 4 and Mar 1. **Rolled:** Mar 2. Harrowed three times: between Mar 12-18. "Early" seed sown and artificials applied to "Early" plots: Mar 22. Harrowed and rolled: Mar 23. "Late" seed sown and artificials applied to "Late" plots: Apr 16. Harrowed and rolled: Apr 16. Dusted against flea beetle: May 5-8, and again May 14. Weeds pulled: June 8. Early plots harvested: Aug 19. Late plots harvested: Sept 9.

Standard Error per plot: ± 1.296 cwt per acre or 14.7% (13 d.f.)

Grain: cwt per acre

Mean yield: 6.79

Responses to treatments: (± 0.648 . Means ± 0.458)

Response to:	Sulphate of ammonia		Super phosphate		Muriate of potash		Sowing		Seed Rate lbs per acre
	Absent	Present	Absent	Present	Absent	Present	Early	Late	
Mean									
Sulphate of ammonia	0.05	-	0.47	-0.37	0.40	-0.30	-1.06	1.16	0.84
Superphosphate	-0.16	-0.58	-	-	0.77	-1.09	-0.32	0.00	-0.84
Muriate of Potash	0.29	-0.06	1.22	-0.64	-	-	0.10	0.48	-0.26
Sowing Late-Early	-1.00	0.11	-1.16	-0.84	-1.19	-0.81	-	-	-0.26
Seed Rate 90-60	0.26	-0.53	0.94	-0.42	-0.29	0.81	1.00	-0.48	-

48/Cf/3.2

48/Cf/4

LINSEED

The effects of rates and methods of application of a compound fertilizer.

W/JL - Broad - Mead I Woburn 1948

System of replication: 4 randomized blocks of 6 plots each.

Area of each plot: 0.0212 acre.

Treatments:

None.

$4\frac{1}{2}$ and 9 cwt compound fertilizer per acre, broadcast

$2\frac{1}{4}$ and $4\frac{1}{2}$ cwt compound fertilizer per acre, drilled.

The compound fertilizer was made up of:

	lb
Sulphate of Ammonia	32
Superphosphate	57
Muriate of Potash	<u>11</u>

Total 100

= 6.7% N.

9.0% P_2O_5

6.7% K_2O

Basal Manuring: None.

Cultivations etc: Ploughed: Feb 4 - Mar 1. Rolled: Mar.2. Harrowed three times: Mar 12-18. Seed sown and fertilizer applied to all plots: Mar 22. Harrowed and ring rolled: Mar 23. Dusted against flea beetle: May 5-8. Weeds pulled: June 8. Harvested: Aug 19. Variety: Royal. Previous crop: Ley.

Standard error per plot: 1.26 cwt. per acre or 13.0% (16 d.f.)

Grain: cwt per acre

	Compound			Mean
	None	Broadcast	Drilled	
Level 1		9.9	10.5	10.2
Level 2		8.9	8.6	8.7
Mean	10.2	9.4	9.5	9.7
Difference		-1.0	-1.9	-1.5
(±0.45)		(±0.63)		(±0.45)
(±0.89)				(±0.63)

48/Cg/1.1

PEAS

The effects of two methods of broadcasting and of two other methods of placement of two levels of a compound fertilizer.

R/PE - Long Hoos, 1948

System of replication: 3 randomized blocks of 12 plots each

Area of each plot: 0.0150 acre

Treatments:

Levels of fertilizer: None, 3, 6 cwt. per acre of a special mixture (10% P₂O₅, 20% K₂O)

Methods of placement: Broadcast before all cultivations, broadcast on seedbed and harrowed in at once, placed in band 2" below and 1" to side of seed, placed in band 2" below and 3" to side of seed.

Basal manuring: None

Cultivations, etc.: Ploughed: Sept 30 - Oct 2. Springtine harrowed then rolled and springtined: Mar 10. Ground chalk, 2 tons per acre, applied: Mar 13. Broadcast fertilizers, before cultivations on appropriate plots: Mar 19. Cultivated: Mar 20. Rolled and springtine harrowed: Mar 22. Harrowed, and ring rolled: Mar 30. Fertilizers broadcast in seedbed, seed drilled and fertilizers placed: Mar 30. Plant almost failed so decision made to resow, thistles cut, and seed redrilled: Apr 20. Harrowed and light ring rolled: Apr 21. Weeds pulled: June 7-9. Hoed: June 12.

Standard errors per plot:

Yield, 1.61 cwt. per acre or 9.2% (25 d.f.)

Plant number, 1.47 tens of thousands per acre or 10.5% (25 d.f.)

Fertilizer (cwt/acre)	48/Cg/1.2				Mean
	Broadcast before cultivations	Broad- cast on seed bed	Band 2" below and 1" to side	Band 2" below and 3" to side	
Yield of threshed peas: cwt. per acre (± 0.927)					(± 0.464)
None					16.3
3	17.8	17.2	18.2	18.6	17.9
6	17.3	17.2	19.2	18.1	17.9
Mean (± 0.656)	17.5	17.2	18.7	18.4	17.4
Plant number: tens of thousands per acre (± 0.851)					(± 0.425)
None					13.4
3	14.0	13.2	13.5	15.5	14.1
6	14.6	13.7	15.3	14.2	14.5
Mean (± 0.601)	14.3	13.5	14.4	14.9	14.0

48/Da/1

MEADOW HAY

17th Season

Direct and residual effects of mixed artificials and compost.

Lady Manners School, Bakewell 1948

System of replication: 4 randomized blocks of 9 plots each

Area of each plot: 0.00492 acre

Treatments: 3 x 3 factorial design.

No manure, mixed artificials (2 cwt nitrate of soda, 3 cwt superphosphate and 1 cwt muriate of potash per acre), 8 tons per acre compost applied in alternate years from 1932 to 1948 or from 1933 to 1947.

Basal Manuring: None

Soil: Limestone

Cultivations: Compost and artificials applied: Mar 22. Harvested: July 1.

Standard error per plot: ± 7.02 cwt. per acre or 16.0% (24 d.f.)

Hay: cwt. per acre

Alternate years 1932 - 1948	Alternate years 1933 - 1947			Mean	Increase
	No Manure	NPK	Compost		
		(± 3.51)		(± 2.03)	(± 2.87)
No manure	29.9	27.7	43.5	33.7	
NPK	41.7	44.0	49.0	44.9	11.2
Compost	51.2	54.4	53.1	52.9	19.2
Mean (± 2.03)	41.0	42.0	48.5	43.8	
Increase (± 2.87)		1.0	7.5		

48/Da/2

MEADOW HAY

18th Season

Effects of nitrate of soda, superphosphate and muriate of potash

Lady Manners School, Bakewell 1948

System of replication: 3 randomized blocks of 8 plots each.

Area of each plot: 0.00723 acre

Treatments: 2 x 2 x 2 factorial design

Nitrate of soda: None, 2 cwt per acre

Superphosphate: None, 3 cwt per acre

Muriate of Potash: None, 1 cwt per acre

Basal Manuring: None

Soil: Limestone.

Artificials applied: Mar 22. Harvested: June 30

Standard error per plot: ± 7.08 cwt per acre of 21.2% (14 d.f.)

Hay: cwt per acre

Mean yield 33.3

Responses to treatments (± 4.09 Means ± 2.89)

	Mean	Nitrate of soda		Superphosphate		Muriate of potash	
		Absent	Present	Absent	Present	Absent	Present
Nitrate of Soda	7.2	-	-	7.4	7.0	2.5	11.9
Super-phosphate	3.7	3.9	3.5			-1.2	8.6
Muriate of potash	9.1	4.4	13.8	4.2	14.0	-	-

48/z1/1

CHEMICAL ANALYSES OF MANURES USED IN THE THREE, FOUR
AND SIX COURSE ROTATIONS 1948

Three Course Rotation

Manures	% Organic matter	% N	% P ₂ O ₅	% K ₂ O
Chaffed Straw	82.0	0.41	0.11	0.66
Adco	21.9	0.43	0.22	0.18
Sulphate of Ammonia		21.0		
Nitrate of Soda		15.5		
Superphosphate			17.7 (total)	
Muriate of Potash				62.0

Four Course Rotation

Manures	% Organic matter	% N	% P ₂ O ₅	% K ₂ O
Chaffed Straw	82.0	0.41	0.11	0.66
Dung	20.6	0.58	0.29	0.92
Adco	21.9	0.43	0.22	0.18
Sulphate of Ammonia		21.0		
Superphosphate			17.7	
Mineral Phosphate			33.3	
Muriate of Potash				62.0

Six Course Rotation

Manures	% Organic matter	% N	% P ₂ O ₅	% K ₂ O
Sulphate of Ammonia		21.0		
Superphosphate			17.7 (total)	62.0
Muriate of Potash				62.0