

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1948

[Full Table of Content](#)



---

## 48/BA/1 Two-course Rotation - Rothamsted

### Rothamsted Research

Rothamsted Research (1949) *48/BA/1 Two-course Rotation - Rothamsted* ; Yields Of The Field Experiments 1948, pp 11 - 15

48/Ba/1.1

## TWO COURSE ROTATION EXPERIMENT

### Cumulative Effects of Agricultural Salt

Rothamsted 1948

(For full details see 1942 Report)

#### Treatments:

All combinations of:

- (1) Agricultural salt: None,  $2\frac{1}{2}$ , 5 and  $7\frac{1}{2}$  cwt per acre applied to sugar beet or at half these rates to barley.
- (2) Muriate of potash: None, the equivalent of half the single dressing of salt (approximately 1 cwt.  $K_2O$  per acre), the equivalent of the single dressings of salt (approximately 2 cwt.  $K_2O$  per acre), applied to sugar beet at sowing.
- (3) Time of application of salt: In seed bed at sowing, before ploughing in winter.
- (4) Salt applied to sugar beet only, salt repeated at half rate on barley.

Note: treatment (3) applies to both sugar beet and barley crops; the barley receives no potash treatment. For ease of analysis  $K_2O$  of the muriate of potash dressings is made equivalent to  $Cl$  of agricultural salt dressings.

Basal dressings, applied to all plots at sowing:

Barley: 0.3 cwt. N per acre as sulphate of ammonia  
Sugar beet: 0.8 cwt. N per acre as sulphate of ammonia  
0.6 cwt.  $P_2O_5$  per acre as superphosphate.

Cultivations, etc:

Sugar Beet. Series I. Long Hoos VII

Agricultural salt applied: Oct. 21. Ploughed: Nov. 10-21. Cultivated: Mar. 16. Rolled and springtime harrowed, ring rolled: Mar. 27. Sulphate of ammonia applied: Mar. 30. Superphosphate, agricultural salt and muriate of potash applied: Apr. 12. Harrowed: Apr. 14. Seed drilled: Apr. 15. Seed harrowed in: Apr. 16. Rolled: Apr. 22. Dusted with Flea beetle dust: May 14. Plots 6, 12, 18, 24, 30, 36, 42, 48 damaged by herbicide spray from adjacent barley; May 14. Bare patches resown: late May. Hoed: May 20-21, 25. Weeds hand pulled: June 9-11. Singled: June 10-11. Hoed: June 21 - July 22. Lifted: Nov. 23-30. Variety: Klein E. Previous crop: Barley.

48/Ba/1.2

Barley. Series II. Long Hoos V.  
Agricultural salt applied: Dec 2. Ploughed: Dec 10-12.  
Springtime harrowed: Mar 1. Harrowed, seed drilled,  
sulphate of ammonia applied: Mar 8. Applied agricultural  
salt, harrowed in, ring rolled: Mar 9. Sprayed with  
"Denocate" to clear charlock, (plots 66, 65, 64, 63, 62,  
61 damaged by an overdose of spray): May 14. Hand weeded:  
June 30. Harvested: Aug. 14. Variety: Plumage Archer.  
Previous crop: Sugar Beet.

Standard errors per plot:

Sugar beet, total sugar,	4.40 cwt. per acre or	10.9%
tops,	1.13 tons per acre or	7.8%
Barley, grain,	2.20 cwt. per acre or	7.1%

All standard errors estimated from 22 d.f.



48/Ba/1.3

Series I: Sugar Beet

Salt in 1948 (cwt. per acre)	Muriate of potash K <sub>2</sub> O cwt. per acre			Salt applied		Mean
	0.0	1.0	2.0	Winter	In seed bed	
Total Sugar: cwt. per acre						
		(±2.20)		(±1.68)		(±1.27)
0	33.4	34.2	35.7	41.6	39.6	34.4
2.5	43.7	37.9	40.2	42.2	43.5	40.6
5.0	41.8	44.6	42.0	43.4	45.1	42.8
7.5	41.9	44.6	46.1			44.2
Mean	40.2	40.3 (±1.10)	41.0	42.4 (±1.02)	42.7	40.5
Sugar percentage						
0	16.24	16.34	16.57			16.38
2.5	16.49	16.46	16.78	16.57	16.58	16.57
5.0	16.75	16.90	16.42	16.58	16.80	16.69
7.5	16.59	16.62	16.84	16.78	16.60	16.69
Mean	16.51	16.58	16.65	16.64	16.66	16.58
Roots (washed): tons per acre						
0	10.26	10.44	10.74			10.48
2.5	13.27	11.50	11.97	12.56	11.93	12.25
5.0	12.49	13.20	12.74	12.69	12.93	12.81
7.5	12.63	13.42	13.71	12.92	13.59	13.25
Mean	12.16	12.14	12.29	12.73	12.81	12.20
Tops: tons per acre						
		(±0.566)		(±0.433)		(±0.327)
0	14.21	14.80	13.90			14.30
2.5	14.57	14.27	14.68	13.05	15.97	14.51
5.0	13.28	13.75	15.36	14.86	13.40	14.13
7.5	14.90	17.08	13.56	13.79	16.57	15.18
Mean	14.24	14.98 (±0.283)	14.37	13.90 (±0.261)	15.31	14.53

48/Bn/1.4

Series I: Sugar Beet (contd.)

Salt in 1948 (cwt. per acre)	Muriate of potash K <sub>2</sub> O cwt. per acre			Salt applied		Mean
	0.0	1.0	2.0	Winter	In seed bed	
	Plant No.: thousands per acre					
0	22.3	22.2	21.4			22.0
2.5	22.0	21.9	22.3	21.9	22.3	22.1
5.0	21.9	22.6	21.6	21.7	22.3	22.0
7.5	22.1	22.4	21.5	22.2	21.8	22.0
Mean	22.1	22.3	21.7	21.9	22.1	22.0
	Noxious Nitrogen: m.g. per 100 grams.					
0	30.2	31.5	31.2			31.0
2.5	28.5	30.2	28.8	28.7	29.7	29.2
5.0	29.8	28.6	28.0	29.3	28.3	28.8
7.5	31.0	29.5	30.8	31.2	29.7	30.4
Mean	29.9	30.0	29.7	29.7	29.2	29.8

48/Ba/1.5

Series II: Barley

Salt in 1947 (cwt. per acre)	Muriate of potash K <sub>2</sub> O cwt. per acre applied in 1947			Salt applied In seed bed		Salt in 1948 None Half Rate		Mean
	0.0	1.0	2.0	Winter				
	Grain: cwt. per acre							
	(±1.10)			(±0.84)				(±0.63)
0	31.3	33.6	30.6					31.8
2.5	32.6	30.8	31.2	31.6	31.5	31.2	31.9	31.5
5.0	30.2	31.1	29.8	30.8	30.0	29.1	31.6	30.4
7.5	29.6	30.0	29.6	30.5	29.0	29.1	30.3	29.7
Mean	30.9	31.4	30.3	31.0	30.2	29.8	31.3	30.9
	(±0.55)			(±0.51)				
	Straw: cwt. per acre							
0	35.9	37.0	35.3					36.1
2.5	36.8	33.2	34.7	34.8	34.9	35.6	34.1	34.9
5.0	32.7	33.4	32.0	31.9	33.5	32.1	33.3	32.7
7.5	32.4	32.2	31.5	31.4	32.7	30.7	33.4	32.0
Mean	34.4	34.0	33.4	32.7	33.7	32.8	33.6	33.9