Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1948



Full Table of Content

48/CA/3 Wheat - Wireworm 1

Rothamsted Research

Rothamsted Research (1949) 48/CA/3 Wheat - Wireworm 1; Yields Of The Field Experiments 1948, pp 65 - 65

48/Ca/3

WHEAT

Wireworm Experiment (1)

The effect of various insecticides, and their methods of application.

RW ~ Little Hoos 1948

System of replication: 3 randomized blocks of 9 plots each

Area of each plot: 0.0289 acre

Treatments:

None

D.D. injected 400 lbsper acre.

Ethylene Dibromide 41% solution injected 15 gallons per acre.

D.D.T. dust combine drilled 4 cwt per acre.

Gammexane; broadcast 2 cwt per acre, combine drilled $\frac{3}{4}$ cwt per acre, or applied as seed dressing.

Basal manuring: 23 cwt per acre sulphate of ammonia.

Cultivations etc.: Ploughed: Sept 16-27. Harrowed and rolled: Oct 7.

DD and Ethylene Dibromide injected: Oct 10. Seed drilled, and remaining treatments applied: Oct 29-30. Harrowed in: Oct 30. Harrowed: Mar 30. Ring rolled: Apr 1. Sulphate of ammonia drilled: May 5. Hand weeded: June 5, 7-8, 22-23, and various days June 28 - July 23. Harvested: Aug 17. Variety: Bersee. Previous crop: Linseed.

Standard errors per plot:

Grain, 2.62 cwt per acre or 12.6% (18 d.f.) Straw, 6.37 cwt per acre or 1335% (18 d.f.)

			Ethylen Dibromid Injected		Broad-	exans Drilled	Treated seed	Mean
	Grain: cwt. per acre							
Mean Yield (±1.51)	8.9(1)	28.3	32.1	20.7	30.6	21+.8	24.0	20.8
Increase (+1.74)		19.4	23.2	20.7	21.7	15.9	15.1	
	Straw: cwt. per acre							
Mean Yield (±3.68) Increase (±4.25)	22.8 ⁽²⁾	64.1	71.6	46.0	65.9	54.2	54.6 31.8	47.2
(24.25)		41.3	48.8	23.2	43.1	31.4	31.8	
Standard			U	1		1		

Standard errors (1) ±0.87 (2) ±2.12