


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1948

[Full Table of Content](#)



48/CA/4 Wheat - Wireworm 2

Rothamsted Research

Rothamsted Research (1949) *48/CA/4 Wheat - Wireworm 2 ; Yields Of The Field Experiments 1948*, pp 66 - 66

43/Ca/4.1

WHEAT

Wireworm Experiment (2)

The effect of treatment of seed with gammexane and of three strengths of gammexane dust.

RW - Little Hoos 1948

System of replication: 3 replicates of 3 incomplete randomized blocks of 3 plots each.

Area of each plot: 0.0289 acre

Treatments:

None.

Seed treated with gammexane dressing.

Gammexane dust, $\frac{1}{4}$, $\frac{1}{2}$ and 1 cwt per acre; combine drilled with seed (filler added where necessary to make total dressing of 1 cwt per acre)

Basal Manuring: $2\frac{3}{4}$ cwt per acre sulphate of ammonia.

Cultivations etc:

Ploughed: Sept 16-27. Harrowed and rolled: Oct 7. Seed and gammexane dust drilled: Oct 29-30. Harrowed in: Oct 30.

Harrowed: Mar 30. Ring rolled: Apr 1. Sulphate of ammonia drilled: May 5. Hand weeded: June 5, 7-8, 22-23, and various days June 23-July 23. Harvested: Aug 17.

Variety: Bersée. Previous crop: Linseed.

Standard errors per plot:

Grain: Block error, 1.61 cwt per acre or 7.15% (4d.f.)
Plot error, 2.28 cwt per acre or 10.1% (16d.f.)

Straw: Block error, 2.29 cwt per acre or 4.63% (4d.f.)
Plot error, 4.86 cwt per acre or 9.86% (16d.f.)