


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1948

[Full Table of Content](#)



48/CA/5 Wheat - Inoculation

Rothamsted Research

Rothamsted Research (1949) *48/CA/5 Wheat - Inoculation* ; Yields Of The Field Experiments 1948, pp 67 - 69

4.8/Ca/4.2

	Mean Yields				0 v S v D	Standard errors	
	Untreated (0)	Treated Seed (S)	Gammexane Dust (D)			D ₁ v D ₂ v D ₃	0, S v D ₁ , D ₂ , D ₃
			$\frac{1}{4}$ cwt/acre (D ₁)	$\frac{1}{2}$ cwt/acre (D ₂)	1 cwt/acre (D ₃)		
Grain: cwt/acre	13.6	26.3	28.4	27.7	±0.76	±1.42	±1.12
Straw: cwt/acre	32.8	56.2	60.2	59.6	±1.63	±2.65	±2.23
			27.5	60.4			
			27.1	58.0			

48/Ca/5.1

SPRING WHEAT

The effects of inoculation and wetting of seed, and of sulphate of ammonia.

RW - Long Hoos 1948

System of replication: 3 randomized blocks of 6 plots each.

Area of each plot: 0.0158 acre.

Treatments:

Seed: untreated, inoculated by wet method, wetted but not inoculated.

Sulphate of ammonia: None, 0.3 cwt. N per acre.

Note: The inoculum was a bacterial culture of French origin.

Basal manuring: 3 cwt. Superphosphate and 1 cwt. muriate of potash per acre.

Cultivations, etc.: Ploughed: Sept. 30 - Oct. 2. Rolled and springtine harrowed: Mar. 10. Sulphate of ammonia applied and basal manuring broadcast by machine: Mar. 12. Harrowed and seed drilled: Mar. 13. Harrowed in: Mar. 15. Rolled: Mar. 20. Weeds hand pulled: June 8. Harvested: Aug. 24. Variety: Bersee.

Standard errors per plot (10 d.f.): Grain, 3.35 cwt. per acre
or 17.1%
Straw, 3.00 cwt. per acre
or 7.20%

48/0a/5.2

	Seed Untreated	Seed inoculated (wet method)	Seed wetted not inoculated	Mean
Grain: cwt. per acre				
Mean (± 1.37)	19.8	18.5	20.5	19.6
Sulphate of Ammonia:				
Absent (± 1.93)	16.5	17.2	17.8	17.2
Present	23.2	19.8	23.1	22.0
Response (± 2.74)	6.7	2.6	5.3	4.8 (± 1.58)
Straw: cwt. per acre				
Mean (± 1.22)	41.3	41.3	42.3	41.6
Sulphate of Ammonia:				
Absent (± 1.73)	38.4	38.6	38.7	38.6
Present	44.2	44.1	45.8	44.7
Response (± 2.45)	5.8	5.5	7.1	6.1 (± 1.41)