


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1948

[Full Table of Content](#)



---

## 48/CF Linseed

### Rothamsted Research

Rothamsted Research (1949) *48/CF Linseed* ; Yields Of The Field Experiments 1948, pp 88 - 88

48/Cf/1.1

## LINSEED

The effects of time of sowing, seed rate, sulphate of ammonia, superphosphate and muriate of potash.

R/JL - Bones Close 1948

System of replication: 4 randomised blocks of 8 plots each; certain high order interactions confounded with block differences.

Area of each plot: 0.0212 acre.

### Treatments:

Time of sowing: 24th March (Early), 20th April (Late).  
Seed rate: 60 and 90 lbs per acre.  
Sulphate of ammonia: None, 0.45 cwt N per acre.  
Superphosphate: None, 0.60 cwt  $P_2O_5$  per acre.  
Muriate of Potash: None, 0.45 cwt  $K_2O$  per acre.

Basal Manuring: None.

Cultivations etc: Ploughed: Nov 10-18. Springtime harrowed: Mar 3-4. Rolled and springtime harrowed: Mar 12. Harrowed: Mar 23. "Early" seed drilled, and artificials applied to "Early" plots: Mar 24. Harrowed in: Mar 25. Ring rolled: Mar 27. "Late" seed drilled, and artificials applied to "Late" plots: Apr 20. Harrowed and light ring rolled: Apr 21. Dusted against flea beetle: May 18. Thistles hand hoed: May 25. Agroxone used against weeds: May 27. Hoed: June 4-7 and 8-9. Harvested: Aug 28. Variety: Royal  
Previous crop: Barley.

Note: The late sown crop was practically a failure, and was discarded.

Standard error per plot:  $\pm 0.979$  per acre or 12.6% (3 d.f.)