


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1948

[Full Table of Content](#)



48/CF/2 Linseed - Rothamsted : Rates and Application of Manures

Rothamsted Research

Rothamsted Research (1949) *48/CF/2 Linseed - Rothamsted : Rates and Application of Manures ; Yields Of The Field Experiments 1948*, pp 90 - 90

48/Cf/2

LINSEED

The effects of rates and methods of application of a compound fertilizer.

R/JL - Bones Close 1948

System of replication: 4 randomized blocks of 6 plots each.

Area of each plot: 0.0191 acre.

Treatments:

None

$4\frac{1}{2}$ and 9 cwt compound fertilizer per acre, broadcast.

$2\frac{1}{4}$ and $4\frac{1}{2}$ cwt compound fertilizer per acre, drilled.

The compound fertilizer was made up of:

	lb
Sulphate of Ammonia	32
Superphosphate	57
Muriate of Potash	<u>11</u>
Total	100

= 6.7% N, 9.0% P₂O₅, 6.7% K₂O.

Basal Manuring: None.

Cultivations etc: Ploughed: Nov 10-18. Springtime harrowed: Mar 3-4.
 Rolled and springtime harrowed: Mar 12. Tooth-harrowed: Mar 23. Seed
 drilled, and compound drilled and broadcast: Mar 24. Harrowed in:
 Mar 25. Ring rolled Mar 27. Dusted against flea beetle: May 18.
 Thistles hand hoed: May 25. Agroxone applied against weeds: May 27.
 Hoed: June 4-7 and again 8-9. Harvested: Aug 28. Variety: Royal.
 Previous crop: Barley.

Standard error per plot: 1.42 cwt per acre or 14.4% (16 d.f.)

Grain: cwt per acre

	None	Compound		Mean
		Broadcast	Drilled	
Level 1		11.2	9.6	10.4
		(±0.71)		(±0.50)
Level 2		8.2	7.9	8.1
Mean (±0.50)	11.2	9.7	8.8	9.9
Difference (±1.01)		-3.0	-1.7	-2.3 (±0.71)