

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1948

[Full Table of Content](#)



48/CF/4 Linseed - Woburn : Rates and Application of Manures

Rothamsted Research

Rothamsted Research (1949) *48/CF/4 Linseed - Woburn : Rates and Application of Manures ; Yields Of The Field Experiments 1948*, pp 93 - 93

48/Cf/4

LINSEED

The effects of rates and methods of application of a compound fertilizer.

W/JL - Broad - Mead I Woburn 1948

System of replication: 4 randomized blocks of 6 plots each.

Area of each plot: 0.0212 acre.

Treatments:

None.

$4\frac{1}{2}$ and 9 cwt compound fertilizer per acre, broadcast

$2\frac{1}{4}$ and $4\frac{1}{2}$ cwt compound fertilizer per acre, drilled.

The compound fertilizer was made up of:

	lb
Sulphate of Ammonia	32
Superphosphate	57
Muriate of Potash	<u>11</u>

Total 100

= 6.7% N.

9.0% P_2O_5

6.7% K_2O

Basal Manuring: None.

Cultivations etc: Ploughed: Feb 4 - Mar 1. Rolled: Mar.2. Harrowed three times: Mar 12-18. Seed sown and fertilizer applied to all plots: Mar 22. Harrowed and ring rolled: Mar 23. Dusted against flea beetle: May 5-8. Weeds pulled: June 8. Harvested: Aug 19. Variety: Royal. Previous crop: Ley.

Standard error per plot: 1.26 cwt. per acre or 13.0% (16 d.f.)

Grain: cwt per acre

	Compound			Mean
	None	Broadcast	Drilled	
Level 1		9.9	10.5	10.2
		(±0.63)		(±0.45)
Level 2		8.9	8.6	8.7
Mean	10.2	9.4	9.5	9.7
(±0.45)				
Difference		-1.0	-1.9	-1.5 (±0.63)
(±0.89)				