Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1950



Full Table of Content

50/CA/3 Wheat - Wireworm 1 - Rothamsted

Rothamsted Research

Rothamsted Research (1951) 50/CA/3 Wheat - Wireworm 1 - Rothamsted; Yields Of The Field Experiments 1950, pp 59 - 59 - DOI: https://doi.org/10.23637/ERADOC-1-185

50/ca/3

Wireworm Experiment (1)

Residual effects of insecticides - Little Hoose 1950.

System of replication: 3 randomized blocks of 9 plots each.

Area of each plot: 0.0269 acre.

Treatments - applied 1948.

None

D.D. injected 400 lb per acre.

Ethylene Dibromide 41% solution, injected 15 gallons per acre.

D.D.T. dust combine drilled 2 cwt per acre.

Gammexane: broadcast, 2 cwt per acre; combine drilled, 3 cwt per acre; applied as seed dressing.

Basal Manuring: $1\frac{1}{2}$ cwt superphosphate per acre combine drilled; $2\frac{1}{2}$ cwt sulphate of ammonia per acre as top dressing.

Cultivations, etc.: Ploughed: Oct 13-17. Springtined, seed and superphosphate drilled, harrowed in: Oct 31 - Nov 1. Harrowed: Mar 31. Rolled: Apr 3. Sulphate of ammonia applied: May 3. Sprayed with D.N.O.C.: May 11. Harvested: Aug 10. Variety: Squareheads Master 13/4. Previous crop: Wheat.

Standard errors per plot:

Grain: 2.16 cwt per acre or 13.6% (18 d.f.) Straw: 3.26 cwt per acre or 12.9% (18 d.f.)

Summary of Results

	Un- treated	D.D In-	Ethylene Dibromide Injected	D.D.T. Dust	Groad-		Dusted seed	Mean
	-		Grain: c	wt per a	cre	a a		
Mean	15.5(1)	16.9	18.8	18.0	16.1	15.0	11.7	15.9
(±1.25) Increase (±1.44)		1.4	Grain: c 18.8	2.5	0.6	-0.5	-3.8	
NATIONAL DISCONSISSION OF THE PROPERTY OF THE	Straw: cwt per acre							
Mean (±1.88) Increase (±2.17)	25.0(2)	27.0	27.8	27.1	25.8	23.0	20.9	29.2
		2.0	2.8	.2.1	0.8	-2.0	-4.1	

Standard errors (1) ±0.72

Note: Wireworm counts were made on all plots and are available.