


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1950

[Full Table of Content](#)



50/CA/5 Wheat - Late Application of Nitrogen - Rothamsted

Rothamsted Research

Rothamsted Research (1951) *50/CA/5 Wheat - Late Application of Nitrogen - Rothamsted* ; Yields Of The Field Experiments 1950, pp 62 - 62 - DOI: <https://doi.org/10.23637/ERADOC-1-185>

50/Ca/5

WHEAT

Late application of nitrogen - West Barnfield I 1950.

System of replication: 4 randomized blocks of 4 plots each.

Area of each plot: 0.0187 acre.

Treatments:

Nitrochalk: None, none but plots walked over as for broadcasting,
 $1\frac{1}{2}$, 3 cwt per acre applied as a top dressing.

Basal manuring: $1\frac{3}{4}$ cwt sulphate of ammonia and 1 cwt superphosphate per acre.

Cultivations, etc.: Ploughed: Sept 45-19. Rolled and springtine harrowed: Oct 14. Harrowed: Oct 15. Seed and superphosphate drilled and harrowed in: Oct 17. Harrowed: Mar 28. Rolled: Mar 31. Sulphate of ammonia drilled: May 3. Sprayed with low volume 2, 4-D: May 18. Nitrochalk applied by hand: June 28. Harvested: Aug 14. Variety: Squareheads Master $1\frac{3}{4}$. Previous crop: Wheat.

Standard errors per plot.

Grain: 2.59 cwt per acre or 10.0% (9 d.f.)

Straw: 8.25 cwt per acre or 16.3% (9 d.f.)

Summary of Results

	Nitrochalk: cwt per acre as top dressing				Mean
	None	None plots walked over	$1\frac{1}{2}$	3	
	cwt per acre				
Grain (± 1.30)	24.5	27.6	25.9	25.5	25.9
Straw (± 4.12)	49.0	51.7	51.3	50.6	50.7

Note

Analytical results showing increases in crude protein due to late nitrogen are given on page 116 of the Station's Annual Report for 1950.