


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1950

[Full Table of Content](#)



50/CB/1 Barley - Late Application of Nitrogen - Rothamsted

Rothamsted Research

Rothamsted Research (1951) *50/CB/1 Barley - Late Application of Nitrogen - Rothamsted* ; Yields Of The Field Experiments 1950, pp 63 - 63 - DOI: <https://doi.org/10.23637/ERADOC-1-185>

BARLEY

Late application of nitrogen - Long Hoos V 1950.

System of replication: 4 randomized blocks of 3 plots each.

Area of each plot: 0.0192 acre.

Treatments:

Nitrochalk: None, $1\frac{1}{2}$, 3 cwt per acre applied as a top dressing.

Basal manuring: $1\frac{1}{2}$ cwt sulphate of ammonia per acre.

Cultivations, etc.: Ploughed: Sept 27-Oct 3. Springtine harrowed: Mar 8 and 11. Harrowed: Mar 13. Sulphate of ammonia drilled: Mar 14. Seed drilled, harrowed in and ring rolled: Mar 16. Sprayed with high volume 2, 4-D: May 12. Nitrochalk applied by hand: June 27. Harvested: Aug 8. Variety: Plumage Archer. Previous crop: Beans.

Standard errors per plot:

Grain: 1.51 cwt per acre or 6.6% (6 d.f.)

Straw: 2.16 cwt per acre or 7.4% (6 d.f.).

Summary of Results

	Nitrochalk: cwt per acre as top dressing			Mean
	None	$1\frac{1}{2}$	3	
	cwt per acre			
Grain (± 0.76)	22.7	22.7	23.1	22.9
Straw (± 1.08)	28.2	29.6	29.5	29.1

Note

Analytical results showing increases in crude protein due to late nitrogen are given on page 116 of the Station's Annual Report for 1950.