


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1950

[Full Table of Content](#)



50/CC/1 Spring Oats - Late Application of Nitrogen - Rothamsted

Rothamsted Research

Rothamsted Research (1951) *50/CC/1 Spring Oats - Late Application of Nitrogen - Rothamsted* ;
Yields Of The Field Experiments 1950, pp 64 - 64 - DOI: <https://doi.org/10.23637/ERADOC-1-185>

50/Cc/1

SPRING OATS

Late application of nitrogen - Great Harpenden II 1950.

System of replication: 4 randomized blocks of 3 plots each.

Area of each plot: 0.0192 acre.

Treatments:

Nitrochalk: None, $1\frac{1}{2}$, 3 cwt per acre as a top-dressing.

Basal Manuring: 2 cwt sulphate of ammonia, and $1\frac{1}{4}$ cwt superphosphate per acre.

Cultivations, etc.: Ploughed: Sept 12-30, and again Dec 6-8.

Springtine harrowed both ways: Mar 14-15. Sulphate of ammonia drilled: Mar 18. Seed and superphosphate drilled and harrowed in: Mar 21-22. Ring rolled: Mar 25. Nitrochalk applied by hand: June 27. Harvested: Aug 10. Variety: Sun II. Previous crop: Wheat.

Standard errors per plot:

Grain: 2.16 cwt per acre or 7.2% (6 d.f.)

Straw: 2.06 cwt per acre or 3.8% (6 d.f.)

Summary of Results

	Nitrochalk: cwt per acre as top dressing			Mean
	None	$1\frac{1}{2}$	3	
	cwt per acre			
Grain (± 1.08)	31.4	30.0	28.8	30.1
Straw (± 1.03)	53.9	54.1	53.0	53.7

Note

Analytical results showing increases in crude protein due to late nitrogen are given on page 116 of the Station's Annual Report for 1950.