


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1950

[Full Table of Content](#)



### 50/CG/1 Lucerne - Fertilizer Placement - Rothamsted

#### Rothamsted Research

Rothamsted Research (1951) *50/CG/1 Lucerne - Fertilizer Placement - Rothamsted* ; Yields Of The Field Experiments 1950, pp 81 - 82 - DOI: <https://doi.org/10.23637/ERADOC-1-185>

50/Cg/1.1

LUCERNE

Fertilizer placement - Long Hoos IV 1950

System of replication: 3 randomized blocks of 12 plots each.

Area of each plot: 0.00798 acre. Area harvested: 0.00644 acre.

Treatments: No fertilizer (duplicate plots); no fertilizer but plots drilled over (duplicate plots); and all combinations of:

Granular PK compound fertilizer (10% P<sub>2</sub>O<sub>5</sub>, 20% K<sub>2</sub>O): 2.5, 5.0 cwt per acre.

Method of placement: Broadcast during preparation of seed bed, drilled in bands 3" below soil surface and 2" to side of seed; half broadcast during preparation of seed bed and half to be broadcast on surface next year; half drilled beside seed, and half to be drilled beside rows of plants next year.

It will be noted that for 1950 there are duplicate plots per block with 2.5 cwt per acre fertilizer broadcast or drilled.

Basal dressing: 10 cwt ground chalk per acre

Cultivations, etc.: Springtined: Mar 8 and again Mar 11. Ground chalk applied: Mar 10. Fertilizers broadcast: Mar 23. Harrowed, seed and fertilizers drilled: Mar 27. Rolled: Mar 28. Ring rolled: Mar 29. Dusted with B.H.C.: May 4. Weeded: June 21-23. First cut: Aug 9. Second cut: Oct 12. Variety: Du Puits. Previous crop: Wheat.

Standard errors per plot:

Dry Matter, 1st cut: 2.43 cwt per acre or 10.5% (26 d.f.)  
2nd cut: 1.13 cwt per acre or 7.3% (26 d.f.)

50/Cg/1.2

Summary of Results

Lucerne, Dry Matter: cwt per acre

Method of placement	Compound Fertilizer: cwt per acre				Mean
	None	1.25	2.5	5.0	
1st cut					
	(a)	(b)	(a)	(b)	(c)
Broadcast	22.5 <sup>+</sup>	27.3	25.0	25.8	25.8*
Drilled	21.4 <sup>+</sup>	22.8	20.1	23.4	21.6*
Mean	22.0 <sup>(c)</sup>	25.0 <sup>(a)</sup>	22.5 <sup>(c)</sup>	24.6 <sup>(a)</sup>	23.1
Difference	-1.1 <sup>(b)</sup>	-4.5 <sup>(d)</sup>	-4.9 <sup>(b)</sup>	-2.4 <sup>(d)</sup>	-4.2 <sup>(a)*</sup>
2nd cut					
	(a)	(b)	(a)	(b)	(c)
Broadcast	15.5 <sup>+</sup>	15.2	15.7	16.5	15.8*
Drilled	14.9 <sup>+</sup>	15.6	15.0	16.2	15.5*
Mean	15.2 <sup>(c)</sup>	15.4 <sup>(a)</sup>	15.4 <sup>(c)</sup>	16.3 <sup>(a)</sup>	15.5
Difference	-0.6 <sup>(b)</sup>	0.4 <sup>(d)</sup>	-0.7 <sup>(b)</sup>	-0.3 <sup>(d)</sup>	-0.3 <sup>(a)*</sup>

Standard errors	1st cut	2nd cut
(a)	0.99	0.46
(b)	1.41	0.65
(c)	0.70	0.33
(d)	1.99	0.92

<sup>+</sup> See Treatment descriptions.

\* Excluding no fertilizer.