


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1951

[Full Table of Content](#)



51/CA/4 Wheat - Control of Powdery Mildew - Rothamsted Research

Rothamsted Research (1952) *51/CA/4 Wheat - Control of Powdery Mildew - Rothamsted ; Yields Of The Field Experiments 1951*, pp 65 - 67 - DOI: <https://doi.org/10.23637/ERADOC-1-171>

51/Ca/4.1

WHEAT

Control of Powdery Mildew - Sawyers III 1951.

System of replication: 2^5 factorial in 4 blocks of 8 plots each, 3 high order interactions being confounded with block differences.

Area of each plot: 0.0252 acre. Area harvested: 0.0197 acre.

Treatments: All combinations of:-

Sulphate of ammonia: None; 0.6 cwt N per acre.

Superphosphate: None; 0.6 cwt P_2O_5 per acre.

Muriate of potash: None; 1.2 cwt K_2O per acre.

Lime sulphur spray, concentration 1 in 80 applied in three doses, each at the rate of 160 gallons per acre: None; in winter; in late spring; in winter and in late spring.

Basal manuring: None.

Cultivations, etc.: Ploughed: Sept 21. Superphosphate and muriate of potash applied: Oct 26. Seed drilled at 3 bushels per acre: Oct 27. Sulphate of ammonia applied: May 25. Spraying dates: Winter:- Jan 24, Feb 28, Mar 14. Late Spring:- Apr 3, May 14, June 18. Harvested: Aug 27. Variety: Bersee. Previous crop: Wheat.

Standard error per plot:

Grain: 2.08 cwt per acre or 25.1% (9 d.f.)

Notes

- (1) The crop was severely infested with eyespot, take-all and weeds.
- (2) Counts of Powdery Mildew infection were made and are available.

Summary of Results

	Spraying				Mean
	None	In winter	In late winter	and in spring	
Grain: cwt per acre					
Sulphate of ammonia					
None (± 1.04)	6.3	6.9	6.9	7.6	6.9
0.6 cwt N per acre	9.7	9.6	10.4	8.7	9.6
Response to N (± 1.47)	3.4	2.7	3.5	1.1	2.7 ⁽¹⁾
Superphosphate					
None (± 1.04)	6.9	6.9	8.6	8.1	7.6
0.6 cwt P ₂ O ₅ per acre	9.1	9.6	8.8	8.1	8.9
Response to P (± 1.47)	2.2	2.7	0.2	0.0	1.3 ⁽¹⁾
Muriate of potash					
None (± 1.04)	8.5	7.9	9.7	8.6	8.7
1.2 cwt K ₂ O per acre	7.5	8.6	7.7	7.7	7.9
Response to K (± 1.47)	-1.0	0.7	-2.0	-0.9	-0.8 ⁽¹⁾
Mean (± 0.73)	8.0	8.2	8.7	8.1	8.3
Straw: cwt per acre					
Sulphate of ammonia					
None	11.6	11.2	11.8	11.7	11.6
0.6 cwt N per acre	20.7	20.4	22.5	20.0	20.9
Response to N	9.1	9.2	10.7	8.3	9.3
Superphosphate					
None	15.1	14.9	16.9	17.0	16.0
0.6 cwt K ₂ O per acre	17.2	16.6	17.4	14.8	16.5
Response to P	2.1	1.7	0.5	-2.2	0.5
Muriate of Potash					
None	15.8	15.1	17.8	15.5	16.0
1.2 cwt K ₂ O per acre	16.6	16.4	16.5	16.2	16.4
Response to K	0.8	1.3	-1.3	0.7	0.4
Mean	16.2	15.8	17.1	15.9	16.2

(1) ± 0.73

51/Ca/4.3

Responses to Treatments

Response to	Sulphate of ammonia abs.	Sulphate of ammonia pres.	Superphosphate abs.	Superphosphate pres.	Muriate of potash abs.	Muriate of potash pres.
Grain: cwt per acre						
Sulphate of ammonia	-	-	2.8	2.6	2.3	3.1
Superphosphate	1.4	1.2	-	-	0.8	1.8
Muriate of potash	-1.2	-0.4	-1.3	-0.3	-	-
Straw: cwt per acre						
Sulphate of ammonia	-	-	10.0	8.6	8.5	10.1
Superphosphate	1.2	-0.2	-	-	-0.2	1.2
Muriate of potash	-0.4	1.2	-0.3	1.1	-	-