


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1951

[Full Table of Content](#)



51/CA/5 Wheat - Late Application of Nitrogen - Rothamsted

Rothamsted Research

Rothamsted Research (1952) *51/CA/5 Wheat - Late Application of Nitrogen - Rothamsted* ; Yields Of The Field Experiments 1951, pp 68 - 68 - DOI: <https://doi.org/10.23637/ERADOC-1-171>

51/Ca/5

WHEAT

Late application of nitrogen - West Barnfield II, 1951.

System of replication: 3 randomized blocks of 3 plots each.

Area of each plot: 0.0136 acre.

Treatments:

Nitrochalk: None; $1\frac{1}{2}$; 3 cwt per acre applied as a late top dressing.

Basal manuring: $\frac{3}{4}$ cwt superphosphate per acre, drilled with seed, $2\frac{1}{2}$ cwt sulphate of ammonia per acre as a spring dressing.

Cultivations, etc.: Ploughed: Oct 5. Seed drilled at 3 bushels per acre, with superphosphate: Oct 24. Sulphate of ammonia drilled: Mar 25. Nitrochalk applied: July 5. Harvested: Aug 27. Variety: Nord Desprez. Previous crop: Spring Beans.

Standard errors per plot:

Grain: 1.58 cwt per acre or 4.3% (14 d. f.)
 Straw: 2.00 cwt per acre or 4.7% (14 d. f.)

Summary of Results

	Nitrochalk: cwt per acre as top dressing			Mean
	None	$1\frac{1}{2}$	3	
Yield: cwt per acre				
Grain (± 0.56)	36.7	35.9	36.4	36.3
Straw (± 0.71)	42.8	44.0	42.0	42.9
Crude protein: cwt per acre				
Grain	3.53	3.51	3.62	
Increase		-0.02	0.09	
Straw	1.28	1.44	1.36	
Increase		0.16	0.08	
Percentage uptake of added Nitrogen				
Grain		-1	3	
Straw		12	3	