


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1951

[Full Table of Content](#)



51/CC/1 Spring Oats - Late Application of Nitrogen - Rothamsted

Rothamsted Research

Rothamsted Research (1952) *51/CC/1 Spring Oats - Late Application of Nitrogen - Rothamsted ; Yields Of The Field Experiments 1951*, pp 70 - 70 - DOI: <https://doi.org/10.23637/ERADOC-1-171>

51/Cc/1

SPRING OATS

Late application of nitrogen - Great Harpenden II 1951.

System of replication: 8 randomized blocks of 3 plots each.

Area of each plot: 0.0186 acre.

Treatments:

Nitrochalk: None; $1\frac{1}{2}$; 3 cwt per acre applied as a late top dressing.

Basal manuring: $1\frac{1}{2}$ cwt Superphosphate per acre drilled with the seed; $1\frac{2}{3}$ cwt Sulphate of ammonia per acre as a top dressing.

Cultivations, etc.: Ploughed: Dec 23. Seed drilled at 4 bushels per acre with Superphosphate: Apr 19. Sulphate of ammonia applied: June 4. Nitrochalk applied: July 5. Harvested: Aug 23. Variety: Sun II. Previous crop: Spring Oats.

Standard errors per plot:

Grain: 0.814 cwt per acre or 4.4% (14 d.f.)
 Straw*: 1.06 cwt per acre or 7.7% (14 d.f.)

Summary of Results

	Nitrochalk: cwt per acre, as top dressing			Mean
	None	$1\frac{1}{2}$	3	
Yield: cwt per acre				
Grain (± 0.29)	17.7	18.4	18.7	18.3
Straw* (± 0.38)	12.8	13.9	14.8	13.8
Crude protein: cwt per acre				
Grain	1.98	2.18	2.29	
Increase		0.20	0.31	
Straw	0.55	0.74	0.94	
Increase		0.19	0.39	
Percentage uptake of added nitrogen				
Grain		14	11	
Straw		13	14	

*Corrected to 85% dry matter owing to variable conditions during harvesting.