


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1951

[Full Table of Content](#)



51/CH/2 Permanent Grass - Nitrophosphates - Rothamsted

Rothamsted Research

Rothamsted Research (1952) *51/CH/2 Permanent Grass - Nitrophosphates - Rothamsted* ; Yields Of The Field Experiments 1951, pp 90 - 90 - DOI: <https://doi.org/10.23637/ERADOC-1-171>

51/Ch/2

PERMANENT GRASS

Nitrophosphates - Highfield 9 1951.

System of replication: 6 x 6 Latin square.

Area of each plot: 0.0102 acre

Treatments: None; Sulphate of ammonia; Superphosphate; Sulphate of ammonia and superphosphate; British nitrophosphate (12.8% N, 15.25% P₂O₅); Dutch nitrophosphate (20% N, 20.3% P₂O₅). The dressings supply 0.39 cwt N and 0.39 cwt P₂O₅ per acre, the British nitrophosphate receiving extra N to reach this standard.

Basal manuring: 1 1/3 cwt muriate of potash per acre.

Cultivations, etc.: Fertilizer applied: Mar 28. Cut: July 3.

Standard errors per plot:

Hay, dry matter: 1.52 cwt per acre or 5.4% (20 d.f.)

P₂O₅ uptake: 0.00803 cwt per acre or 6.7% (20 d.f.)

Summary of Results

	No fert- ilizer	Sulphate of Ammonia	Super- phosphate	Sulphate of Ammonia and Super- phosphate	British Nitro- phosphate	Dutch Nitro- phosphate	Mean
Hay, dry matter: cwt per acre							
Mean (±0.62)	24.9	29.9	23.5	30.5	29.6	29.8	28.1
Increase (±0.88)		5.0	-1.4	5.6	4.7	4.9	
P ₂ O ₅ uptake: cwt per acre							
Mean (±0.0033)	0.101	0.115	0.109	0.137	0.125	0.134	0.120
Increase (±0.0046)		0.014	0.008	0.036	0.024	0.033	

Mean Dry Matter %: 72.6