


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1952

[Full Table of Content](#)



52/CJ/1 Various - Krilium - Rothamsted and Woburn

Rothamsted Research

Rothamsted Research (1953) *52/CJ/1 Various - Krilium - Rothamsted and Woburn* ; Yields Of The Field Experiments 1952, pp 103 - 105 - DOI: <https://doi.org/10.23637/ERADOC-1-178>

52/Cj/1.1

CLOVER, RYEGRASS, RED BEET AND CARROTS

Krillium - Rothamsted, Fosters and Barnfield; Woburn, Stackyard and Warren Field.

System of replication: 4 x 4 Latin square.

Area of each plot: 0.00207 acre. Area harvested: Clover and Ryegrass - 0.000275 acre, Red beet and Carrots - 0.000689 acre.

Treatments:

Krillium per acre: None; 3 cwt broadcast and twice rotary cultivated in (K_1); 6 cwt broadcast and twice rotary cultivated in (K_2); 3 cwt broadcast and raked into seed bed (K_3).

Basal manuring: 5 cwt Compound fertilizer (7% N, 7% P_2O_5 , 10.5% K_2O) per acre.

Cultivations, etc.: Rothamsted

Ploughed: Feb 13. K_1 and K_2 applied, all plots rotor tilled: May 14. Rotor tilled 2nd time, applied K_3 , seeds drilled: May 15. Dusted red beet with DDT: June 4. 1st cut ryegrass: Fosters - July 25, Barnfield - Aug 27. Clover cut: Aug 29. 2nd cut ryegrass: Sept 18. Carrots lifted: Nov 13. Beet, Barnfield lifted: Nov 14. The beet on Fosters were damaged by sheep and the yields were not taken. Varieties: Clover - Crimson; Ryegrass - Western Walth; Carrots - James Scot Intermediate; Red Beet - Detroit.

Woburn

Ploughed: Stackyard - Sept 17 1951 and Jan 4 1952. Warren Field - Feb 22. K_1 and K_2 applied, all plots rotor tilled: May 14. Applied K_3 and basal manures, rotor tilled, seeds drilled: May 15. Owing to rabbit damage the clover ryegrass and carrots on Warren Field were not harvested. Clover and ryegrass harvested: Aug 22. Beet lifted: Sept 12. Carrots lifted: Oct 30. Varieties: Clover - Scarlet Isles Trifolium; Ryegrass - Italian; Carrots - Scarlet Intermediate; Red Beet - Detroit.

52/Cj/1.2

Standard errors per plot:

Clover, fresh weight.

Fosters:	9.66 cwt per acre or 14.1% (6 d.f.)
Barnfield:	3.73 cwt per acre or 12.0% (6 d.f.)
Stackyard:	19.3 cwt per acre or 31.6% (6 d.f.)

Ryegrass, fresh weight.

Fosters, 1st cut:	2.09 cwt per acre or 22.0% (6 d.f.)
2nd cut:	10.9 cwt per acre or 17.3% (6 d.f.)
Barnfield, 1st cut:	5.86 cwt per acre or 9.9% (5 d.f.)*
2nd cut:	1.37 cwt per acre or 9.7% (6 d.f.)
Stackyard:	14.3 cwt per acre or 11.4% (6 d.f.)

Red Beet, weight of bulbs.

Barnfield:	0.960 tons per acre or 18.8% (6 d.f.)
Stackyard:	3.62 tons per acre or 39.9% (6 d.f.)
Warren Field:	0.436 tons per acre or 8.2% (6 d.f.)

Carrots, roots.

Fosters:	1.02 tons per acre or 7.8% (6 d.f.)
Barnfield:	0.525 tons per acre or 19.9% (6 d.f.)
Stackyard:	1.15 tons per acre or 33.5% (6 d.f.)

*1 missing value.

Summary of Results

	Krilium: cwt per acre				Mean
	None	Broadcast 3	Broadcast in 6	Broadcast in seedbed 3	
Clover, fresh weight: cwt per acre					
	<u>Fosters</u>				
Mean (± 4.83)	67.0	66.8	68.2	72.8	68.7
Increase (± 6.83)		-0.2	+1.2	+5.8	
	<u>Barnfield</u>				
Mean (± 1.87)	30.4	31.0	31.0	32.2	31.2
Increase (± 2.64)		+0.6	+0.6	+1.8	
	<u>Stackyard</u>				
Mean (± 9.6)	60.7	63.2	62.4	58.3	61.1
Increase (± 13.6)		+2.5	+1.7	-2.4	

52/Cj/1.3

	Kriilium: cwt per acre				Mean
	None	Broadcast 3	Broadcast 6	Broadcast in seedbed 3	

Ryegrass, fresh weight: cwt per acre

	<u>Fosters, 1st cut</u>				
Mean (± 1.05)	9.2	9.2	9.7	9.9	9.5
Increase (± 1.48)		0.0	+0.5	+0.7	
	<u>Fosters, 2nd cut</u>				
Mean (± 5.43)	71.1	60.8	61.3	58.2	62.8
Increase (± 7.68)		-10.3	-9.8	-12.9	
	<u>Barnfield, 1st cut</u>				
Mean (± 2.93)	58.7	57.4	58.3	62.1	59.1
Increase (± 4.14)		-1.3	-0.4	+3.4	
	<u>Barnfield, 2nd cut</u>				
Mean (± 0.68)	15.5	14.2	12.4	14.5	14.1
Increase (± 0.97)		-1.3	-3.1	-1.0	
	<u>Stackyard</u>				
Mean (± 7.1)	132.2	123.0	124.5	123.0	125.7
Increase (± 10.1)		-9.2	-7.7	-9.2	

Red Beet, weight of bulbs: tons per acre

	<u>Barnfield</u>				
Mean (± 0.480)	5.56	5.49	4.57	4.83	5.11
Increase (± 0.679)		-0.07	-0.99	-0.73	
	<u>Stackyard</u>				
Mean (± 1.81)	9.01	10.26	9.61	7.47	9.09
Increase (± 2.56)		+1.25	+0.60	-1.54	
	<u>Warren Field</u>				
Mean (± 0.218)	5.61	4.12	5.93	5.64	5.32
Increase (± 0.308)		-1.49	+0.32	+0.03	

Carrots, roots: tons per acre

	<u>Fosters</u>				
Mean (± 0.510)	13.37	11.59	13.77	13.74	13.12
Increase (± 0.721)		-1.78	+0.40	+0.37	
	<u>Barnfield</u>				
Mean (± 0.263)	2.50	2.88	2.31	2.88	2.65
Increase (± 0.371)		+0.38	-0.19	+0.38	
	<u>Stackyard</u>				
Mean (± 0.577)	3.34	3.35	3.49	3.60	3.45
Increase (± 0.816)		+0.01	+0.15	+0.26	