

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1952

[Full Table of Content](#)



52/CA/4 Wheat - Control Wireworm - Rothamsted

Rothamsted Research

Rothamsted Research (1953) *52/CA/4 Wheat - Control Wireworm - Rothamsted* ; Yields Of The Field Experiments 1952, pp 74 - 74 - DOI: <https://doi.org/10.23637/ERADOC-1-178>

WHEAT

The control of wireworm by insecticides - Geescroft 1952.

System of replication: 3 randomized blocks of 8 plots each.

Area of each plot: 0.0289 acre.

Treatments:

O = None (triplicate plots)
 S = Gammexane seed dressing 2 oz per bushel
 G = Gammexane combine drilled with seed at 56 lb per acre 3.5% dust
 A = Aldrin " " " " 200 lb " " 1.78% "
 C = Chlordane " " " " 100 lb " " 5% "
 D = D.D.T. " " " " 150 lb " " 5% "

Basal dressing: 3 cwt sulphate of ammonia per acre.

Cultivations, etc: Ploughed: Nov 9. Seed drilled at 3 bushels per acre with insecticides: Dec 1. Sulphate of ammonia applied: Apr 21. Sprayed with low volume M.C.P.A: May 7. Harvested: Aug 31. Variety: Nord Desprez. Previous crop: Old Grass.

Standard error per plot:

Grain: 1.24 cwt per acre or 3.8% (16 d.f.)

Summary of Results

	O	S	G	A	C	D	Mean
Grain: cwt per acre							
Mean (± 0.71)	31.6 ⁽¹⁾	32.9	33.6	36.3	33.7	31.7	32.9
Increase (± 0.83)		1.3	2.0	4.7	2.1	0.1	
Straw: cwt per acre							
Mean	43.6	46.7	45.6	47.0	44.6	42.6	44.7
Increase		3.1	2.0	3.4	1.0	-1.0	

Standard error (1) 0.41.