


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1952

[Full Table of Content](#)



52/CA/5 Wheat - Late Application of Nitrogen - Rothamsted

Rothamsted Research

Rothamsted Research (1953) *52/CA/5 Wheat - Late Application of Nitrogen - Rothamsted* ; Yields Of The Field Experiments 1952, pp 75 - 76 - DOI: <https://doi.org/10.23637/ERADOC-1-178>

52/Ca/5.1

WHEAT

Late application of nitrogen and methods of harvesting - Fosters 1952.

System of replication: 4 randomized blocks of 3 plots each, blocks and plots being split into 2 strips for different methods of harvesting.

Area of each plot: 0.0242 acre. Area harvested: 'binder' plots - 0.0202 acre, 'combine' plots - 0.0091 acre.

Treatments:

Nitrochalk: None; $1\frac{1}{2}$; 3 cwt per acre applied as top dressing.
Methods of harvesting: Binder; N.I.A.E. combine harvester.

Basal manuring: 3 cwt sulphate of ammonia per acre.

Cultivations, etc.: Ploughed: Sept 11. Seed drilled at 3 bushels per acre: Oct 24. Sulphate of ammonia applied: Apr 18. Sprayed with D.N.O.C.: May 15. Nitrochalk applied: June 28. Harvested: 'binder' plots - Aug 12, 'combine' plots - Aug 13. Variety: Yeoman. Previous crop: Wheat.

Standard errors per plot. Grain:

Strip: 1.72 cwt per acre or 8.9% (3 d.f.)
Whole plot: 1.32 cwt per acre or 6.9% (6 d.f.)
Sub plot: 1.68 cwt per acre or 8.7% (6 d.f.)

52/Oa/5.2

Summary of Results

	Nitrochalk: cwt per acre as top dressing			Mean
	None	1.5	3.0	
Grain: cwt per acre (a) and (b)				
Binder	19.0	20.9	20.8	20.2
Combine	18.7	18.4	17.5	18.2
Mean (± 0.66)	18.8	19.7	19.2	19.2
Difference (± 1.55)	-0.3	-2.5	-3.3	-2.0 (± 1.21)

(a) ± 0.89 for use in horizontal comparisons only

(b) ± 1.13 for use in diagonal comparisons only

Binder Plots

Straw: cwt per acre				
	33.7	34.1	35.9	34.6
Crude Protein: cwt per acre				
Grain	2.11	2.44	2.43	
Increase		0.33	0.32	
Straw	0.68	0.68	0.74	
Increase		0.00	0.06	
Percentage uptake of added nitrogen				
Grain		23	11	
Straw		1	2	