


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1952

[Full Table of Content](#)



---

### 52/CB/1 Barley - Late Application of Nitrogen - Rothamsted

#### Rothamsted Research

Rothamsted Research (1953) *52/CB/1 Barley - Late Application of Nitrogen - Rothamsted* ; Yields Of The Field Experiments 1952, pp 77 - 78 - DOI: <https://doi.org/10.23637/ERADOC-1-178>

52/Cb/1.1

BARLEY

Late application of nitrogen and methods of harvesting - Great Field II 1952.

System of replication: 4 randomized blocks of 3 plots each, blocks and plots being split into 2 strips for different methods of harvesting.

Area of each plot: 0.0242 acre. Area harvested: 'binder' plots - 0.0242 acre, 'combine' plots - 0.0091 acre.

Treatments:

Nitrochalk: None;  $1\frac{1}{2}$ ; 3 cwt per acre applied as top dressing.  
Methods of harvesting: Binder; N.I.A.E. combine harvester.

Basal manuring per acre:  $2\frac{1}{4}$  cwt sulphate of ammonia; 1 cwt super-phosphate combine drilled with seed.

Cultivations, etc.: Sulphate of ammonia applied to linseed: Mar 19.  
Harrowed in linseed which had failed: May 12. Seed and super-phosphate drilled: May 13. Sprayed with M.C.P.A., medium volume, 5 pints in 30 gallons water per acre: June 9. Nitrochalk applied: June 16. Harvested: 'binder' plots - Sept 16, 'combine' plots - Sept 19. Variety: Plumage Archer. Previous crop: Permanent grass.

Standard errors per plot. Grain:

Strip: 1.09 cwt per acre or 7.6% (3 d.f.)  
Whole plot: 1.05 cwt per acre or 7.4% (6 d.f.)  
Sub plot: 1.18 cwt per acre or 8.3% (6 d.f.)

52/Cb/1.2

Summary of Results

	Nitrochalk: cwt per acre as top dressing			Mean
	None	1.5	3.0	
Grain: cwt per acre				
	(a) and (b)			
Binder	13.1	13.5	13.1	13.2
Combine	14.7	15.6	15.5	15.3
Mean ( $\pm 0.53$ )	13.9	14.5	14.3	14.2
Difference ( $\pm 1.03$ )	1.6	2.1	2.4	2.1 ( $\pm 0.77$ )

(a)  $\pm 0.67$  for use in horizontal comparisons only  
 (b)  $\pm 0.79$  for use in diagonal comparisons only

	Straw: cwt per acre			
Binder	19.6	18.3	19.2	19.0

	Crude Protein: cwt per acre		
Grain: Binder	1.76	1.84	1.78
Increase		0.08	0.02
Combine	2.07	2.16	2.14
Increase		0.09	0.07
Straw: Binder	1.17	1.11	1.12
Increase		-0.06	-0.05

	Percentage uptake of added nitrogen	
Grain: Binder	6	1
Combine	6	2
Straw: Binder	-4	-2