


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1952

[Full Table of Content](#)



52/CC/1 Spring Oats - Late Application of Nitrogen - Rothamsted

Rothamsted Research

Rothamsted Research (1953) *52/CC/1 Spring Oats - Late Application of Nitrogen - Rothamsted* ;
Yields Of The Field Experiments 1952, pp 80 - 80 - DOI: <https://doi.org/10.23637/ERADOC-1-178>

SPRING OATS

Late application of nitrogen - Long Hoos III 1952.

System of replication: 8 randomized blocks of 3 plots each.

Area of each plot: 0.0145 acre.

Treatments: Nitrochalk: None; $1\frac{1}{2}$; 3 cwt per acre applied as a late top dressing.

Basal dressing: $2\frac{1}{2}$ cwt sulphate of ammonia per acre; 1 cwt superphosphate per acre drilled with the seed.

Cultivations, etc.: Ploughed: Dec 10. Sulphate of ammonia applied: Mar 3. Seed drilled at 4 bushels per acre with superphosphate: Mar 15. Sprayed with M.C.P.A. low volume, $2\frac{1}{2}$ pints in 10 gallons of water: May 13. Nitrochalk applied: June 30. Harvested: July 24. Variety: Star. Previous crop: Wheat.

Standard errors per plot:

Grain*: 1.74 cwt per acre or 4.9% (14 d.f.)
 Straw*: 2.39 cwt per acre or 5.2% (14 d.f.)

Summary of Results

	Nitrochalk: cwt per acre, as top dressing			Mean
	None	$1\frac{1}{2}$	3	
Yield: cwt per acre				
Grain* (± 0.62)	35.5	36.4	35.2	35.7
Straw* (± 0.84)	45.5	45.5	45.6	45.5
Crude protein: cwt per acre				
Grain	3.57	3.87	3.81	
Increase		0.30	0.24	
Straw	0.87	0.92	0.97	
Increase		0.05	0.10	
Percentage uptake of added nitrogen				
Grain		21	8	
Straw		3	3	

*Corrected to 85% dry matter owing to variable conditions during harvesting.