Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1952



Full Table of Content

52/CD/2 Spring Beans - Control of Back Aphis - Rothamsted

Rothamsted Research

Rothamsted Research (1953) 52/CD/2 Spring Beans - Control of Back Aphis - Rothamsted; Yields Of The Field Experiments 1952, pp 83 - 83 - DOI: https://doi.org/10.23637/ERADOC-1-178

52/ca/2

SPRING BEANS

Control of Black Aphis - Great Harpenden II 1952.

System of replication: 2* randomized blocks of 7 plots each.

Area of each plot: 0.00831 acre.

Treatments - Insecticidal sprays at 300 gallons per acre.

None.

Parathion, 0.02% technical W/V.

Pyrethrum, 0.05% W/V total pyrethrins.

DDT emulsion, 0.1% W/V DDT.

DDT suspension, 0.1% W/V DDT

Systox, 0.05% W/V.

Nicotine, 0.1% V/V technical nicotine alkaloid. (F)

Basal dressing, per acre: 1½ cwt nitrochalk, 3 cwt superphosphate, 1½ cwt sulphate of potash.

Cultivations, etc.: Ploughed: Oct 30. 40 cwt ground chalk per acre applied: Mar 13. Basal fertilizers applied: Mar 15. Beans drilled at 4 bushels per acre: Mar 18. Spray treatments applied: June 12. Harvested: Aug 2. Variety: Ashwells selection. Previous crop: Oats.

Standard error per plot:
Grain: 4.50 cwt per acre or 49.7% (6 d.f.)

Note. The experiment was originally designed as one of 4 blocks of 7, but as at spraying time 2 of the blocks carried a very poor crop these were not used in the experiment.

Summary of Results

	Spray							
	0	A	В	C	D	E	F	Mean
Grain: cwt per acre								
Mean (±3.19)	1.	4 11.8	10.3	11.8	1.3	12.0	14.8	9.1
Increase (±4.50)		10.4	8.9	10.4	-0.1	10.6	13.4	
Straw: cwt per acre								
Mean	14.	2 19.3	14.4	15.4	15.4	16.7	18.3	16.2
Increase		5.1	0.2	1.2	1.2	2.5	4.1	