


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1952

[Full Table of Content](#)



52/CD/3 Broad Beans - Fertilizer Placement - Rothamsted

Rothamsted Research

Rothamsted Research (1953) *52/CD/3 Broad Beans - Fertilizer Placement - Rothamsted* ; Yields Of The Field Experiments 1952, pp 84 - 84 - DOI: <https://doi.org/10.23637/ERADOC-1-178>

52/Ca/3

BROAD BEANS

Fertilizer placement - Great Harpenden II 1952.

System of replication: 4 randomized blocks of 5 plots each.

Area of each plot: 0.00344 acre.

Treatments: No fertilizer; and all combinations of:-

Compound granular PK fertilizer (13 $\frac{3}{4}$ % P₂O₅; 13 $\frac{3}{4}$ % K₂O): 2.7; 4.9 cwt per acre.

Method of placement: Broadcast on seed bed; drilled in band beside seed.

Basal dressing per acre: 1 $\frac{1}{2}$ cwt nitrochalk; 3 cwt superphosphate; 1 $\frac{1}{2}$ cwt sulphate of potash.

Cultivations, etc.: Ploughed: Oct 30. 40 cwt ground chalk per acre applied: Mar 13. Basal fertilizers applied: Mar 15. Beans sown: Mar 18. Harvested: July 1. Variety: Seville Long Pod. Previous crop: Spring Oats.

Standard error per plot:

Yield of green beans in pod: 2.45 cwt per acre or 9.1% (12 d.f.)

Summary of Results

	Compound PK fertilizer: cwt per acre					Mean
	None	2.7 Broadcast	2.7 Drilled	4.9 Broadcast	4.9 Drilled	
Yield of green beans in pod: cwt per acre						
Mean (± 1.22)	22.8	24.5	29.5	26.6	31.8	27.0
Increase (± 1.73)		1.7	6.7	3.8	9.0	