

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1952

[Full Table of Content](#)



52/CH/1 Globe Beet - Fertilizer Placement - Rothamsted

Rothamsted Research

Rothamsted Research (1953) *52/CH/1 Globe Beet - Fertilizer Placement - Rothamsted* ; Yields Of The Field Experiments 1952, pp 98 - 98 - DOI: <https://doi.org/10.23637/ERADOC-1-178>

52/Ch/1

GLOBE BEET

Fertilizer placement - Long Hoos IV 1952.

System of replication: 4 randomized blocks of 5 plots each.

Area of each plot: 0.00344 acre.

Treatments: No fertilizer; and all combinations of:-

National Compound fertilizer No. 1A (8% N; 6% P₂O₅; 10½% K₂O):
2.5; 5.0 cwt per acre.

Method of placement: Broadcast on seed bed; drilled in band
beside seed.

Note. On calibration, the levels of fertilizer actually drilled were
found to be 2.0 and 4.5 cwt per acre.

Basal dressing: None.

Cultivations, etc.: Ploughed: Feb 5. Seed drilled at 8 lb per acre,
fertilizer applied: May 13. Lifted: Aug 6. Variety: Crimson
Globe. Previous crop: Barley.

Standard error per plot:

Saleable produce*: 1.60 tons per acre or 45.2% (12 d.f.)

Summary of Results

	Compound fertilizer: cwt per acre					Mean
	None	2.5 Broadcast	2.0 Drilled	5.0 Broadcast	4.5 Drilled	
Saleable produce*: tons per acre						
Mean (±0.80)	3.99	3.18	3.25	3.20	4.03	3.53
Increase (±1.13)		-0.81	-0.74	-0.79	+0.04	
Total produce*: tons per acre						
Mean	5.50	4.72	4.58	4.65	5.31	4.95
Increase		-0.78	-0.92	-0.85	-0.19	

*Bulbs and tops.