Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1953



Full Table of Content

53/A/8 Stackyard Woburn - Barley

Rothamsted Research

Rothamsted Research (1954) 53/A/8 Stackyard Woburn - Barley; Yields Of The Field Experiments 1953, pp 12 - 12 - DOI: https://doi.org/10.23637/ERADOC-1-173

53/1/8

BARLEY - WOBURN STACKYARD 1953

For history, details of treatments etc. see "Results of the Field Experiments 1939-47" Vol. I, Section A/7 with the exception that in 1952 it was decided to divide each plot into two, and sow one half with a Winter variety, and the other as before with a Spring variety.

Cultivations, etc.: Ploughed: Aug 26, 1952. 'Winter' seed drilled at 3 bushels per acre: Oct 9. 'Spring' seed drilled at 3 bushels per acre: Mar 5, 1953. Nitrochalk applied: Mar 13. Sprayed with DNOC 6 lb in 80 gallons per acre: May 22. Harvested: Aug 17. Varieties: Winter - Pioneer, Spring - Plumage Archer.

Summary of Results

Plot	Nitrochalk	Grain: cwi	per acre	Straw: cw	per acre
	dressing	Winter	Spring	Winter	Spring
	cwt per acre	sown	sown	sown	sown
3	2	16.3	8.4	12.8	7.6
7	4	9.0	15.9	10.6	13.6
1	6	8.3	16.6	11.8	14.0
9 4 6	2	19.8	8.9	14.8	9.0
	4	17.6	18.9	21.7	16.2
	6	25.9	25.2	26.9	22.2
11b(1)	2	20.3	14.2	21.0	12.1
11b(2)	4	26.4	19.7	28.4	17.9
11b(3)	6	31.1	24.4	34.5	22.0
11a	2	15.3	12.1	14.1	9.1
10a	4	6.6	8.7	10.1	9.1
10b	6	15.1	11.6	16.1	9.8