


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1953

[Full Table of Content](#)



## 53/BA/1 Three-course Rotation - Rothamsted

### Rothamsted Research

Rothamsted Research (1954) *53/BA/1 Three-course Rotation - Rothamsted ; Yields Of The Field Experiments 1953*, pp 13 - 19 - DOI: <https://doi.org/10.23637/ERADOC-1-173>

53/Ba/1.1

THREE COURSE ROTATION EXPERIMENT

2nd year of revised scheme

For details of rotation, treatments etc., see "Results of the Field Experiments 1952", Section 52/Ba/1.

Area of each plot: Potatoes (sub-plot) - 0.0092 acre; Barley - 0.0200 acre; Sugar beet - 0.0205 acre.

Cultivations, etc.:

Potatoes: Straw applied, ploughed all plots: Jan 22, 1953.

Fertilizers applied: Mar 25. Ridged: Mar 27. Potatoes

planted with mechanical dropper: Apr 11. Earthed up ridges:

June 25. Sprayed with copper fungicide,  $5\frac{1}{2}$  lb per acre:

July 28 and again Aug 8. Haulm destroyed mechanically:

Sept 17. Lifted: Sept 28. Variety: Majestic.

Barley: Straw applied, ploughed all plots: Jan 22, 1953. Seed drilled at 3 bushels per acre, fertilizers applied: Feb 28.

Harvested: Aug 11. Variety: Plumage Archer.

Sugar beet: Straw applied, ploughed all plots: Jan 22, 1953.

Fertilizers applied: Mar 18. Seed drilled at 18 lb per acre:

Mar 26. Sprayed with D.D.T. emulsion, 3 pints in 10 gallons:

May 11. Singled: May 31. Lifted: Nov 12. Variety: Klein E.

Treatment symbols

Ar Complete artificials only.

St1 Straw ploughed in in autumn, artificials applied in spring.

St2 Straw ploughed in in autumn, artificials applied half in autumn, half in spring.

Ad Adco ploughed in in autumn with supplementary artificials.

St  $53\frac{1}{3}$  cwt cut straw in autumn.

Nitrogen dressing: 0.2, 0.4, 0.6 cwt N as sulphate of ammonia.

K<sub>s</sub> Muriate of potash equivalent to K<sub>2</sub>O in straw.

K 0.5 cwt K<sub>2</sub>O as muriate of potash.

53/Ba/1.2

## Summary of Results

## Potatoes

Treatments applied	1953	1953	0	0.4N	St + 0.2N	St + 0.6N	$K_s$ + 0.4N
	-	-	K	-	K	-	K
Ar	0	0.4N	8.49	8.24	9.92	10.79	
Ar	0	0.4N	7.54	8.44	10.04	10.28	
St1	St2	0	9.60	8.68	11.54	11.88	
St1	St2	0.4N	9.60	8.68	9.31	9.21	
St1	St2	0	9.70	9.85	9.75	10.28	
St1	St2	0.4N	9.70	9.85	10.67	10.62	
St1	St2	St+0.2N	9.17	8.88	9.36	9.99	
St1	St2	St+0.6N	10.43	10.09	10.33	9.60	
Ad	Ad	0	0	0	10.09	8.78	
Ad	Ad	0.4N	8.34	9.31			
Ad	Ad	St+0.6N	8.29	7.76			
Ad	Ad	$K_s$ 0.4N	11.30	11.15			

53/Ba/1.3

				Potatoes					
Treatments applied		1953	1952	0		0.4N	St + 0.2N	St + 0.6N	K <sub>S</sub> + 0.4N
		1953	-	K	-	K	-	K	-
1950	1951	Ar	0	82.5	87.3	86.8	84.6		
		Ar	0.4N	82.2	86.3	85.3	87.4		
		St1	0	82.0	82.8	87.7	91.2	89.6	
		St2	0	82.0	82.8	84.9	84.8	88.3	
		St1	0	89.6	84.9	84.9	84.3		
		St2	0.4N	87.4	86.4	87.6	85.5		
		K <sub>S</sub>	0.4N	84.8	88.1	86.9	90.1		
		Ad	0	88.4	82.8	88.4	82.8		
		Ad	0.4N	81.4	87.7	85.9	84.6		
			St+0.6N	84.6	84.6	84.6	83.8		
			K <sub>S</sub> 0.4N	84.6	83.8				

53/Ba/1.4

Barley

Treatments applied		1953	0	0.4N	St + 0.2N	St + 0.6N	K <sub>s</sub>	K <sub>s</sub> + 0.4N
1950	1951	1952	Grain: cwt per acre					
	Ar	0		31.2				
		0.4N		28.6				
Ar		0		30.2				
		0.4N		31.5				
	St1 St2	0		27.4				
		0.4N		21.9		31.4		27.3
St1 St2		0		26.0		26.3		
		0.4N		30.4				
		St+0.2N		30.3				
		St+0.6N		30.7				
		K <sub>s</sub>		30.6				
		0.4N		31.3				
	Ad	0		26.0				
		0		28.0				
Ad		0.4N		30.0				31.4
		St+0.6N						
		K <sub>s</sub> 0.4N		22.3				
Straw: cwt per acre								
	Ar	0		44.8				
		0.4N		31.9				
Ar		0		41.3				
		0.4N		33.7				
	St1 St2	0		40.2				
		0.4N		25.9		41.4		36.8
St1 St2		0		30.0		29.6		
		0.4N		41.9				
		St+0.2N		37.1				
		St+0.6N		46.5				
		K <sub>s</sub>		38.0				
		0.4N		40.8				
	Ad	0		39.7				
Ad		0		35.9		41.7		39.4
		0.4N						
		St+0.6N		27.7				
		K <sub>s</sub> 0.4N		30.2				
				25.5				

53/Ba/1.5

## Sugar beet

Treatments applied	1953	0	0.4N	St + 0.2N	St + 0.6N	K <sub>S</sub>	K <sub>S</sub> + 0.4N
1950	1951	1952	Roots (washed): tons per acre				
Sugar percentage							
Ar	0	15.84					
	0.4N	12.07					
Ar	0	15.12					
	0.4N	12.45					
St1 St2	0	16.04		14.25		14.53	
	0.4N	12.09		12.31		14.62	
St1 St2	0	15.19					
	0.4N	13.73		14.82			
	St+0.2N						
	St+0.6N	12.81					
	K <sub>S</sub>	15.39					
	0.4N	13.05					
Ad	0	15.63		13.64		16.11	
	0.4N	12.31					
	St+0.6N	13.36					
	K <sub>S</sub> 0.4N	12.83					

53/Ba/1.6

Sugar beet

Treatments applied	1953	0	0.4N	St + 0.2N	St + 0.6N	K <sub>s</sub>	K <sub>s</sub> + 0.4N
1950	1951	1952	Total sugar: cwt per acre				
Tops: tons per acre							
Ar	0		57.0				
	0.4N	45.0					
Ar	0		55.9				
	0.4N	46.5					
St1 St2	0		60.2		53.3		53.7
	0.4N	45.0		45.4		54.1	
St1 St2	0		55.4				
	0.4N	49.8					
St1 St2	St+0.2N		54.6				
	St+0.6N	47.1					
Ad	K <sub>s</sub>		54.6				
	0.4N	47.6					
Ad	0		57.2		50.6		57.2
	0.4N	45.0					
Ad	St+0.6N	47.6					
	K <sub>s</sub> 0.4N	47.1					

53/Ba/1.7

Sugar beet

Treatments applied	1953	0	0.4N	St + 0.2N	St + 0.6N	K <sub>s</sub>	K <sub>s</sub> + 0.4N
1950	1951	1952	Plant number: thousands per acre				
Ar	0		26.9				
	0.4N	29.2					
Ar	0		29.6				
	0.4N	28.2					
St1 St2	0		28.4	28.9	27.3		28.0
	0.4N	28.4				28.5	
St1 St2	0		27.9				
	0.4N	28.3					
	St+0.2N		25.9				
	St+0.6N	27.5					
	K <sub>s</sub>		26.7				
	0.4N	27.7					
Ad	0		28.2		27.3		28.3
	0.4N	29.6					
	St+0.6N	28.6					
	K <sub>s</sub>	29.0					
Noxious nitrogen: mg. per 100 g.							
Ar	0		15.0				
	0.4N	15.0					
Ar	0		15.0				
	0.4N	15.0					
St1 St2	0		15.0	15.0	15.0		15.0
	0.4N	15.0				15.0	
St1 St2	0		15.0				
	0.4N	15.0					
	St+0.2N		25.0				
	St+0.6N	15.0					
	K <sub>s</sub>		25.0				
	0.4N	15.0					
Ad	0		15.0		25.0		15.0
	0.4N	15.0					
	St+0.6N	15.0					
	K <sub>s</sub>	25.0					