

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1953

[Full Table of Content](#)



Classical Experiments

Rothamsted Research

Rothamsted Research (1954) *Classical Experiments* ; Yields Of The Field Experiments 1953, pp 4 - 12 - DOI: <https://doi.org/10.23637/ERADOC-1-173>

53/A/1

WHEAT - BROADBALK 1953

The 110th year

For history, details of treatments etc. see "Results of the Field Experiments 1939-47" Vol. I, Section A/1.

Cultivations, etc.:

Cropped sections. Dung applied and all plots ploughed: Sept 3, 1952.
Autumn fertilizers applied: Nov 11. Seed drilled at 3 bushels per acre: Nov 12. Spring fertilizers applied: Apr 23, 1953.
Second dressing of nitrate of soda applied to plot 16: May 13.
Harvested: Aug 25. Variety: Squareheads Master 13/4.

Fallow section. Ploughed: Sept 3, 1952, Mar 22 and June 10, 1953.

Summary of Results

Section Years after Fallow	Grain: cwt per acre					Straw: cwt per acre				
	II	I	III	IV	Mean	II	I	III	IV	Mean
	1	2	3	4		1	2	3	4	
2A	29.3	20.9	23.0	25.0	24.6	60.3	48.1	46.3	47.7	50.6
2B	24.8	22.4	23.4	25.0	23.9	57.9	62.1	46.2	53.5	54.9
3	16.5	13.7	13.2	12.4	14.0	35.1	28.8	21.0	21.2	26.5
5	11.6	12.9	8.4	13.8	11.7	32.6	33.9	22.8	26.0	28.8
6	17.9	17.3	16.3	14.9	16.6	46.5	40.6	35.2	32.5	38.7
7	16.3	22.3	22.6	20.4	20.4	46.9	53.4	47.0	44.0	47.8
8	21.9	22.6	22.9	28.1	23.9	48.8	58.8	53.8	55.7	54.3
9	19.9	18.5	17.9	16.4	18.2	45.8	46.1	35.3	30.3	39.4
10	16.4	25.5	20.2	19.1	20.3	34.7	44.4	35.4	35.8	37.6
11	16.0	24.0	19.8	17.2	19.2	31.7	41.2	35.8	34.1	35.7
12	18.6	24.7	21.4	19.0	20.9	36.3	50.3	41.1	38.8	41.6
13	24.2	24.3	22.6	18.6	22.4	46.6	47.9	45.3	35.7	43.9
14	16.8	23.5	22.6	19.1	20.5	37.9	39.8	41.5	39.8	39.8
15	25.7	19.9	19.4	14.9	20.0	51.6	37.9	40.2	31.0	40.2
16	24.1	22.5	20.3	17.6	21.1	51.9	49.6	48.7	48.9	49.8
17	20.2	14.7	14.9	8.5	14.6	36.2	25.5	25.4	19.8	26.7
18	23.9	26.1	20.7	21.0	22.9	49.2	47.5	44.5	42.8	46.0
19	19.3	19.5	15.8	15.4	17.5	38.1	40.7	35.1	32.6	36.6
20	21.1	27.8	-	-	24.4	35.5	49.8	-	-	42.6

53/A/2

BARLEY - HOOSFIELD 1953

The 102nd year

For history, details of treatments etc. see "Results of the Field Experiments 1939-47" Vol. I, Section A/2.

Cultivations, etc.: Ploughed: Sept 19, 1952. Dung applied and ploughed in: Sept 23. Fertilizers applied: Mar 3, 1953. Seed drilled at 3 bushels per acre: Apr 9. Sprayed with D.N.O.C.: May 26. Variety: Plumage Archer.

Note: Owing to the very heavy infestation with Wild Oats (*Avena Fatua*) the whole crop was cut and carted green before the oats ripened. No yields therefore were recorded.

53/A/3

WHEAT AFTER FALLOW - HOOSFIELD 1953

Without manure 1851 and since

For history, details of treatments etc. see "Results of the Field Experiments 1939-47", Vol. I, Section A/3.

Cultivations, etc.:

Cropped plots. Ploughed: Sept 6, 1952. Seed drilled at 3 bushels per acre: Oct 27. Harvested: Aug 20, 1953. Variety: Squareheads Master 13/4.

Fallow plots. Ploughed: Sept 6, 1952 and Sept 3, 1953.

Summary of Results

Plot No. of years of fallow	Mean yields: cwt per acre			Mean
	B ₂ 1	B ₃ 1	B ₄ 3	
Grain	18.7	8.3	13.7	13.6
Straw	24.9	14.5	23.6	21.0

CROPS IN ROTATION - AGDELL 1953

Barley - 2nd crop of 27th course 1952-55

For history, details of treatments etc. see "Results of the Field Experiments 1939-47", Vol. I, Section A/4.

Cultivations, etc.: Ploughed: Sept 5, 1952, Seed drilled at 3 bushels per acre: Mar 2, 1953. Sprayed with MCPA, 2 pints per acre: May 26. Harvested: Aug 28. Variety: Plumage Archer.

Note: No turnips were grown and no manures were applied in 1952, the plots being fallowed.

Summary of Results

Manure to Turnips 1948	Mean yields: cwt per acre					
	None since 1848		Mineral manure No nitrogen		Complete mineral and nitrogenous manure	
	Fallow 5	Clover 6	Fallow 3	Clover 4	Fallow 1	Clover 2
Grain	5.7	11.1	7.6	16.2	3.6	5.0
Straw	5.8	11.5	8.3	16.4	4.8	8.9

MANGOLDS AND SUGAR BEET - BARNFIELD 1953

53/A/4.1

The 78th and 8th years

For history, details of treatments etc, see "Results of the Field Experiments 1939-47" Vol.I, Section A/5.

Cultivations, etc.: Dung applied: Jan 16. Ploughed: Jan 20.
Fertilizers applied: Mar 19. Seed drilled, mangolds - 9 lb per acre, sugar beet - 18 lb per acre: Apr 10. Sprayed with D.D.T. emulsion: May 11. Singling commenced: June 22. Top dressings applied: July 14. Lifted: mangolds - Oct 29, sugar beet - Nov 11. Varieties: mangolds - Yellow Globe, sugar beet - Klein E.

Erratum to "Results of the Field Experiments 1939-47" Vol.I, page A/5.1, 6th line up: dressing for plot 4(b) should read "PCa" not "NPCa".

Summary of Results

Strip	Cross Dressing				
	O	N	A	AC	C
Mangolds, Roots: tons per acre					
1	7.02	26.33	29.20	26.20	27.83
2	15.18	31.31	30.11	33.65	33.08
4	3.07	(a) 22.32 (b) 20.79*	17.85	20.74	26.26
5	2.33	19.18	12.96	13.61	21.61
6	2.43	17.75	14.93	24.72	25.46
7	1.94	21.05	19.42	27.83	27.83
8	1.19	15.13	10.28	10.53	17.81
9	19.69				
Mangolds, Leaves: tons per acre					
1	1.15	2.52	5.26	4.09	4.80
2	2.35	2.84	3.79	3.94	3.45
4	0.71	(a) 2.50 (b) 2.08*	2.94	2.72	2.84
5	0.64	1.91	2.45	2.89	3.01
6	0.69	1.74	1.96	3.47	2.74
7	0.66	2.10	2.84	4.28	3.33
8	0.71	2.08	2.40	3.43	3.79
9	2.25				
Mangolds, Plant number: thousands per acre					
1	14.4	18.0	16.9	15.9	17.8
2	18.7	18.2	18.6	17.3	17.1
4	15.7	(a) 19.5 (b) 20.7*	18.9	17.9	18.8
5	16.6	18.5	17.8	16.6	18.8
6	15.6	19.3	19.1	18.4	19.1
7	14.3	19.4	19.4	20.3	18.8
8	13.9	18.4	16.7	15.6	18.2
9	20.4				

* No nitrate of soda. Nitrogen applied as calcium and potassium nitrates.

53/A/4.2

Strip	Cross Dressing				
	O	N	A	AC	C
Sugar Beet, Roots: tons per acre					
1	7.28	18.24	18.67	18.32	19.18
2	7.71	15.24	13.02	15.76	17.13
4	2.46	(b) 12.33 [‡]	9.59	15.93	14.98
5	2.50	9.08	8.56	12.76	13.61
6	2.02	9.59	8.82	16.18	12.50
7	1.82	10.96	10.36	16.01	12.42
8	1.63	9.33	8.22	11.05	10.36
Sugar Beet, Tops: tons per acre					
1	3.03	8.61	11.50	10.28	8.91
2	3.72	7.24	7.19	8.56	7.29
4	1.08	(b) 5.19 [‡]	3.72	8.86	6.65
5	1.13	4.55	4.45	10.28	6.65
6	0.83	4.99	4.75	7.54	5.68
7	0.64	5.24	4.99	10.67	5.28
8	0.83	5.09	4.75	11.69	5.68
Sugar Beet, Plant number: thousands per acre					
1	18.6	21.5	20.4	19.8	20.2
2	21.4	21.6	22.1	20.6	19.3
4	20.6	(b) 21.8 [‡]	22.5	21.1	20.0
5	19.8	21.9	22.3	20.0	19.9
6	19.7	22.3	21.1	21.7	20.7
7	20.5	22.3	23.2	23.0	21.3
8	21.2	22.9	22.3	21.9	19.6
Sugar Beet, Sugar percentage					
1	18.00	16.91	17.45	17.22	17.66
2	17.37	17.37	17.02	16.88	17.43
4	19.13	(b) 18.06 [‡]	17.60	17.48	17.86
5	18.12	16.39	17.66	17.25	17.14
6	17.83	16.99	17.51	17.14	18.72
7	17.71	18.12	17.83	17.77	18.09
8	17.22	16.85	17.60	17.05	17.37

[‡] No nitrate of soda. Nitrogen applied as calcium and potassium nitrates.

HAY - THE PARK GRASS PLOTS 1953

The 98th year

For history, details of treatments etc. see "Results of the Field Experiments 1939-47", Section A/6.

Cultivations, etc.: Dung applied: Dec 2, 1952. Mineral fertilizers applied: Dec 11. Nitrogenous fertilizers applied: 1st dressing - Mar 23, 1953, 2nd dressing - April 22. Cut: 1st - July 21, 2nd - Nov 19.

Summary of Results

Yield of Hay: cwt per acre

Plot	Not limed			Limed		
	1st Crop	2nd Crop	Total	1st Crop	2nd Crop	Total
1	14.5	4.1	18.6	20.4	2.8	23.2
2	17.2	0.0	17.2	19.7	0.0	19.7
3	16.1	0.0	16.1	16.7	0.0	16.7
4-1	20.6	1.9	22.5	20.0	1.9	21.9
4-2	22.3	4.5	26.8	36.3	4.5	40.8
5-1	10.0	3.0	13.0			
5-2	26.4	4.1	30.5			
6	36.6	4.4	41.0			
7	30.0	4.4	34.4	34.1	4.9	39.0
8	24.9	1.0	25.9	18.8	0.6	19.4
9	43.0	1.5	44.5	47.2	2.2	49.4
10	20.8	2.5	23.3	39.6	3.4	43.0
11-1	45.5	13.4	58.9	61.2	8.2	69.4
11-2	63.4	11.9	75.3	71.6	10.9	82.5
12	19.0	0.0	19.0			
13	41.5	5.2	46.7	36.9	5.6	42.5
14	60.6	8.1	68.7	60.6	4.6	65.2
15	27.4	2.2	29.6	33.7	3.9	37.6
16	47.2	3.0	50.2	45.7	2.2	47.9
17	26.7	3.8	30.5	25.0	1.1	26.1
18	18.9	3.1	22.0	27.0*	2.2*	29.2*
				31.4†	2.9†	34.3†
19	63.3	4.6	67.9	23.7*	3.0*	26.7*
				43.6†	3.4†	47.0†
20	38.5	3.4	41.9	38.3*	2.5*	40.8*
				39.8†	2.4†	42.2†

*Heavy liming

†Light liming

Note: The second crop was carted green; hay yields were estimated from the dry matter.

53/A/6

BARLEY - EXHAUSTION LAND HOOSFIELD 1953

For history, details of treatments etc. see "Results of the Field Experiments 1952", Section A/6.

Cultivations, etc.: Ploughed: Sept 4, 1952. $2\frac{1}{2}$ cwt sulphate of ammonia per acre applied: Feb 26, 1953. Seed drilled at 3 bushels per acre: Feb 27. Sprayed with M.C.P.A. 2 pints per acre: May 26. Variety: Plumage Archer.

Summary of Results

Manuring to Potatoes 1876-1901*	Grain cwt per acre	Straw cwt per acre
1 Unmanured	15.8	18.2
2 Unmanured after 6 years dung	11.8	18.0
3 Dung	22.4	25.6
4 Dung	22.8	26.4
5 Ammonium salts	17.4	19.3
6 Nitrate of soda	14.6	18.2
7 Ammonium salts and complete minerals	21.4	24.2
8 Nitrate of soda and complete minerals	20.2	23.4
9 Superphosphate	20.0	24.4
10 Complete minerals	21.8	25.7

*For certain changes see history.

53/4/7

WHEAT - WOBURN STACKYARD 1953

For history, details of treatments etc., see "Results of the Field Experiments 1939-47" Vol I, Section 4/7.

Cultivations, etc.: Ploughed: Aug 27, 1952. Seed drilled at 3 bushels per acre: Oct 10. Nitrochalk applied: Apr 20, 1953. Sprayed with DNOC 6 lb in 80 gallons: May 22. Harvested: Aug 18. Variety: Squareheads Master (13/4).

Summary of Results

Plot	Nitrochalk dressing cwt per acre	Grain: cwt per acre	Straw: cwt per acre
1	2	7.9	16.0
7	4	10.1	20.8
3	6	13.8	21.6
9	2	7.4	15.8
4	4	12.1	23.9
6	6	17.9	27.8
11b (3)	2	7.9	15.9
11b (1)	4	13.8	24.2
11b (2)	6	17.1	26.9
10a	2	7.6	14.2
10b	4	10.2	19.2
11a	6	17.7	24.6

53/4/8

BARLEY - WOBURN STACKYARD 1953

For history, details of treatments etc. see "Results of the Field Experiments 1939-47" Vol. I, Section A/7 with the exception that in 1952 it was decided to divide each plot into two, and sow one half with a Winter variety, and the other as before with a Spring variety.

Cultivations, etc.: Ploughed: Aug 26, 1952. 'Winter' seed drilled, at 3 bushels per acre: Oct 9. 'Spring' seed drilled at 3 bushels per acre: Mar 5, 1953. Nitrochalk applied: Mar 13. Sprayed with DNOC 6 lb in 80 gallons per acre: May 22. Harvested: Aug 17. Varieties: Winter - Pioneer, Spring - Plumage Archer.

Summary of Results

Plot	Nitrochalk dressing cwt per acre	Grain: cwt per acre		Straw: cwt per acre	
		Winter sown	Spring sown	Winter sown	Spring sown
3	2	16.3	8.4	12.8	7.6
7	4	9.0	15.9	10.6	13.6
1	6	8.3	16.6	11.8	14.0
9	2	19.8	8.9	14.8	9.0
4	4	17.6	18.9	21.7	16.2
6	6	25.9	25.2	26.9	22.2
11b(1)	2	20.3	14.2	21.0	12.1
11b(2)	4	26.4	19.7	23.4	17.9
11b(3)	6	31.1	24.4	34.5	22.0
11a	2	15.3	12.1	14.1	9.1
10a	4	6.6	8.7	10.1	9.1
10b	6	15.1	11.6	16.1	9.8