


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1953

[Full Table of Content](#)



### 53/CA/5 Wheat - Control Wireworm - Rothamsted

#### Rothamsted Research

Rothamsted Research (1954) *53/CA/5 Wheat - Control Wireworm - Rothamsted* ; Yields Of The Field Experiments 1953, pp 75 - 75 - DOI: <https://doi.org/10.23637/ERADOC-1-173>

53/Ca/5

WHEAT

The residual effect of insecticides on the control of wireworm - Geescroft 1953.

System of replication: 3 randomized blocks of 8 plots each.

Area of each plot: 0.0289 acre. Area harvested: 0.0178 acre.

Treatments (applied autumn 1951 to previous wheat):

None (triplicate plots)							(O)
Gammexane seed dressing, 2 oz per bushel							(S)
Gammexane combine drilled with seed at	56 lb per acre,	3.5% dust					(G)
Aldrin " " " "	200 lb per acre,	1.78% "					(A)
Chlordane " " " "	100 lb per acre,	5% "					(C)
D. D. T. " " " "	150 lb per acre,	5% "					(D)

Basal dressing: 3 cwt sulphate of ammonia per acre.

Cultivations, etc.: Ploughed: Sept 8, 1952. Seed drilled at 2 1/4 bushels per acre: Oct 27. Sulphate of ammonia applied: Apr 25, 1953. Combine harvested: Sept 26. Variety: Nord Desprez. Previous crop: Wheat.

Standard error per plot:

Grain: 2.76 cwt per acre or 10.4% (16 d.f.)

Note: Wireworm counts were made and are available.

For details of original experiment see "Results of the Field Experiments 1952", Section Ca/4.

Summary of Results

	O	S	G	A	C	D	Mean
Grain: cwt per acre							
Mean (±1.60)	21.6 <sup>(1)</sup>	22.9	30.2	34.6	32.9	26.7	26.5
Increase (±1.84)		1.3	8.6	13.0	11.3	5.1	

Standard error (1) ±0.92