

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1953

[Full Table of Content](#)



---

### 53/CA/6 Wheat - Late Application of Nitrogen - Rothamsted

#### Rothamsted Research

Rothamsted Research (1954) *53/CA/6 Wheat - Late Application of Nitrogen - Rothamsted* ; Yields Of The Field Experiments 1953, pp 76 - 76 - DOI: <https://doi.org/10.23637/ERADOC-1-173>

53/Ca/6

WHEAT

Late application of nitrogen - Highfield 1953.

System of replication: 4 randomized blocks of 3 plots each.

Area of each plot: 0.0204 acre.

Treatments:

Nitrogen: None; 0.23; 0.46 cwt per acre applied as nitrochalk top dressing.

Basal dressing: 2 cwt sulphate of ammonia per acre.

Cultivations, etc.: Ploughed: Aug 23, 1952. Seed drilled at 3 bushels per acre: Oct 30. Sulphate of ammonia applied: Apr 29, 1953. Nitrochalk applied: June 26. Harvested: Aug 19. Variety: Nord Desprez. Previous crop: Wheat.

Standard error per plot:

Grain: 1.12 cwt per acre or 3.5% (6 d.f.)

Summary of Results

	Nitrogen (as nitrochalk top dressing): cwt per acre			Mean
	None	0.23	0.46	
Yield: cwt per acre				
Grain ( $\pm 0.56$ )	31.8	32.2	31.8	31.9
Straw <sup>x</sup>	44.6	43.3	43.3	43.7
Crude protein: cwt per acre				
Grain	3.79	3.91	3.88	
Increase		+0.12	+0.09	
Straw	1.78	1.73	1.82	
Increase		-0.05	+0.04	
Percentage uptake of added nitrogen				
Grain		+8	+3	
Straw		-3	+1	

<sup>x</sup>Corrected to 85% dry matter. Mean dry matter %: 83.6.