


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1953

[Full Table of Content](#)



53/CE/3 Potatoes - Control of Blight - Rothamsted

Rothamsted Research

Rothamsted Research (1954) *53/CE/3 Potatoes - Control of Blight - Rothamsted* ; Yields Of The Field Experiments 1953, pp 96 - 97 - DOI: <https://doi.org/10.23637/ERADOC-1-173>

53/0e/3.1

POTATOES

Control of Blight - Deacons Field 1953.

System of replication: 4 x 4 Latin Square, plots being split into 2 for determination of tractor damage.

Area of each sub plot: 0.0175 acre.

Treatments:

Whole plots: No spray; Copper fungicide 5 lb in 40 gallons per acre sprayed twice; 100 gallons Sulphuric acid, 15% B.O.V. sprayed to destroy haulms; Copper fungicide and sulphuric acid sprayed as above. The tractor used for spraying was driven over all the plots on each occasion.

Sub plots: The 4 inner rows damaged by three passages of the tractor were compared with the 4 outer and undamaged rows.

Basal dressing: 15 cwt compound granular fertilizer (7% N; 7% P₂O₅; 10 $\frac{1}{2}$ % K₂O) per acre.

Cultivations, etc.: Ploughed: Sept 15, 1952. Cultivated twice: Mar 17, 1953. Ridged: Mar 26. Potatoes planted: Apr 16. Earthed up: June 26. Sprayed appropriate plots with copper fungicide: Aug 1 and again Aug 13. Sprayed appropriate plots with sulphuric acid: Sept 14. Lifted: Oct 6. Variety: Majestic. Previous crop: Barley.

Standard errors per plot: Total tubers.

Whole plot: 0.361 tons per acre or 2.7% (6 d.f.)

Sub plot: 0.454 tons per acre or 3.4% (12 d.f.)

Blight counts were made and are available. The mean level of infection was only 0.3%.

53/Ce/3.2

Summary of Results

	Spray				Mean
	None	Copper fungicide	Sulphuric Acid	Copper fungicide and sulphuric Acid	
Total tubers: tons per acre (± 0.242) [†]					
Undamaged rows	13.45	13.87	13.28	14.66	13.82
Damaged rows	12.49	13.67	12.71	13.52	13.10
Mean (± 0.181)	12.97	13.77	13.00	14.09	13.46
Difference (± 0.321)	-0.96	-0.20	-0.57	-1.14	-0.72 (± 0.161)

Percentage Ware

Undamaged rows	91.4	88.4	91.8	89.7	90.3
Damaged rows	90.6	89.6	90.7	92.5	90.9
Mean	91.0	89.0	91.2	91.1	90.6
Difference	-0.8	+1.2	-1.1	+2.8	+0.6

[†] for use in comparisons other than vertical.