


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1954

[Full Table of Content](#)



---

## Long-term Experiments

### Rothamsted Research

Rothamsted Research (1955) *Long-term Experiments* ; Yields Of The Field Experiments 1954, pp 13 - 62 - DOI: <https://doi.org/10.23637/ERADOC-1-184>

54/Ba/1.1

### THREE COURSE ROTATION EXPERIMENT

3rd year of revised scheme

For details of treatments and rotation see "Results of the Field Experiments 1952", Section Ba/1.

Area of each plot: Potatoes (sub-plot) - 0.0092 acre;  
Barley - 0.0200 acre; Sugar beet - 0.0204 acre.

#### Cultivations, etc.:

Potatoes. Straw applied, ploughed all plots: Dec 28, 1953.  
Fertilizers applied: Apr 8, 1954. Ridged: Apr 9. Potatoes planted: Apr 12. Earthed up: July 6. Sprayed (low volume) with copper fungicide at 5 lb powder per acre: July 27. Sprayed (low volume) with copper fungicide at  $\frac{1}{2}$  gallon paste per acre: Aug 23. Sprayed with 15% sulphuric acid: Sept 28. Lifted: Oct 2. Variety: Majestic.

Barley. Straw applied, ploughed all plots: Dec 28, 1953.  
Ground chalk applied at 22 cwt per acre: Jan 5, 1954. Fertilizers applied, seed drilled at 3 bushels per acre: Mar 15. Harvested: Aug 26. Variety: Plumage Archer.

Sugar beet. Straw applied, ploughed all plots: Dec 28, 1953.  
Fertilizers applied: Mar 29, 1954. Seed drilled at 18 lb per acre: Apr 5. Sprayed (at low volume) with D.D.T. emulsion at  $\frac{1}{2}$  lb per acre: May 12. Singled: June 1. Sprayed again as above: June 24. Lifted: Nov 16. Variety: Klein E.

#### Treatment symbols

Ar Complete artificials only  
St1 Straw ploughed in in autumn, artificials applied in spring  
St2 Straw ploughed in in autumn, artificials applied half in autumn, half in spring  
Ad Adco ploughed in in autumn with supplementary artificials  
St  $5\frac{1}{3}$  cwt cut straw in autumn  
Nitrogen dressing: 0.2; 0.4; 0.6 cwt N as sulphate of ammonia  
K<sub>3</sub> Muriate of potash equivalent to K<sub>2</sub>O in straw  
K 0.5 cwt K<sub>2</sub>O as muriate of potash.



Summary of Results

Potatoes

Treatments applied	1950	1953		1954		1952 & 1954		0	0.4N	St+ 0.2N	St+ 0.6N	Ks	Ks + 0.4N
		-	K	-	K	-	K						
		6.15	6.24	4.92	5.98								
	Ar	5.19	6.72	4.18	5.62								
	St1 St2	6.43	5.76	5.19	5.38	5.91	6.24	6.63	5.52	7.30	7.20	4.66	7.25
	St1 St2	4.75	5.57	4.37	4.47								
		7.54	8.55	7.68	7.44								
		5.57	7.15	6.00	5.67								
	Ad	4.23	7.30	5.47	6.82	7.68	5.28	4.95	6.10				
		5.38	5.52										
		6.24	7.30										

Total tubers: tons per acre

54/Ba/1.2







54/Ba/1.4

Barley

Treatments applied		1953	0	0.4N	St + 0.2N	St + 0.6N	K <sub>S</sub>	K <sub>S</sub> + 0.4N
1950	1951	1952 & 1954	Grain: cwt per acre					
	Ar	0		30.1				
		0.4N	29.9					
Ar		0		30.8				
		0.4N	30.5					
	St1 St2	0	24.3		30.5			28.3
		0.4N	34.7		31.4		34.3	
St1 St2		0		29.9				
		0.4N	30.7					
		St+ 0.2N		23.5				
		St+ 0.6N		33.0				
		K <sub>S</sub>		28.5				
		K <sub>S</sub> +0.4N		34.3				
	Ad	0		28.9		27.0		32.6
		0.4N	32.0					
Ad		St+ 0.6N	29.7					
		K <sub>S</sub> +0.4N	35.3					
			Straw: cwt per acre					
	Ar	0		33.5				
		0.4N	41.8					
Ar		0		35.1				
		0.4N	40.2					
	St1 St2	0	27.6		42.0			32.8
		0.4N	42.7		43.6		39.6	
St1 St2		0		43.8				
		0.4N	34.3					
		St+ 0.2N		26.5				
		St+ 0.6N		44.6				
		K <sub>S</sub>		31.8				
		K <sub>S</sub> +0.4N		44.1				
	Ad	0		33.5		30.9		34.9
		0.4N	43.2					
Ad		St+ 0.6N	37.7					
		K <sub>S</sub> +0.4N	43.9					



54/Ba/1.5

Treatments applied 1953			Sugar Beet					
1950	1951	1952 & 1954	0	0.4N	St + 0.2N	St + 0.6N	K <sub>S</sub>	K <sub>S</sub> + 0.4N
			Roots (washed): tons per acre					
	Ar	0		7.90				
		0.4N	8.90					
Ar		0		6.46				
		0.4N	9.42					
	St1 St2	0		7.40		8.21		8.23
		0.4N	9.45		9.32		10.30	
St1 St2		0		8.29				
		0.4N	9.93					
		St+ 0.2N		8.75				
		St+ 0.6N	9.78					
		K <sub>S</sub>		8.29				
		K <sub>S</sub> +0.4N	10.93					
	Ad	0		7.77		6.66		8.07
Ad		0.4N	8.95					
		St+ 0.6N	9.65					
		K <sub>S</sub> +0.4N	8.34					

Treatments applied 1953			Sugar Percentage					
1950	1951	1952 & 1954	0	0.4N	St + 0.2N	St + 0.6N	K <sub>S</sub>	K <sub>S</sub> + 0.4N
	Ar	0		17.00				
		0.4N	17.08					
Ar		0		17.51				
		0.4N	17.25					
	St1 St2	0		16.82		17.48		17.08
		0.4N	17.19		17.22		17.34	
St1 St2		0		17.14				
		0.4N	17.08					
		St+ 0.2N		17.19				
		St+ 0.6N	17.16					
		K <sub>S</sub>		16.88				
		K <sub>S</sub> +0.4N	17.16					
	Ad	0		16.85		17.14		17.44
Ad		0.4N	16.93					
		St+ 0.6N	17.77					
		K <sub>S</sub> +0.4N	17.14					



54/Ba/1.6

Sugar Beet

Treatments applied			1953	0	0.4N	St + 0.2N	St + 0.6N	K <sub>S</sub>	K <sub>S</sub> + 0.4N
1950	1951	1952 & 1954							
Total sugar: cwt per acre									
	Ar	0		26.8					
		0.4N	30.4						
Ar		0		22.6					
		0.4N	32.5						
	St1 St2	0		24.9		28.7			28.1
		0.4N	32.5		32.1		35.7		
St1 St2		0		28.4					
		0.4N	33.9						
		St + 0.2N		30.1					
		St + 0.6N		33.6					
		K <sub>S</sub>		28.0					
		K <sub>S</sub> + 0.4N		37.5					
	Ad	0		26.2		22.8			28.2
Ad		0.4N	30.3						
		St + 0.6N		34.3					
		K <sub>S</sub> + 0.4N		28.6					
Tops: tons per acre									
	Ar	0		8.28					
		0.4N	8.90						
Ar		0		6.07					
		0.4N	10.14						
	St1 St2	0		6.70		7.14			7.46
		0.4N	7.51		7.81		8.71		
St1 St2		0		9.62					
		0.4N	10.61						
		St + 0.2N		8.47					
		St + 0.6N		10.72					
		K <sub>S</sub>		7.22					
		K <sub>S</sub> + 0.4N		13.33					
	Ad	0		6.35		9.47			7.59
Ad		0.4N	9.76						
		St + 0.6N		8.47					
		K <sub>S</sub> + 0.4N		6.72					



54/Ba/1.7

Sugar Beet

Treatments applied 1953			0	0.4N	St + 0.2N	St + 0.6N	K <sub>S</sub>	K <sub>S</sub> + 0.4N
1950	1951	1952 & 1954						
Plant number: thousands per acre								
	Ar	0		27.1				
		0.4N	27.4					
Ar		0		27.0				
		0.4N	27.0					
	St1 St2	0		27.2		27.9		27.0
		0.4N	25.8		26.8		27.6	
St1 St2		0		27.7				
		0.4N	26.7					
		St + 0.2N		26.6				
		St + 0.6N	27.4					
		K <sub>S</sub>		27.1				
		K <sub>S</sub> +0.4N	27.5					
	Ad	0		26.3		27.6		26.6
Ad		0.4N	26.5					
		St + 0.6N	26.2					
		K <sub>S</sub> +0.4N	27.0					

Noxious nitrogen:mg per 100g.

	Ar	0		20.0				
		0.4N	15.0					
Ar		0		15.0				
		0.4N	17.5					
	St1 St2	0		15.0		20.0		15.0
		0.4N	15.0		15.0		15.0	
St1 St2		0		20.0				
		0.4N	15.0					
		St + 0.2N		20.0				
		St + 0.6N	25.0					
		K <sub>S</sub>		15.0				
		K <sub>S</sub> +0.4N	20.0					
	Ad	0		15.0		15.0		15.0
Ad		0.4N	20.0					
		St + 0.6N	15.0					
		K <sub>S</sub> +0.4N	15.0					



54/Ba/2.1

FOUR COURSE ROTATION EXPERIMENT

The 25th year

Direct and residual effects of organic manures and phosphatic fertilizers - Hoosfield 1954.

For details of treatments and rotation see "Results of the Field Experiments 1939-47" Vol.I, Section Ba/3.

Area of each plot: Potatoes (whole plot): 0.0228 acre. Barley, ryegrass and wheat: 0.0244 acre.

Manures (cwt per acre) applied 1953-54

Treatment	Organic manures and phosphates				Supplementary fertilizers		
	Organic matter	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	N as Sulph. of amm.	P <sub>2</sub> O <sub>5</sub> as Super	K <sub>2</sub> O as Mur. of potash
Dung (a)	50	1.236	0.534	2.331	0.564	0.666	0.669
Dung (b)	50	1.189	0.514	2.406	0.611	0.686	0.594
Adco	50	1.302	0.754	0.822	0.498	0.446	2.178
Straw (a)	167	0.533	0.100	0.550	1.267	1.100	2.450
Straw (b)	167	0.833	0.250	1.250	0.967	0.950	1.750
Super			1.2		0.36*		0.6*
Rock phosphate			1.2		0.36*		0.6*

Dung and straw: (a) applied to wheat and ryegrass; (b) to barley and potatoes.

\*Yearly dressings totalling to the standard rates in the 5 year period.

Cultivations, etc.:

Potatoes.

Ploughed: Sept 2, 1953. Dung and adco applied: Nov 26. Straw, first dressing of fertilizers to straw plot and supplementary fertilizers to dung and adco plots applied, all plots ploughed: Nov 30. Second dressing of fertilizers to straw plot applied: Jan 18, 1954. Ridged, spring fertilizers including third dressing to straw plot and sulphate of ammonia to half plots applied, potatoes planted: Apr 20. Earthed up: July 21. Sprayed with copper fungicide,  $\frac{1}{2}$  gallon paste in 10 gals. water per acre: July 27. Sprayed with copper fungicide, 5 lb per acre: Aug 13. Sprayed with sulphuric acid, 15% B.O.V. 100 gallons per acre: Sept 28. Lifted: Oct 9. Variety: Majestic.

Barley.

Dung and adco applied: Nov 26, 1953. Straw, first dressing of fertilizers to straw plot and supplementary fertilizers to dung and adco plots applied, all plots ploughed: Dec 1. Ground chalk applied at 21 cwt per acre: Dec 5. Second dressing of fertilizer to straw plot: Jan 18, 1954. Spring fertilizers



54/Ba/2.2

including third dressing to straw plot applied: Mar 12. Seed drilled at 3 bushels per acre: Mar 15. Harvested: Aug 27.

Variety: Plumage Archer.

Ryegrass.

Dung and adco with supplementary fertilizers, straw with first dressing of fertilizers applied, all plots ploughed: Sept 23, 1953.

Autumn fertilizers applied: Oct 8. Seed sown at 112 lb per acre:

Oct 9. Second dressing of fertilizers to straw plot applied:

Jan 18, 1954. Sulphate of ammonia, and third dressing of fertilizers to straw plot applied: Apr 29. Harvested: June 29.

Variety: Western Wolths.

Wheat.

Dung and adco with supplementary fertilizers, straw with first dressing of fertilizers applied, all plots ploughed: Sept 23, 1953.

Autumn fertilizers applied: Oct 9. Seed drilled at 3 bushels per acre: Oct 22. Second dressing of fertilizers to straw plot

applied: Jan 18, 1954. Sulphate of ammonia and third dressing of fertilizers to straw plot applied: May 1. Harvested: Sept 6.

Variety: Squareheads Master 13/4.



Summary of Results

Manure +	Year of Cycle	Potatoes				Percentage Ware		Response to N	Barley Grain Straw # cwt per acre	Ryegrass Dry matter cwt per acre	Wheat Grain # Straw * cwt per acre
		Total tubers, tons per acre	Additional N Without	With	Mean	Additional N Without	With				
Dung	I	5.02	5.81	5.42	+0.79	78.6	74.3	76.4	32.2	36.9	28.0
	II	4.77	5.46	5.12	+0.69	86.0	78.1	82.0	19.2	13.1	17.3
	III	4.24	4.32	4.28	+0.08	73.8	77.4	75.6	24.8	21.0	16.6
	IV	3.45	4.80	4.12	+1.35	78.3	73.3	75.8	15.0	13.0	16.5
	V	3.24	4.06	3.65	+0.82	76.9	81.7	79.3	17.0	17.3	18.9
Adco (Straw compost)	I	4.17	5.07	4.62	+0.90	76.8	83.6	80.2	28.9	32.2	27.0
	II	2.78	3.51	3.14	+0.73	69.5	72.0	70.8	25.7	20.7	15.4
	III	2.29	2.78	2.54	+0.49	69.5	78.5	74.0	22.1	17.4	16.8
	IV	2.53	2.36	2.44	-0.17	72.9	74.7	73.8	20.8	13.1	18.0
	V	3.21	3.24	3.22	+0.03	69.7	77.6	73.6	20.6	16.7	18.3
Straw	I	4.58	5.48	5.03	+0.90	76.7	76.2	76.4	26.0	46.6	29.4
	II	2.86	4.06	3.46	+1.20	74.0	83.4	78.7	20.0	11.9	13.2
	III	3.73	4.82	4.28	+1.09	74.1	85.5	79.8	23.5	20.0	17.8
	IV	3.65	4.02	3.84	+0.37	78.8	77.4	78.1	20.5	13.6	15.3
	V	5.46	5.40	5.43	-0.06	77.0	82.5	79.8	20.0	17.1	15.5
Super-phosphate	I	4.36	5.56	4.96	+1.20	78.1	76.1	77.1	23.0	25.7	20.2
	II	3.62	3.53	3.58	-0.09	78.2	74.3	76.2	24.8	28.7	21.8
	III	4.24	2.70	3.47	-1.54	72.6	74.8	73.7	25.4	28.4	22.3
	IV	3.54	4.11	3.82	+0.57	80.5	79.9	80.2	23.8	30.7	18.2
	V	3.43	3.99	3.71	+0.56	74.0	82.1	78.0	24.8	26.4	21.8
Rock phosphate	I	1.70	2.18	1.94	+0.48	62.4	67.6	65.0	25.9	34.0	24.3
	II	2.14	2.66	2.40	+0.52	76.1	73.7	74.9	23.2	24.9	22.3
	III	2.41	2.51	2.46	+0.10	79.5	70.0	74.8	25.9	27.2	19.6
	IV	1.92	1.95	1.94	+0.03	70.1	69.5	69.8	26.1	25.7	20.3
	V	2.33	1.83	2.08	-0.50	76.3	65.8	71.0	24.9	23.4	22.8
Mean Dry Matter % <span style="float: right;">54/Ba/2.3</span>											
* At 65% Dry Matter <span style="float: right;">82.5</span>											

+ Note: All manures are supplemented by fertilizers as shown in table on page 54/Ba/2.1



54/Ba/3.1

SIX COURSE ROTATION EXPERIMENT

The 25th year

Seasonal effects of fertilizers - Rothamsted Long Hoos IV and Woburn Stackyard 1954.

For details of treatments, rotation etc., see "Results of the Field Experiments 1939-47" Vol.I, Section Ba/4.

Area of each plot: Rothamsted - 0.0250 acre. Woburn - 0.0266 acre.

Cultivations, etc.:

Rothamsted

Sugar beet.

Ploughed: Aug 21, 1953, Oct 20 and Dec 1. Fertilizers applied: Mar 26, 1954. Seed drilled at 18 lb per acre: Mar 30. Singled: May 31. Lifted: Nov 17. Variety: Klein E.

Barley.

Ploughed: Nov 27, 1953. Ground chalk applied at 22 cwt per acre: Dec 4. Fertilizers applied, seed drilled at 3 bushels per acre: Mar 15, 1954. Harvested: Aug 25. Variety: Plumage Archer.

Clover.

Seed undersown in barley at 40 lb per acre: Apr 23, 1953. Autumn fertilizers applied: Nov 2. Sulphate of ammonia applied: Apr 30, 1954. Cut: July 13. Variety: Late flowering Montgomery Red.

Wheat.

Ploughed: July 10, 1953 and again Sept 17. Autumn fertilizers applied: Oct 9. Seed drilled at 3 bushels per acre: Oct 17. Sulphate of ammonia applied: Apr 30, 1954. Harvested: Aug 26. Variety: Yeoman.

Potatoes.

Ploughed: Aug 29, 1953, Oct 20 and Nov 30. Ridged: Apr 13, 1954. Fertilizers applied, potatoes planted: Apr 14. Earthed up: July 6. Sprayed (low volume) with copper fungicide paste  $\frac{1}{2}$  gallon per acre: July 27 and again Aug 18. Sprayed with sulphuric acid 15% B.O.V: Sept 28. Lifted: Oct 2. Variety: Majestic.

Rye.

Ploughed: Sept 30, 1953. Ground chalk applied at 22 cwt per acre: Oct 3. Autumn fertilizers applied: Oct 10. Seed drilled at 3 bushels per acre: Oct 17. Sulphate of ammonia applied: Apr 30, 1954. Harvested: Aug 25. Variety: King II.

Note:- Clover: 6 plots were carted off in error when green. No summary of results can be shown.



54/Ba/3.2

Woburn

Sugar beet.

Ploughed: Aug 20, 1953, Nov 3 and Jan 20, 1954. Fertilizers applied: Mar 29. Seed drilled at 18 lb per acre: Mar 30. Dusted with D.D.T; May 12. Singled: May 31. Sprayed with parathion  $\frac{1}{2}$  pint per acre: June 21. Lifted: Nov 2. Variety: Klein E. Previous crop: Rye. N.B. The crop was attacked by flea beetle and mangold fly.

Barley.

Ploughed: Nov 4, 1953. 2nd ploughing: Jan 20, 1954. Fertilizers applied: Mar 11. Seed drilled at  $2\frac{1}{2}$  bushels per acre: Mar 17. Harvested: Aug 28. Variety: Plumage Archer. Previous crop: Sugar beet.

Clover.

Seed undersown in barley at 40 lb per acre: Mar 26, 1953. Autumn fertilizers applied: Oct 23. Sulphate of ammonia applied: Apr 27, 1954. Cut: June 30. Variety: Montgomery Red. Previous crop: Barley. N.B. The crop was damaged by Sclerotinia and was very weedy.

Wheat.

Ploughed: July 3, 1953. 2nd ploughing: Sept 16. Autumn fertilizers applied: Oct 22. Drilled at  $2\frac{1}{2}$  bushels per acre: Oct 23. Sprayed with D.N.O.C. at 6 lb per acre in 80 gallons: Apr 27. Sulphate of ammonia applied: Apr 29. Harvested: Sept 7. Variety: Squareheads Master  $13/4$ . Previous crop: Clover. N.B. The crop was very weedy.

Potatoes.

Ploughed: Sept 15, 1953, Nov 3 and Jan 20, 1954. Ridged, fertilizers applied, seed planted: Apr 18. Sprayed with copper fungicide: July 30. Sprayed with 15% sulphuric acid: Sept 23. Lifted: Oct 8. Variety: Majestic. Previous crop: Wheat.

Rye.

Ploughed: Oct 1, 1953. Autumn fertilizer applied, drilled seed at  $2\frac{1}{2}$  bushels per acre: Oct 22. Sulphate of ammonia applied: Apr 29. Harvested: Aug 28. Variety: King II. Previous crop: Potatoes.



54/Ba/3.3

Summary of Results

Mean yields per acre and responses in yield per cwt of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O

	Rothamsted	Woburn	Rothamsted	Woburn
Sugar Beet, roots (washed): tons per acre			Barley, grain: cwt per acre	
Mean	8.55	4.57	31.7	29.1
Response to: N	+2.88	+1.14	+4.6	+25.6
P	+3.05	+4.46	+0.4	-1.9
K	-0.08	-0.07	-1.2	-1.0
Sugar Beet, sugar percentage			Barley, straw: cwt per acre	
Mean	16.30	16.29	36.0	29.9
Response to: N	+0.22	+0.56	+12.8	+40.3
P	-1.21	+1.03	+1.1	-7.1
K	+0.28	+0.58	-1.9	+2.8
Sugar Beet, total sugar: cwt per acre			Clover, hay: dry matter cwt per acre	
Mean	27.9	14.9		37.6
Response to: N	+9.9	+4.2		-29.0
P	+8.1	+15.7		+11.3
K	+0.2	+0.2		+8.6
Sugar Beet, tops: tons per acre			Wheat, grain: cwt per acre	
Mean	10.93	4.88	26.5	14.8
Response to: N	+5.88	+0.69	+8.0	+19.3
P	+1.93	+5.21	-10.7	-9.6
K	+0.02	-0.55	+5.2	+10.1
Sugar Beet, plant number: thousands per acre			Wheat, straw: cwt per acre	
Mean	26.0		48.2	18.6
Response to: N	-1.1	†	+22.1	+22.7
P	+3.1		-2.1	-9.8
K	+1.7		-0.8	+12.7
Potatoes, total tubers: tons per acre			Rye, grain: cwt per acre	
Mean	6.08	4.21	32.8	26.4
Response to: N	+3.07	+4.49	+32.2	+24.6
P	+0.57	+1.39	-3.7	+3.3
K	+1.32	+0.30	+0.1	-1.1
Potatoes, percentage ware			Rye, straw: cwt per acre	
Mean	78.9	80.5	40.2	30.9
Response to: N	-10.7	+10.8	+23.6	+23.5
P	-10.7	+6.6	-3.9	+8.9
K	+6.5	+12.7	+2.0	-0.5

\* corrected to 85% Dry Matter. Mean Dry Matter %: 83.6 † not recorded.



54/Bb/1.1

## DEEP CULTIVATION ROTATION EXPERIMENT

The 11th year

Deep ploughing, fertilizers and dung - Long Hoos I and II 1954.

For details of rotation and treatments etc. see "Results of the Field Experiments 1939-47" Vol. I, Section Bc/1.

Area of each plot: 0.0312 acre. Area harvested: Sugar beet (half plot), 0.0119 acre; barley, wheat, spring oats, 0.0265 acre; ley, 0.0275 acre; potatoes (half plot), 0.0107 acre.

Cultivations, etc.:

Sugar beet (Series 4)

Dung and fertilizers for ploughing in "deep" applied: Sept 25, 1953. "Deep" plots ploughed: Sept 29. Dung and fertilizers for ploughing in "shallow" applied, "shallow" plots ploughed: Sept 30. Fertilizers for surface application applied: Mar 29, 1954. Seed drilled at 18 lb per acre: Mar 30. Sprayed with DDT oil emulsion (15% DDT) at 3 pints per acre: May 12 and again June 24. Singled: June 3. Lifted: Nov 25. Variety: Klein E.

Barley (Series 5)

Ploughed: Dec 16, 1953. Ground chalk at 21 cwt per acre applied: Dec 21. Basic slag and sulphate of ammonia applied: Mar 12, 1954. Seed drilled at  $2\frac{3}{4}$  bushels per acre: Mar 13. Harvested: Aug 27. Variety: Plumage Archer.

Ley (Series 1)

Seeds undersown in barley: Apr 23, 1953. Cut: July 23, 1954. Seeds mixture (per acre): 18 lb S.24 perennial ryegrass, 8 lb Montgomery red clover, 2 lb American Alsike clover.

Wheat (Series 6)

"Deep" and "shallow" plots ploughed: Sept 30, 1953. Seed drilled at 3 bushels per acre: Oct 17. Sulphate of ammonia applied: Apr 30, 1954. Sprayed with MCPA, 2 pints per acre at medium volume: May 27. Harvested: Aug 27. Variety: Yeoman.

Potatoes (Series 2)

Dung and fertilizers for ploughing in "deep" applied: Sept 24, 1953. "Deep" plots ploughed: Sept 25. Dung and fertilizers for ploughing in "shallow" applied, "shallow" plots ploughed: Sept 26. Ridged: Apr 13, 1954. Fertilizers for surface application applied, potatoes planted: Apr 14. Earthed up: July 5. Sprayed with copper fungicide,  $\frac{1}{2}$  gallon paste in 10 gallons water per acre: July 27 and again Aug 18. Sprayed with sulphuric acid, 15% BOV: Sept 28. Lifted: Oct 25. Variety: Majestic.



54/Bb/1.2

Spring oats (Series 3)

Ploughed: Sept 28, 1953. Ground chalk at 21 cwt per acre applied:  
Dec 4. Sulphate of ammonia applied: Mar 12, 1954. Seed drilled  
at 4 bushels per acre: Mar 13. Harvested: Aug 27. Variety:  
Star.

Standard errors per plot:

Sugar beet, Total sugar.	Whole plot: 3.08 cwt per acre or 8.2% (4 d.f.)
	Sub plot: 2.88 cwt per acre or 7.7% (7 d.f.)
Tops.	Whole plot: 0.658 tons per acre or 6.0% (4 d.f.)
	Sub plot: 2.50 tons per acre or 22.6% (7 d.f.)
Barley, Grain:	1.82 cwt per acre or 5.5% (4 d.f.)
Ley, Hay:	5.31 cwt per acre or 6.4% (4 d.f.)
Wheat, Grain:	2.34 cwt per acre or 6.1% (4 d.f.)
Potatoes, Total tubers,	Whole plot: 0.919 tons per acre or 10.6% (4 d.f.)
	Sub plot: 0.991 tons per acre or 11.4% (7 d.f.)
Spring Oats, Grain:	3.82 cwt per acre or 11.0% (4 d.f.)



54/Bb/1.3

Summary of Results

Series 4: Sugar Beet

Responses to treatments

Response to	Mean	Ploughing		Dung		Phosphate		Potash	
		Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.

Roots (washed): Mean yield 11.16 tons per acre

Ploughing									
deep-shallow	-0.31	-	-	+0.05	-0.67	-1.05	+0.43	-0.03	-0.59
Dung	+1.48	+1.84	+1.12	-	-	+0.76	+2.20	+2.19	+0.77
Phosphate	+0.68	-0.06	+1.42	-0.04	+1.40	-	-	+0.02	+1.34
Potash	+0.67	+0.95	+0.39	+1.38	-0.04	+0.01	+1.33	-	-

Sugar Percentage: Mean 16.85

Ploughing									
deep-shallow	-0.25	-	-	+0.14	-0.64	-0.29	-0.21	-0.31	-0.19
Dung	-0.29	+0.10	-0.68	-	-	-0.24	-0.34	-0.04	-0.54
Phosphate	+0.04	0.00	+0.08	+0.09	-0.01	-	-	+0.35	-0.27
Potash	-0.02	-0.08	+0.04	+0.23	-0.27	+0.29	-0.33	-	-

Total Sugar: Mean yield 37.6 cwt per acre

(±1.54)

(±2.18)

Ploughing									
deep-shallow	-1.6	-	-	+0.6	-3.8	-4.1	+0.9	-0.8	-2.4
Dung	+4.4	+6.6	+2.2	-	-	+2.1	+6.7	+7.4	+1.4
Phosphate	+2.3	-0.2	+4.8	0.0	+4.6	-	-	+0.7	+3.9
Potash	+2.2	+3.0	+1.4	+5.2	-0.8	+0.6	+3.8	-	-

Tops: Mean yield 11.03 tons per acre

(±0.329)

(±0.465)

Ploughing									
deep-shallow	+0.92	-	-	+0.57	+1.27	+1.17	+0.67	+0.05	+1.79
Dung	+1.18	+0.83	+1.53	-	-	+0.99	+1.37	-0.05	+2.41
Phosphate	-0.27	-0.02	-0.52	-0.46	-0.08	-	-	-0.72	+0.18
Potash	-0.19	-1.06	+0.68	-1.42	+1.04	-0.64	+0.26	-	-

Plant Number: Mean 27.1 thousands per acre

Ploughing									
deep-shallow	-0.3	-	-	0.0	-0.6	0.0	-0.6	+0.3	-0.9
Dung	-0.4	-0.1	-0.7	-	-	-0.1	-0.7	-0.4	-0.4
Phosphate	+0.2	+0.5	-0.1	+0.5	-0.1	-	-	+0.2	+0.2
Potash	-0.1	+0.5	-0.7	-0.1	-0.1	-0.1	-0.1	-	-

Noxious Nitrogen: Mean 33.1 mg. per 100 g.

Ploughing									
deep-shallow	+5.0	-	-	+5.0	+5.0	+6.2	+3.8	+5.6	+4.4
Dung	0.0	0.0	0.0	-	-	-2.5	+2.5	+1.9	-1.9
Phosphate	+1.2	+2.4	0.0	-1.3	+3.7	-	-	-0.7	+3.1
Potash	-5.6	-5.0	-6.2	-3.7	-7.5	-7.5	-3.7	-	-



54/Bb/1.4

Series 4: Sugar Beet

	Phosphate			Potash			Mean
	None	Ploughed in	In seed bed	None	Ploughed in	In seed bed	
Roots (washed): tons per acre							
Shallow	11.34	11.43	11.12	10.84	11.37	12.20	11.31
Deep	10.29	12.39	11.04	10.81	11.20	11.21	11.00
No dung	10.44	10.70	10.09	9.72	11.11	11.10	10.42
Dung	11.20	13.13	12.07	11.92	11.45	12.31	11.90
Mean	10.82	11.91	11.08	10.82	11.23	11.71	11.16
Sugar Percentage							
Shallow	16.97	17.03	16.92	17.01	17.02	16.84	16.97
Deep	16.68	16.65	16.87	16.70	16.68	16.80	16.72
No dung	16.95	17.10	16.98	16.38	17.13	17.08	16.99
Dung	16.71	16.58	16.81	16.84	16.57	16.56	16.70
Mean	16.83	16.84	16.90	16.86	16.85	16.82	16.85
Total Sugar: cwt per acre							
	(a)	(b) and (c)		(a)	(b) and (c)		
Shallow	38.4	39.0	37.5	36.9	38.7	41.0	38.3
Deep	34.4	41.1	37.2	36.1	37.2	37.7	36.8
No dung	35.4	36.6	34.1	32.8	38.0	37.8	35.4
Dung	37.5	43.5	40.6	40.2	37.9	40.8	39.8
Mean	36.4	40.0	37.4	36.5	38.0	39.3	37.6
Tops: tons per acre							
	(a)	(b) and (c)		(a)	(b) and (c)		
Shallow	10.58	9.84	11.28	11.10	9.54	10.55	10.57
Deep	11.75	11.07	11.41	11.15	11.87	11.80	11.50
No dung	10.67	9.22	11.22	11.15	9.25	10.22	10.44
Dung	11.66	11.69	11.48	11.10	12.17	12.13	11.62
Mean	11.17	10.45	11.35	11.13	10.71	11.17	11.03
Plant Number: thousands per acre							
Shallow	27.1	27.5	27.5	27.0	27.5	27.6	27.3
Deep	27.0	27.1	26.7	27.3	26.6	26.6	27.0
No dung	27.1	27.5	27.6	27.3	27.1	27.5	27.3
Dung	27.0	27.1	26.6	27.0	27.0	26.7	26.9
Mean	27.0	27.3	27.1	27.2	27.1	27.1	27.1

	Total Sugar	Tops	
(a)	±1.54	±0.329	for use in comparisons other than horizontal
(b)	±1.44	±1.243	for use in horizontal comparisons
(c)	±1.85	±0.942	as (a).



54/Bb/1.5

Series 4: Sugar Beet

	Phosphate			Potash			Mean
	None	Ploughed in	In seed bed	None	Ploughed in	In seed bed	
Noxious Nitrogen: mg. per 100 mg.							
Shallow	29.4	27.5	36.2	33.1	26.2	30.0	30.6
Deep	35.6	35.0	36.2	33.3	32.5	32.5	35.6
No dung	33.8	26.2	38.8	35.0	27.5	35.0	33.1
Dung	31.2	36.2	33.8	36.9	31.2	27.5	33.1
Mean	32.5	31.2	36.2	35.9	29.4	31.2	33.1

Responses to treatments to previous Sugar Beet

Series 5: Barley

Response to	Mean	Ploughing		Dung		Phosphate		Potash	
		Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.

Grain: Mean yield 33.1 cwt per acre

Ploughing	(±0.91)	(±1.29)							
deep-shallow	+0.2	-	-	+0.9	-0.5	-0.7	+1.1	+0.5	-0.1
Dung	+2.5	+3.2	+1.8	-	-	+2.4	+2.6	+2.6	+2.4
Phosphate	+0.4	-0.5	+1.3	+0.3	+0.5	-	-	+0.1	+0.7
Potash	+0.8	+1.1	+0.5	+0.9	+0.7	+0.5	+1.1	-	-

Straw: Mean yield 38.0 cwt per acre

Ploughing									
deep-shallow	+1.1	-	-	+1.6	+0.6	+1.4	+0.8	+3.3	-1.1
Dung	+6.6	+7.1	+6.1	-	-	+6.0	+7.2	+6.1	+7.1
Phosphate	+1.1	+1.4	+0.8	+0.5	+1.7	-	-	+0.7	+1.5
Potash	+2.2	+4.4	0.0	+1.7	+2.7	+1.8	+2.6	-	-

Series 1: Ley

Hay: Mean yield 83.2 cwt per acre

Ploughing	(±2.66)	(±3.76)							
deep-shallow	+1.9	-	-	+5.6	-1.8	-1.0	+4.8	+1.9	+1.9
Dung	+3.6	+7.3	-0.1	-	-	+3.4	+3.8	+7.7	-0.5
Phosphate	+3.4	+0.5	+6.3	+3.2	+3.6	-	-	+6.4	+0.4
Potash	+2.1	+2.1	+2.1	+6.2	-2.0	+5.1	-0.9	-	-



54/Bb/1.6

Series 6: Wheat\*

Response to	Mean	Ploughing		Dung		Phosphate		Potash	
		Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.

Grain: Mean yield 38.4 cwt per acre

(±1.17)

(±1.66)

Ploughing deep-shallow	+2.1	-	-	+0.8	+3.4	+4.8	-0.6	+1.7	+2.5
Dung	+5.6	+4.3	+6.9	-	-	+5.7	+5.5	+3.6	+2.6
Phosphate	-1.4	+1.3	-4.1	-1.3	+1.5	-	-	-2.1	-0.7
Potash	+1.7	+1.3	+2.1	+4.7	-1.3	+1.0	+2.4	-	-

Straw: Mean yield 51.4 cwt per acre

Ploughing deep-shallow	+2.7	-	-	+2.6	+2.8	+4.7	+0.7	+4.9	+0.5
Dung	+8.3	+8.2	+8.4	-	-	+9.7	+6.9	+11.4	+5.2
Phosphate	+0.3	+2.3	-1.7	+1.7	-1.1	-	-	+1.1	-0.5
Potash	-0.4	+1.8	-2.6	+2.7	-3.5	+0.4	-1.2	-	-

Series 2: Potatoes

Total tubers: Mean yield 8.70 tons per acre

(±0.459)

(±0.650)

Ploughing deep-shallow	-1.11	-	-	-0.86	-1.36	-0.76	-1.46	-0.42	-1.80
Dung	+2.45	+2.70	+2.20	-	-	+3.35	+1.55	+3.36	+1.54
Phosphate	+0.62	+0.97	+0.27	+1.52	-0.28	-	-	+1.31	-0.07
Potash	+1.79	+2.48	+1.10	+2.70	+0.88	+2.48	+1.10	-	-

Percentage Ware (1½" riddle): Mean 84.1

Ploughing deep-shallow	-1.0	-	-	+0.2	-2.2	+2.9	-4.9	+0.1	-2.1
Dung	+0.8	+2.0	-0.4	-	-	+2.8	-1.2	+1.9	-0.3
Phosphate	-3.2	+0.7	-7.1	-1.2	-5.2	-	-	-0.8	-5.6
Potash	+3.5	+4.6	+2.4	+4.6	+2.4	+5.9	+1.1	-	-

\*Cultivation treatments direct to wheat, manures to previous sugar beet.



54/Bb/1.7

Series 2: Potatoes

	Phosphate			Potash			Mean
	None	Ploughed in	In ridges	None	Ploughed in	In ridges	
Total tubers: tons per acre							
	(a)	(b) and (c)		(a)	(b) and (c)		
Shallow	8.78	9.45	10.03	8.02	10.39	10.60	9.26
Deep	8.01	8.52	8.05	7.60	8.67	8.73	8.15
No dung	6.72	7.75	8.73	6.13	8.48	9.18	7.48
Dung	10.07	10.22	9.35	9.49	10.58	10.15	9.93
Mean	8.40	8.99	9.04	7.81	9.53	9.67	8.70
Percentage Ware (1½" riddle)							
Shallow	84.2	84.4	85.4	82.3	86.9	86.9	84.6
Deep	87.1	81.0	79.0	82.4	85.8	83.7	83.6
No dung	84.3	83.2	82.9	81.4	85.1	86.8	83.7
Dung	87.0	82.3	81.5	83.3	87.6	83.7	84.5
Mean	85.7	82.7	82.2	82.3	86.4	85.3	84.1

Responses to treatments to previous potatoes

Series 3: Spring Oats

Response to	Mean	Floughing		Dung		Phosphate		Potash	
		Shallow	Deep	Abs.	Pres.	Abs.	Pres.	Abs.	Pres.
Grain: Mean yield 34.8 cwt per acre									
	(±1.91)			(±2.70)					
Ploughing deep-shallow	-2.0	-	-	-1.2	-2.8	-3.2	-0.8	-3.2	-0.8
Dung	+1.9	+2.7	+1.1	-	-	+0.6	+3.2	-0.3	+4.1
Phosphate	+2.1	+0.9	+3.3	+0.8	+3.4	-	-	+2.6	+1.6
Potash	-0.5	-1.7	+0.7	-2.7	+1.7	0.0	-1.0	-	-
Straw: Mean yield 47.2 cwt per acre									
Ploughing deep-shallow	-1.7	-	-	-2.0	-1.4	-3.8	+0.4	-2.6	-0.3
Dung	+4.3	+4.0	+4.6	-	-	+4.2	+4.4	+3.7	+4.9
Phosphate	+0.3	-1.8	+2.4	+0.2	+0.4	-	-	+2.7	-2.1
Potash	-0.5	-1.4	+0.4	-1.1	+0.1	+1.9	-2.9	-	-

Total tubers

- (a) ±0.459 for use in comparisons other than horizontal.
- (b) ±0.495 for use in horizontal comparisons.
- (c) ±0.578 as (a).



54/Bc/1.1

## LEY AND ARABLE ROTATIONS

Highfield and Fosters Field 1954 - the 6th year.

For details of treatments, rotations, etc., see "Results of the Field Experiments 1952", Section Bc/1.

Cultivations, etc.:

### HIGHFIELD

#### 1st year Treatment Crops

Cut grass, Grazed ley, Lucerne, Hay. Ploughed: Aug 31 and for leys again, Oct 3, 1953. Basal fertilizer applied: Hay plots - Dec 2, remainder - Apr 7, 1954.

Cut grass: Nitrochalk applied, seeds sown at 38 lb per acre: Apr 7, 1954. Cut: 4 times - June 21, July 21, Aug 11, Oct 11.

Nitrochalk applied after each cut except the last.

Grazed ley: Nitrochalk applied, seeds sown at 55 lb per acre: Apr 7.

Nitrochalk applied: June 28. Grazed: 9 circuits, June 13-Oct 20.

Lucerne: Seed drilled at 28 lb per acre: Apr 9. Cut twice: Aug 11 and Nov 18. Variety: Du Puits.

Hay: Undersown seeds failed. Resown at 38 lb per acre: Sept 18, 1953.

Nitrochalk applied: Apr 2, 1954. Cut: June 22.

#### 2nd year Treatment Crops

Cut grass, Grazed ley, Lucerne, Potatoes. Basal fertilizer to leys applied: Dec 2, 1953.

Cut grass: Nitrochalk applied: Apr 2, 1954 and after each cut except the last. Cut: 5 times, May 20, June 21, July 22, Aug 17, Oct 11.

Grazed ley: Nitrochalk applied: Apr 6 and July 7. Grazed: 10 circuits, Apr 13 - Oct 20.

Lucerne: Cut: 3 times, June 23, Aug 11, Nov 18.

Potatoes: For cultivations see Potato Test Crop.

#### 3rd year Treatment Crops

Cut grass, Grazed ley, Lucerne, Barley. Basal fertilizer to leys applied: Dec 2, 1953.

Cut grass: Nitrochalk applied: Apr 2, 1954 and after each cut except the last. Cut: 5 times, May 20, June 21, July 23, Aug 17, Oct 8.

Grazed ley: Nitrochalk applied: Apr 2 and June 28. Grazed: 10 circuits, Apr 17 - Oct 20.

Lucerne: Cut: 3 times, June 22, Aug 11, Oct 8.

Barley: For cultivations see Barley Test Crop.



54/Bc/1.2

1st Test Crop, Wheat

Ploughed after barley: Aug 31 and again Oct 3, 1953. Ploughed leys: Oct 16. Seed drilled at 3 bushels per acre with basal fertilizer: Oct 22. Nitrochalk applied: Apr 27, 1954. Combine harvested: Sept 2 and 7. Variety: Yeoman.

2nd Test Crop, Potatoes

Ploughed: Aug 24 and again Oct 26, 1953. Ridged, basal fertilizers applied: Apr 20, 1954. Dung and sulphate of ammonia applied, potatoes planted: Apr 22. Earthed up: July 9. Sprayed with copper fungicide, low volume, 5 lb in 10 gallons per acre: July 28 and again Aug 23. Sprayed with 20% sulphuric acid: Sept 30. Lifted: Oct 26. Variety: Majestic.

3rd Test Crop, Barley

Ploughed: Oct 3, 1953. Ground chalk applied to blocks 2 and 3: Dec 3. Seed drilled at 2 bushels per acre with basal fertilizer, nitrochalk applied: Mar 17, 1954. Combine harvested: Sept 3. Variety: Proctor. Note. Crop yields were not taken owing to the condition of the crop after being laid since early July.

Permanent Grasses

- 4th year Reseeded. Permanent grass, Blocks 9-12.  
Basal fertilizers applied: Dec 2, 1953. Nitrochalk applied: Apr 7, 1954 and July 7. Grazed: 8 circuits, Apr 21 - Oct 28.
- 5th year Reseeded. Permanent grass, Blocks 5-8.  
Basal fertilizers applied: Dec 2, 1953. Nitrochalk applied: Apr 6 and June 28. Reseeded, grazed: 9 circuits, Apr 13 - Oct 28.  
Permanent, grazed: 8 circuits, Apr 13 - Oct 20.
- 6th year Reseeded. Permanent grass, Blocks 1-4.  
Basal fertilizers applied: Dec 2, 1953. Nitrochalk applied: Apr 2.  
Cut: June 22. Nitrochalk applied: June 25. Grazed: 4 circuits, July 16 - Oct 28.

FOSTERS

1st year Treatment Crops

- Cut grass, Grazed ley, Lucerne, Hay. Ploughed (not hay plots): Aug 20, 1953 and again Sept 24. Basal fertilizer applied: Hay plots - Dec 1, remainder - Apr 7.
- Cut grass: Nitrochalk applied, seeds sown at 38 lb per acre: Apr 7, 1954. Cut: 4 times, June 29, July 20, Aug 13, Oct 11. Nitrochalk applied after each cut except the last.
- Grazed ley: Nitrochalk applied, seeds sown at 55 lb per acre: Apr 7. Nitrochalk applied: June 29. Grazed: 7 circuits, May 7 - Oct 21.
- Lucerne: Seed drilled at 28 lb per acre: Apr 9. Cut twice: Aug 16, Nov 18. Variety: Du Puits.
- Hay: Seeds undersown in barley at 38 lb per acre: May 4, 1953. Nitrochalk applied: Apr 1, 1954. Cut: June 23.



54/Bc/1.3

### 2nd year Treatment Crops

Cut grass, Grazed ley, Lucerne, Potatoes. Basal fertilizers to leys applied: Dec 1, 1953.

Cut grass: Nitrochalk applied: Apr 1, 1954 and after each cut except the last. Cut: 5 times, May 21, June 21, July 20, Aug 16, Oct 11.

Grazed ley: Nitrochalk applied: Apr 5 and July 9. Grazed: 10 circuits, May 7 - Oct 21.

Lucerne: Cut: 3 times, June 23, Aug 16, Nov 18.

Potatoes: For cultivations see Potato Test Crop.

### 3rd year Treatment Crops

Cut grass, Grazed ley, Lucerne, Barley. Basal fertilizer to leys applied: Dec 1, 1953.

Cut grass: Nitrochalk applied: Apr 1, 1954 and after each cut except the last. Cut: 5 times, May 21, June 21, July 20, Aug 16, Oct 8.

Grazed ley: Nitrochalk applied: Apr 5 and June 29. Grazed: 8 circuits, May 7 - Oct 21.

Lucerne: Cut: 3 times, June 24, Aug 16, Oct 8.

Barley: For cultivations see Barley Test Crop.

### 1st Test Crop, Wheat

Ploughed after barley: Aug 20, 1953 and again Sept 29. Ploughed leys: Oct 17. Seed drilled at 3 bushels per acre with basal fertilizer: Oct 22. Nitrochalk applied: Apr 28, 1954. Combine harvested: Sept 1. Variety: Yeoman.

### 2nd Test Crop, Potatoes

Ploughed: Aug 21, 1953 and again Oct 24. Ridged, basal fertilizers applied: Apr 20, 1954. Dung and sulphate of ammonia applied, potatoes planted: Apr 21. Earthed up: July 6. Sprayed with copper fungicide, low volume, 5 lb in 10 gallons: July 28 and again Aug 23. Sprayed with 1% sulphuric acid: Sept 29. Lifted: Oct 9 and Oct 16. Variety: Majestic.

### 3rd Test Crop, Barley

Ploughed: Oct 2, 1953. Nitrochalk applied: Mar 16, 1954. Seed sown at 2 bushels per acre with basal fertilizer: Mar 17. Combine harvested: Sept 2. Variety: Proctor.

### Permanent grasses

#### 4th year reseeded grass

Basal fertilizer applied: Dec 1, 1953. Nitrochalk applied: Apr 7, 1954 and July 9. Grazed: 7 circuits, May 7 - Oct 21.

#### 5th year reseeded grass

Basal fertilizer applied: Dec 1, 1953. Nitrochalk applied: Apr 5, 1954 and June 29. Grazed: 7 circuits, May 7 - Oct 21.

#### 6th year reseeded grass

Basal fertilizer applied: Dec 2, 1953. Nitrochalk applied: Apr 1, 1954. Cut: June 23. Nitrochalk applied: June 25. Grazed: 4 circuits, July 25 - Oct 21.



54/Bc/1.4

Standard errors per  $\frac{1}{4}$  plot. Test crops.

Wheat, grain (at 85% dry matter). Highfield: 2.48 cwt per acre or 8.9% (13 d.f.)  
 Fosters: 1.82 cwt per acre or 5.2% (13 d.f.)  
 Potatoes, total tubers. Highfield: 1.073 tons per acre or 10.1% (15 d.f.)  
 Fosters: 0.876 tons per acre or 8.6% (15 d.f.)  
 Barley, grain (at 85% dry matter). Highfield: Crop failed  
 Fosters: 1.87 cwt per acre or 4.2% (15 d.f.)

Summary of Results

cwt N per acre	Wheat 1st test crop Previous rotation 1951, 1952, 1953				Mean
	Lucerne	Ley	Cut Grass	Arable with hay	
Grain (at 85% Dry Matter): cwt per acre					
<u>Highfield</u>					
Mean	28.6	32.7	22.7	27.9	28.0
To test crop					
0.3	29.5	34.8	24.8	28.6	29.4
0.6	27.8	30.6	20.6	27.1	26.5
Difference ( $\pm 1.75$ )	-1.7	-4.2	-4.2	-1.5	-2.9 ( $\pm 0.88$ )
To treatment crops					
Single rate		35.5	25.7	24.8	28.7
Double rate		29.9	19.7	30.9	26.8
Difference ( $\pm 1.75$ )		-5.6	-6.0	+6.1	-1.9 ( $\pm 1.01$ )
<u>Fosters</u>					
Mean	37.4	39.5	38.2	24.7	35.0
To test crop					
0.3	36.0	37.6	38.4	23.9	34.0
0.6	38.7	41.5	38.0	25.6	36.0
Difference ( $\pm 1.28$ )	+2.7	+3.9	-0.4	+1.7	+2.0 ( $\pm 0.64$ )
To treatment crops					
Single rate		39.4	39.1	22.3	33.6
Double rate		39.7	37.4	27.1	34.7
Difference ( $\pm 1.28$ )		+0.3	-1.7	+4.8	+1.1 ( $\pm 0.74$ )



Wheat 1st test crop

54/Bc/1.5

cwt N per acre	Excluding Lucerne N to previous treatment crop			Arable with hay only Dung: tons per acre to potatoes 1952		
	Single rate	Double rate	Mean	None	12	Mean

Grain (at 85% Dry Matter): cwt per acre

Highfield

To test crop	(±1.01)		(±0.71)	(±1.75)		(±1.24)
0.3	30.6	28.2	29.4	29.2	28.0	28.6
0.6	26.8	25.4	26.1	22.6	31.7	27.1
Mean	28.7	26.8	27.8	25.9	29.9	27.9
	(±0.71)			(±1.24)		
To previous treatment crops				(±1.75)		(±1.24)
Single rate				25.8	23.9	24.8
Double rate				26.0	35.8	30.9
Mean				25.9	29.9	27.9
				(±1.24)		

Fosters

To test crop	(±0.74)		(±0.52)	(±1.28)		(±0.91)
0.3	32.7	33.9	33.3	24.0	23.8	23.9
0.6	34.5	35.5	35.0	28.1	23.1	25.6
Mean	33.6	34.7	34.2	26.1	23.4	24.7
	(±0.52)			(±0.91)		
To previous treatment crops				(±1.28)		(±0.91)
Single rate				25.0	19.7	22.3
Double rate				27.1	27.1	27.1
Mean				26.1	23.4	24.7
				(±0.91)		



cwt N per acre	<u>Wheat 1st test crop</u>				54/Bc/1.6
	Previous rotation 1951, 1952, 1953				Mean
	Lucerne	Ley	Cut Grass	Arable with hay	

Straw (at 85% Dry Matter): cwt per acre

	<u>Highfield</u>				
Mean	48.1	45.9	32.8	43.8	42.7
To test crop					
0.3	47.3	44.0	34.1	46.4	42.9
0.6	49.0	47.8	31.5	41.2	42.4
Difference	+1.7	+3.8	-2.6	-5.2	-0.5
To treatment crops					
Single rate		53.7	35.6	45.5	45.0
Double rate		38.1	29.9	42.1	36.7
Difference		-15.6	-5.7	-3.4	-8.3
	<u>Fosters</u>				
Mean	61.6	55.9	50.6	47.9	54.0
To test crop					
0.3	61.2	53.0	48.9	43.7	51.7
0.6	62.1	58.8	52.3	52.2	56.3
Difference	+0.9	+5.8	+3.4	+8.5	+4.6
To treatment crops					
Single rate		56.9	51.2	46.8	51.6
Double rate		54.9	50.1	49.1	51.4
Difference		-2.0	-1.1	+2.3	-0.2



Wheat 1st test crop

54/Bc/1.7

cwt N per acre	Excluding Lucerne N to previous treatment crop			Arable with hay only Dung: tons per acre to potatoes 1952		
	Single rate	Double rate	Mean	None	12	Mean

Straw (at 85% Dry Matter): cwt per acre

Highfield

To test crop						
0.3	46.0	37.0	41.5	46.2	46.7	46.4
0.6	43.9	36.5	40.2	41.5	40.9	41.2
Mean	45.0	36.7	40.8	43.8	43.8	43.8
To previous treatment crops						
Single rate				46.9	44.2	45.5
Double rate				40.8	43.4	42.1
Mean				43.8	43.8	43.8

Fosters

To test crop						
0.3	49.1	48.0	48.5	40.5	46.9	43.7
0.6	54.1	54.7	54.4	51.6	52.7	52.2
Mean	51.6	51.4	51.5	46.0	49.8	47.9
To previous treatment crops						
Single rate				45.6	47.9	46.8
Double rate				46.5	51.7	49.1
Mean				46.0	49.8	47.9



54/Bc/1.8

Potatoes 2nd test crop. Total tubers: tons per acre

	Previous rotation 1950, 1951, 1952				Mean																									
	Lucerne	Ley	Cut Grass	Arable with hay																										
<u>Highfield</u>																														
N: cwt per acre																														
0.5	9.88	10.90	10.42	10.41	10.40																									
1.0	11.04	11.15	10.41	10.94	10.89																									
Difference ( $\pm 0.759$ )	+1.16	+0.25	-0.01	+0.53	+0.49 ( $\pm 0.379$ )																									
Dung: tons per acre																														
None	8.89	9.18	8.87	8.59	8.88																									
12	12.03	12.88	11.95	12.76	12.40																									
Difference ( $\pm 0.759$ )	+3.14	+3.70	+3.08	+4.17	+3.52 ( $\pm 0.379$ )																									
Mean	10.46	11.03	10.41	10.68	10.64																									
<u>Fosters</u>																														
N: cwt per acre																														
0.5	9.58	9.63	9.22	10.17	9.65																									
1.0	10.82	10.73	9.80	11.52	10.72																									
Difference ( $\pm 0.619$ )	+1.24	+1.10	+0.58	+1.35	+1.07 ( $\pm 0.310$ )																									
Dung: tons per acre																														
None	7.59	9.23	7.74	9.75	8.58																									
12	12.81	11.12	11.28	11.95	11.79																									
Difference ( $\pm 0.619$ )	+5.22	+1.89	+3.54	+2.20	+3.21 ( $\pm 0.310$ )																									
Mean	10.20	10.18	9.51	10.85	10.18																									
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th colspan="2">Highfield</th> <th colspan="2">Fosters</th> </tr> <tr> <th>N: cwt per acre</th> <th>0.5</th> <th>1.0</th> <th>0.5</th> <th>1.0</th> </tr> </thead> <tbody> <tr> <td>Dung: tons per acre</td> <td colspan="2" style="text-align: center;">(<math>\pm 0.379</math>)</td> <td colspan="2" style="text-align: center;">(<math>\pm 0.310</math>)</td> </tr> <tr> <td>None</td> <td>8.63</td> <td>9.14</td> <td>7.99</td> <td>9.17</td> </tr> <tr> <td>12</td> <td>12.18</td> <td>12.63</td> <td>11.31</td> <td>12.27</td> </tr> </tbody> </table>							Highfield		Fosters		N: cwt per acre	0.5	1.0	0.5	1.0	Dung: tons per acre	( $\pm 0.379$ )		( $\pm 0.310$ )		None	8.63	9.14	7.99	9.17	12	12.18	12.63	11.31	12.27
	Highfield		Fosters																											
N: cwt per acre	0.5	1.0	0.5	1.0																										
Dung: tons per acre	( $\pm 0.379$ )		( $\pm 0.310$ )																											
None	8.63	9.14	7.99	9.17																										
12	12.18	12.63	11.31	12.27																										



54/Bc/1.9

Potatoes 2nd test crop. Percentage ware

	Previous rotation Lucerne	1950, 1951, 1952 Ley	Cut Grass	Arable with hay	Mean																														
<u>Highfield</u>																																			
N: cwt per acre																																			
0.5	82.6	81.2	81.0	82.1	81.7																														
1.0	84.4	81.2	77.8	82.8	81.6																														
Difference	+1.8	0.0	-3.2	+0.7	-0.1																														
Dung: tons per acre																																			
None	79.3	79.6	80.0	78.9	79.5																														
12	87.6	82.8	78.7	86.0	83.8																														
Difference	+8.3	+3.2	-1.3	+7.1	+4.3																														
Mean	83.5	81.2	79.4	82.5	81.6																														
<u>Fosters</u>																																			
N: cwt per acre																																			
0.5	83.6	83.2	84.3	84.5	83.9																														
1.0	84.4	78.5	83.0	86.4	83.1																														
Difference	+0.8	-4.7	-1.3	+1.9	-0.8																														
Dung: tons per acre																																			
None	82.7	80.1	82.8	84.8	82.6																														
12	85.4	81.6	84.5	86.1	84.4																														
Difference	+2.7	+1.5	+1.7	+1.3	+1.8																														
Mean	84.0	80.9	83.6	85.5	83.5																														
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th colspan="2" style="text-align: center;"><u>Highfield</u></th> <th colspan="2" style="text-align: center;"><u>Fosters</u></th> </tr> <tr> <th></th> <th colspan="2" style="text-align: center;">N: cwt per acre</th> <th colspan="2" style="text-align: center;">N: cwt per acre</th> </tr> <tr> <th></th> <th style="text-align: center;">0.5</th> <th style="text-align: center;">1.0</th> <th style="text-align: center;">0.5</th> <th style="text-align: center;">1.0</th> </tr> </thead> <tbody> <tr> <td>Dung: tons per acre</td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>None</td> <td style="text-align: center;">79.3</td> <td style="text-align: center;">79.6</td> <td style="text-align: center;">83.1</td> <td style="text-align: center;">82.1</td> </tr> <tr> <td>12</td> <td style="text-align: center;">84.1</td> <td style="text-align: center;">83.5</td> <td style="text-align: center;">84.7</td> <td style="text-align: center;">84.1</td> </tr> </tbody> </table>							<u>Highfield</u>		<u>Fosters</u>			N: cwt per acre		N: cwt per acre			0.5	1.0	0.5	1.0	Dung: tons per acre					None	79.3	79.6	83.1	82.1	12	84.1	83.5	84.7	84.1
	<u>Highfield</u>		<u>Fosters</u>																																
	N: cwt per acre		N: cwt per acre																																
	0.5	1.0	0.5	1.0																															
Dung: tons per acre																																			
None	79.3	79.6	83.1	82.1																															
12	84.1	83.5	84.7	84.1																															



54/Bc/1.10

Barley 3rd test crop. Grain (at 85% Dry Matter): cwt per acre

	Previous rotation 1949, 1950, 1951				Mean
	Lucerne	Ley	Cut Grass	Arable with hay	
<u>Fosters</u>					
N: cwt per acre					
0.2	46.6	43.5	44.4	44.3	44.7
0.4	44.8	44.5	44.7	45.2	44.8
Difference ( $\pm 1.32$ )	-1.8	+1.0	+0.3	+0.9	+0.1 ( $\pm 0.66$ )
Dung to potatoes 1953: tons per acre					
None	45.8	43.9	44.7	44.6	44.8
12	45.7	44.1	44.4	44.9	44.7
Difference ( $\pm 1.32$ )	-0.1	+0.2	-0.3	+0.3	-0.1 ( $\pm 0.66$ )
Mean	45.7	44.0	44.6	44.7	44.8

	Fosters	
	N: cwt per acre	
	0.2	0.4
	( $\pm 0.66$ )	
Dung to Potatoes 1953: tons per acre		
None	44.7	44.8
12	44.7	44.8

Note: Highfield: No yields of grain and straw recorded.  
Fosters: No straw yields recorded.



54/Bc/1.11

Treatment crops Arable and Hay rotation  
(values based on Mean of 2 sub plots only)

	Highfield			Fosters		
	N: cwt per acre applied in 1954			N: cwt per acre applied in 1954		
	Single rate	Double rate	Mean	Single rate	Double rate	Mean
Hay (dry matter): cwt per acre						
No dung	59.1	68.8	63.9	55.1	65.8	60.5
Dung in 1952	62.4	62.3	62.4	58.0	64.5	61.3
Mean	60.7	65.6	63.1	56.6	65.2	60.9
Potatoes, total tubers: tons per acre						
No dung	8.03	8.61	8.32	9.25	10.85	10.05
Dung in 1954	12.11	12.42	12.27	10.59	11.77	11.18
Mean	10.07	10.51	10.29	9.92	11.31	10.61
Potatoes, percentage ware						
No dung	77.8	81.0	79.4	82.6	84.4	83.5
Dung in 1954	79.6	83.1	81.4	88.1	85.6	86.8
Mean	78.7	82.1	80.4	85.4	85.0	85.2
Barley, grain (at 85% dry matter): cwt per acre						
No dung	No yields of grain recorded			41.6	44.2	42.9
Dung in 1953	No yields of grain recorded			43.6	43.3	43.4
Mean	No yields of grain recorded			42.6	43.8	43.2
	No yields of straw recorded			No yields of straw recorded		



Cut grass. Dry Matter: cwt per acre

54/Bc/1.12

1st year	Highfield					Fosters				
	N: to previous 3 test crops		Dung to potatoes 1952 tons per acre		Mean	N: to previous 3 test crops		Dung to potatoes 1952 tons per acre		Mean
Single rate	Double rate	None	12	Single rate		Double rate	None	12		
N(1) to cut grass										
Single rate	77.4	80.7	77.8	80.3	79.1	60.6	60.0	63.5	57.1	60.3
Double rate	81.6	89.4	89.0	82.1	85.5	73.9	76.4	73.6	76.7	75.1
N to test crops										
Single rate			80.1	78.9	79.5			67.2	67.3	67.3
Double rate			86.6	83.5	85.0			69.8	66.6	68.2
Mean			83.4	81.2	82.3			68.5	66.9	67.7

	Highfield			Fosters		
	N to cut grass (1)		Mean	N to cut grass (1)		Mean
	Single rate	Double rate		Single rate	Double rate	
2nd year (5 cuts)	59.1	75.3	67.2	71.1	76.6	73.8
3rd year (5 cuts)	57.5	72.7	65.1	65.5	74.4	70.0

(1) 0.15 v. 0.3 cwt N as Nitrochalk for every cut.

Lucerne. Dry Matter: cwt per acre

1st year (2 cuts)	Highfield			Fosters		
	N to 3 previous test crops		Mean	N to 3 previous test crops		Mean
Single rate	Double rate	Single rate		Double rate		
Dung to potatoes 1952						
None	62.0	58.4	60.2	56.3	56.9	56.6
12 tons	57.6	57.3	57.4	56.3	60.6	58.4
Mean	59.8	57.9	58.8	56.3	58.7	57.5

	Highfield	Fosters
2nd year (3 cuts)	Mean 103.8	Mean 97.5
3rd year (3 cuts)	Mean 100.0	Mean 94.4



54/Bc/1.13

Grazed Ley. Dry Matter: cwt per acre (estimated from sampling cuts)

	Highfield			Fosters		
	N: cwt per acre (yearly)		Mean	N: cwt per acre (yearly)		Mean
	Single rate	Double rate		Single rate	Double rate	
	0.15	0.30		0.15	0.30	
1st year	40.8	45.9	43.3	31.2	30.4	30.8
2nd year	39.2	52.2	45.7	40.5	53.4	46.9
3rd year	43.6	48.9	46.2	33.5	36.7	35.1

Reseeded Grass. Dry Matter: cwt per acre

	Cut for hay			Grazed Estimated from sampling cuts		
	N		Mean	N		Mean
	Single rate	Double rate		Single rate	Double rate	
4th year, grazing				49.9	47.6	48.7
5th year, grazing				42.5	45.7	44.1
6th year, hay	55.4	55.5	55.5	17.2*	21.2*	19.2*
	<u>Highfield</u>					
4th year, grazing				41.4	41.6	41.5
5th year, grazing				44.1	43.3	43.7
6th year, hay	46.8	51.8	49.3	20.0*	20.4*	20.2*

Permanent Grass. Dry Matter: cwt per acre

	Highfield					
	Single rate	Double rate	Mean	Single rate	Double rate	Mean
Grazing, Blocks 9-12				40.0	39.9	39.9
Grazing, Blocks 5-8				36.6	36.6	36.6
Hay, Blocks 1-4	44.8	47.1	45.9	19.7*	20.7*	20.2*

\*Aftermath grazing.



54/Bd/1.1

GREEN MANURING EXPERIMENT

Woburn Stackyard - 1954, the 1st year of revised scheme

Original scheme: Details are given in "Results of the Field Experiments 1939-47", Vol I, Section Be/1. In 1950 the fallow, lupins and ryegrass plots were split for early and late planting of cabbages. The original scheme ended with the harvesting of barley in 1953 and the cabbages in 1953-4.

Revised scheme: From 1954 onwards the rotation is: early potatoes, barley. As before each of these two crops is grown in every year on one of two randomized blocks of 40 plots each. The green manuring crops are grown according to the following scheme which is repeated every two years:

1st main crop	Early Potatoes	Early Potatoes	Early Potatoes	Early Potatoes	Early Potatoes
Green manure	-	Ryegrass	Ryegrass	Trefoil	Trefoil
2nd main crop and green manure	Barley	Barley undersown with Ryegrass	Barley	Barley undersown with Trefoil	Barley

8 plots of each block are allocated to each of these sequences. Half the plots of each group carrying ryegrass or trefoil after early potatoes are ploughed in autumn and the remainder are ploughed in the spring before the barley seedbed is prepared. The undersown green manures are ploughed in after February 1st for early potatoes.

In addition chaffed barley straw at the rate of 30 cwt per acre is applied after harvesting the barley to the plots receiving straw in the original scheme. Two levels of nitrogen are tested on each of the two main crops.

0.23 v. 0.46 cwt N per acre as nitrochalk to barley  
0.6 v. 1.2 cwt N per acre as nitrochalk to potatoes

the higher level in each case being applied to the same plots.

The fallow plots of the original scheme remain fallow between each main crop in the revised scheme. The new green manuring treatments are superimposed on the plots carrying the original treatments in such a way that one comparison of the latter (lupins and rape v. clover and ryegrass) can be examined for possible residual effects. Residual effects of the original dung treatment, now discontinued, can also be determined, but any residual effects of the nitrogen treatments applied prior to 1954 have been eliminated by randomization. The green manuring and subsidiary treatments are arranged on the 32 non-fallow plots of each block in a quarter replicate with identities:



54/Bd/1.2

$I \equiv (D) SPUGN \equiv (D)(X)UN \equiv S(X)PG$

where (D) = (residual) dung.  
(X) = (residual) rape and lupins v. clover and ryegrass.  
S = straw.  
P = time of ploughing green manures after early potatoes.  
U = green manures undersown (in addition to those sown after early potatoes).  
G = trefoil v. ryegrass.  
N = nitrogen levels to both crops.

Basal dressing: Early potatoes, 0.75 cwt  $P_2O_5$ , 1.5 cwt  $K_2O$  per acre as granular compound fertilizer, broadcast on the flat before machine planting. Barley and green manures, nil.

Varieties: Early potatoes: Ulster Chieftain  
Barley: Herta  
Trefoil: English  
Ryegrass: English Leafy Italian.

Plot area: 0.0395 acre.

Transition Period:

The barley of 1954 received two levels of nitrogen and was undersown according to the new scheme.

The early potatoes of 1954 received two levels of nitrogen according to the new scheme. It is not possible to make full comparisons of the old treatments in either of these crops. A full analysis of the new scheme will first be possible in 1955.

Cultivations, etc.:

Green manures: Clover and ryegrass undersown in barley: Apr 24, 1953.

Barley: Ploughed: Mar 5, 1954. Ground chalk applied: Mar 11.

Nitrochalk applied: Mar 16. Seed drilled at 3 bushels per acre: Mar 17. Trefoil and Italian ryegrass undersown: Apr 27.

Harvested: Aug 30. Variety: Herta.

Early potatoes: Ploughed (except for green manured plots):

Aug 21, 1953, Oct 2, Dec 2, and Mar 1, 1954. Ploughed (all plots): Mar 10. Nitrochalk and basal fertilizers applied:

Mar 24. Potatoes mechanically planted: Mar 26. Earthed up: June 15 and again June 25. Lifted: July 28. Variety: Ulster Chieftain.

Standard errors per plot:

Early potatoes, Total tubers: 0.973 tons per acre or 21.8% (22 d.f.)

Barley, Grain: 3.04 cwt per acre or 7.8% (18 d.f.)



Summary of Results

Early potatoes, total tubers: tons per acre

N in 1954: cwt per acre	Green manures and treatments to cabbages 1952							Mean
	Fallow	Clover* + Rye- grass*	Lupins + Rape	Dung: tons per acre		Straw: tons per acre		
				None	10	None	1½	
	(±0.486)	(±0.344)		(±0.308)				(±0.217)
0.6	3.69	4.22	4.19	3.24	4.96	4.03	4.17	4.10
1.2	4.37	4.82	4.65	4.02	5.30	4.32	5.00	4.66
Mean	4.03 (±0.344)	4.52 (±0.243)	4.42	3.63	5.13 (±0.217)	4.18	4.59	4.38

\*Also undersown in barley 1953.

Barley, grain: cwt per acre

N in 1954: cwt per acre	Green manures and treatments to cabbages 1953							Mean
	Fallow	Clover + Rye- grass	Lupins + Rape	Dung: tons per acre		Straw: tons per acre		
				None	10	None	1½	
	(±1.50)	(±1.06)		(±0.95)				(±0.67)
0.23	32.2	36.9	35.5	33.5	37.2	35.6	35.1	35.4
0.46	36.8	40.3	41.7	40.2	40.1	40.5	39.8	40.1
Mean	34.5 (±1.06)	38.6 (±0.75)	38.6	36.8	38.7 (±0.67)	38.1	37.5	37.8

Excluding fallow plots

N in 1954: cwt per acre	Undersown in barley 1954			Mean
	None	Trefoil	Eye-grass	
	(±1.06)	(±1.50)		(±0.75)
0.23	36.0	36.9	35.8	36.2
0.46	40.0	41.7	42.1	41.0
Mean	38.0 (±0.75)	39.3 (±1.06)	39.0	38.6



54/Bd/1.4

Barley, straw: cwt per acre

N in 1954: cwt per acre	Green manures and treatments to cabbages 1953							Mean
	Fallow	Clover + Rye-grass	Lupins + Rape	Dung: tons per acre		Straw: tons per acre		
				None	10	None	1½	
0.23	31.5	34.6	33.3	30.2	36.7	33.5	33.4	33.5
0.46	35.8	39.3	39.4	37.6	39.7	38.0	39.3	38.6
Mean	33.7	37.0	36.3	33.9	38.2	35.8	36.3	36.1

Excluding fallow plots

N in 1954: cwt per acre	Undersown in barley 1954			Mean
	None	Trefoil	Ryegrass	
0.23	34.0	36.9	31.0	33.9
0.46	38.7	40.2	39.7	39.4
Mean	36.4	38.5	35.3	36.6



54/Be/1.1

## LEY AND ARABLE ROTATIONS

Woburn Stackyard - 1954 the 17th year.

For details of rotations and treatments etc., see "Results of the Field Experiments 1939-47", Vol. I, Section Bf/1 with the following exceptions:-

In 1949 and subsequently Rye replaced Wheat.

In 1954 the Seeds Hay plots were split into two after the first crop, for testing 0.15 v. 0.30 cwt N per acre applied as nitro-chalk.

Cultivations, etc.:

### Treatment crops

#### Ley rotations

- Ley 1st year. Ploughed twice: Sept 16 and Dec 30, 1953.  
Basal fertilizers applied: Apr 5, 1954. Seeds mixture hand sown: Apr 7. Nitrochalk applied: July 16. Grazed 7 times: June 17-21, July 7-15, July 24-Aug 1, Aug 11-21, Sept 8-17, Sept 25-Oct 1, Oct 19-27. Seeds mixture per acre: 21 lb S23 Perennial Ryegrass, 12 lb S143 Cocksfoot, 6 lb Late flowering Montgomery Red Clover, 3 lb S100 White Clover.
- Ley 2nd year. Nitrochalk applied: May 25 and July 26.  
Grazed 9 times: May 17-25, June 8-17, June 21-29, July 15-24, Aug 1-11, Aug 21-30. Sept 17-25, Oct 3-11, Oct 27-Nov 4.
- Ley 3rd year. Nitrochalk applied: May 25 and July 21.  
Grazed 5 times: May 13-17, May 31-June 8, June 29-July 7, Aug 30-Sept 8, Oct 11-19.
- Lucerne 1st year. Ploughed twice: Sept 16 and Dec 30, 1953.  
Basal fertilizers applied: Apr 2, 1954. Seeds sown at 28 lb per acre: Apr 7. Dusted with 2% DDT: Apr 8. Cut twice: Aug 11 and Nov 3. Variety: Du Puits.
- Lucerne 2nd year. Cut three times: June 24, Aug 11, Nov 3.
- Lucerne 3rd year. Cut three times: June 24, Aug 11, Nov 3.

#### Arable rotations

- Potatoes 1st course. Ploughed: Sept 16 and again Dec 30, 1953.  
Ridged, basal fertilizers applied: Apr 2, 1954. Potatoes planted with dropper: Apr 6. Earthed up: June 25. Sprayed with copper fungicide, 5 lb per acre: July 30, Aug 16 and Aug 27. Sprayed with sulphuric acid: Sept 23. Lifted: Oct 4. Variety: Majestic.
- Rye 2nd course. Ploughed: Oct 1, 1953. Seed drilled at 3 bushels per acre: Oct 22. Nitrochalk applied: Apr 29, 1954. Seeds hay mixture undersown on 4 plots: May 7. Harvested: Aug 23.
- Seeds Hay 3rd course. Seeds undersown in Rye: Apr 9, 1953.  
Basal nitrochalk applied: Apr 6, 1954. 1st cut: June 24. Nitrochalk applied: June 30. 2nd cut: Nov 3. Seeds mixture per acre: 27 lb S24 Perennial Ryegrass, 12 lb Montgomery Red Clover, 3 lb Canadian Alsike Clover.



54/Be/1.2

Sugar beet 3rd course. Ploughed: Aug 21 and again Nov 5, 1953.  
 Basal nitrate of soda applied: Mar 29, 1954. Seed drilled  
 at 18 lb per acre: Mar 30. Dusted with DDT: May 12.  
 Singled: May 28. Sprayed with systemic insecticide  $\frac{1}{2}$  pint  
 per acre, high volume: June 21. Lifted: Nov 2. Variety:  
 Klein E.

Test Crops.

Potatoes 1st test crop. Ploughed: Nov 4, 1953. Ridged, dung  
 applied: Apr 7, 1954. Basal fertilizers applied, potatoes  
 hand planted: Apr 8. Earthed up: June 25. Sprayed with  
 copper fungicide, 5 lb per acre: July 30, Aug 16 and Aug 27.  
 Sprayed with 20% sulphuric acid: Sept 23. Lifted: Oct 4.  
 Variety: Majestic.

Barley 2nd test crop. Ploughed: Oct 2 and again Dec 30, 1953.  
 Ground chalk applied: Mar 8, 1954. Nitrochalk applied:  
 Mar 11. Seed drilled at 3 bushels per acre: Mar 17.  
 Harvested: Aug 28. Variety: Plumage Archer.

Standard errors per plot, Test crops:

Potatoes, Total tubers, whole plot:	0.797 tons per acre or 7.3%	(4 d.f.)
sub plot:	0.881 tons per acre or 8.1%	(4 d.f.)
Barley, Grain, whole plot:	2.50 cwt per acre or 7.8%	(4 d.f.)
sub plot:	2.12 cwt per acre or 6.7%	(4 d.f.)

Summary of Results

Treatment crops

Ley, Sheep days of grazing per acre

1st year	2nd year	3rd year
1817	2616	1018



54/Be/1.3

Treatment crops

Lucerne, yield of hay (at 85% D.M.): cwt per acre

	1st crop	2nd crop	3rd crop	Total
<u>1st year</u>				
No dung	18.7	11.4		30.1
Dung in 1952	26.1	14.1		40.2
Increase	7.4	2.7		10.1
Previous Rotation				
Lucerne	20.9	12.3		33.2
Arable with Hay	23.9	13.2		37.1
Mean	22.4	12.8		35.2
<u>2nd year</u>				
No dung	38.0	24.0	9.0	71.0
Dung in 1951	44.3	25.8	11.8	81.9
Increase	6.3	1.8	2.8	10.9
Previous Rotation				
Lucerne	38.8	24.3	9.6	72.7
Arable with Sugar beet	43.5	25.5	11.2	80.2
Mean	41.2	24.9	10.4	76.5
<u>3rd year</u>				
No dung	32.5	18.9	6.4	57.8
Dung in 1950	39.2	24.9	11.0	75.1
Increase	6.7	6.0	4.6	17.3
Previous Rotation				
Lucerne	35.4	23.7	9.4	68.5
Arable with Hay	36.3	20.1	8.0	64.4
Mean	35.8	21.9	8.7	66.4
	Potatoes		Rye	
	Total tubers: tons per acre	Percentage ware	Grain: cwt per acre	Straw:
No dung	8.52	85.3	32.0	33.5
Dung <sup>#</sup>	11.71	88.3	35.3	37.9
Increase	3.19	3.0	3.3	4.4
Previous Rotation				
Ley	11.84	91.3	33.7	34.8
Lucerne	10.00	87.9	36.7	38.1
Arable with Hay	9.48	83.5	34.5	35.6
Arable with Sugar beet	9.13	84.4	29.8	34.3
Mean	10.12	86.8	33.7	35.7

<sup>#</sup>Dung applied: Potatoes:- 1952 Rye:- 1951.



54/Be/1.4

Treatment crops				
	Hay			
	Yield (at 85% D.M.): cwt per acre			
	1st crop	2nd crop	Total	2nd crop Response to N
No dung	39.4	2.7	42.1	0.5
Dung in 1950	40.5	4.9	45.4	2.0
Increase	1.1	2.2	3.3	1.5
Previous Rotation				
Lucerne	39.8	4.4	44.2	1.0
Arable with Hay	40.1	3.2	43.3	1.5
Mean	40.0	3.8	43.8	1.2
	Sugar Beet			
	Roots (washed): tons per acre	Sugar percent- age	Total sugar: cwt per acre	Tops: tons per acre
No dung	9.06	17.0	30.9	8.06
Dung in 1950	10.18	17.0	34.7	8.44
Increase	1.12	0.0	3.8	0.38
Previous Rotation				
Ley	10.46	17.0	35.6	9.06
Arable with Sugar beet	8.78	17.1	30.0	7.44
Mean	9.62	17.0	32.8	8.25



54/Be/1.5

Test Crops

		Previous Rotation				Mean
		Ley	Lucerne	Arable with hay	Arable with sugar beet	
Potatoes, Total tubers: tons per acre						
No dung	( $\pm 0.715$ ) <sup>*</sup>	11.90	8.90	8.34	5.78	8.73
Dung in 1954		14.14	14.86	13.02	10.44	13.12
Mean	( $\pm 0.564$ )	13.02	11.88	10.68	8.11	10.92
Increase	( $\pm 0.881$ )	2.24	5.96	4.68	4.66	4.39 ( $\pm 0.440$ )
Potatoes, Percentage ware						
No dung		94.5	95.2	91.2	88.0	92.2
Dung in 1954		95.4	95.7	95.8	94.4	95.3
Mean		95.0	95.5	93.5	91.2	93.8
Increase		0.9	0.5	4.6	6.4	3.1
Barley, Grain: cwt per acre						
No dung	( $\pm 2.06$ ) <sup>*</sup>	33.9	32.0	30.1	25.5	30.4
Dung in 1953		35.6	32.2	33.6	31.8	33.3
Mean	( $\pm 1.77$ )	34.7	32.1	31.9	28.6	31.8
Increase	( $\pm 2.12$ )	1.7	0.2	3.5	6.3	2.9 ( $\pm 1.06$ )
Barley, Straw: cwt per acre						
No dung		42.1	35.2	30.5	27.4	33.8
Dung in 1953		48.9	44.4	41.9	39.7	43.7
Mean		45.5	39.8	36.2	33.5	38.8
Increase		6.8	9.2	11.4	12.3	9.9

<sup>\*</sup> for use in comparisons other than vertical.



54/Bf/1.1

#### WOBURN MARKET GARDEN EXPERIMENT

Organic manures and N - Lansome 1954 the 13th year.

The present cropping comprises two series, each carrying in turn the crops of a two course rotation: 1st year - Globe beet followed by spring cabbages; 2nd year - Leeks.

Note: The results for the spring cabbages 1954-55 will be included in the 1955 report.

System of replication (each series): 4 randomized blocks of 10 plots each, certain interactions being confounded with block differences.

Area of each plot: 0.0125 acre.

Treatments applied to each crop:

Organic manures: Dung; sewage sludge compost; sewage sludge (West Middlesex); vegetable compost, each at 10 and 20 tons per acre.

N (applied as nitrochalk): None; 0.3 cwt per acre on plots receiving organic manure. None; 0.3; 0.6; 0.9 cwt per acre on plots not receiving organic manure. The last two rates are applied in two equal dressings.

Basal manuring per acre to each crop: 0.3 cwt  $P_2O_5$ ; 0.3 cwt  $K_2O$ , applied as granular fertilizer (13%  $P_2O_5$ ; 13%  $K_2O$ ).

Cultivations, etc.:

Globe beet. Organic manures applied and ploughed in: Mar 31. Nitrochalk and basal fertilizers applied: May 10. Seed drilled at 13 lb per acre: May 11. Singled: June 16. Second dressing of nitrochalk: June 25. Sprayed with systemic insecticide: June 21. Harvested: Aug 13 - Sept 21. Variety: Detroit.

Leeks. Organic manures applied and ploughed in: July 22. Nitrochalk and basal fertilizers applied: July 27. Planted: Aug 3. Second dressing of nitrochalk applied: Sept 22. Harvested: Mar 8 - 31, 1955. Variety: Musselburgh.

Standard errors per plot.

Globe beet, saleable bulbs: 1.15 tons per acre or 17.3% (17 d.f.)  
Leeks, saleable produce: 0.316 tons per acre or 10.8% (17 d.f.)



54/Bf/1.2

Summary of Results

Globe Beet

Organic manures	Level of manuring tons per acre	N: cwt per acre				Mean
		None	0.3	0.6	0.9	
Saleable bulbs: tons per acre						
			(±0.813)			(±0.575)
None		0.59	3.17	3.63	0.69	1.88*
Dung	10	2.86	8.47			5.66
	20	12.64	12.69			12.67
Sludge compost	10	3.35	6.93			5.14
	20	6.85	8.46			7.66
Sludge	10	6.56	6.98			6.77
	20	8.18	9.97			9.08
Vegetable compost	10	4.92	6.20			5.56
	20	6.46	13.15			9.81
Mean (±0.288)		6.48 <sup>+</sup>	9.11 <sup>+</sup>			6.64

Total produce: tons per acre

None		2.16	6.03	6.91	2.28	4.09*
Dung	10	5.23	13.52			9.37
	20	19.80	19.73			19.77
Sludge compost	10	6.05	11.32			8.68
	20	11.80	13.87			12.84
Sludge	10	11.67	12.28			11.97
	20	13.85	16.64			15.25
Vegetable compost	10	8.20	9.92			9.06
	20	10.42	20.56			15.49
Mean		10.88 <sup>+</sup>	14.73 <sup>+</sup>			11.11

Plant number: thousands per acre

None		46.3	64.8	63.2	37.9	55.6*
Dung	10	49.8	69.5			59.7
	20	82.8	75.2			79.0
Sludge compost	10	52.9	75.8			64.4
	20	72.8	66.4			69.6
Sludge	10	66.6	75.6			71.1
	20	66.9	73.0			70.0
Vegetable compost	10	59.0	64.6			61.8
	20	59.4	83.2			71.3
Mean		63.8 <sup>+</sup>	72.9 <sup>+</sup>			65.3

\*Mean over None and 0.3 cwt N per acre only.

<sup>+</sup>Excluding 'No organics'.



54/Bf/1.3

		Leeks				
Organic manures	Level of manuring tons per acre	N: cwt per acre				Mean
		None	0.3	0.6	0.9	
Saleable produce: tons per acre						
			(±0.224)			(±0.158)
None		1.09	2.08	2.47	2.61	1.58*
Dung	10	2.87	3.47			3.17
	20	3.06	3.69			3.37
Sludge compost	10	2.85	3.46			3.16
	20	3.64	3.19			3.42
Sludge	10	2.92	2.83			2.87
	20	2.87	3.29			3.08
Vegetable compost	10	2.71	3.06			2.89
	20	3.05	3.54			3.30
Mean (±0.079)		3.00 <sup>+</sup>	3.32 <sup>+</sup>			2.94
Percentage saleable (by number)						
None		75.3	89.2	94.9	93.4	82.2*
Dung	10	97.0	98.1			97.6
	20	97.0	96.7			96.8
Sludge compost	10	95.7	98.7			97.2
	20	99.6	95.3			97.4
Sludge	10	97.0	95.4			96.2
	20	96.0	98.7			97.3
Vegetable compost	10	93.8	95.7			94.8
	20	96.1	98.1			97.1
Mean		96.5 <sup>+</sup>	97.1 <sup>+</sup>			95.1

\*Mean over None and 0.3 cwt N per acre only.

<sup>+</sup>Excluding 'No organics'.



54/Bg/1.1

IRRIGATION EXPERIMENT

The 4th year

The effects of irrigation and nitrogen - Woburn Butt Close 1954.

The cropping comprises four series; three of these in turn carry the crops of a 3-course rotation:-

- 1st year: Potatoes
- 2nd year: Sugar beet
- 3rd year: Barley

The fourth series (formerly 3-year ley) was resown to 3-year S37 in 1954. Cocksfoot ley for cutting.

In 1954 maincrop potatoes replaced early potatoes followed by cabbages.

System of replication: 3 randomised blocks of 4 plots each, plots being split into two for the application of nitrogen.

Area of each sub plot: Cut grass - 0.0264, remainder - 0.0278 acre.

Area harvested: Cut grass - 0.0165, potatoes - 0.0155, sugar beet - 0.0176, barley - 0.0168 acre.

Treatments: All combinations of:-

Whole plots. Irrigation: None (0) and 3 other treatments A, B and C as specified below

N.B: On potatoes 0 = B. On sugar beet, 0 = B and A = C. On cut grass, 0 plots received .50" irrigation.

Sub plots. Nitrogen: 2 levels applied to crop as below.

Rainfall and Irrigation: inches

Week- ending	Rainfall	Potatoes			Sugar beet	Barley			Cut Grass			
		A	B	C	A & C	A	B	C	0	A	B	C
May 17	0.04					.50		.50	.50	.50	.50	.50
24	0.18				.50	.50		.50		.25	.25	.25
31	1.40	.50		.50							.25	.50
June 7	1.08											
14	1.73											
21	0.01											
28	0.08			.50							.50	.67
July 5	0.26	.85		.75	.75		.75	.75		.25	.25	.51
12	0.26	.40								.17		
19	0.90			.20							.27	.51
26	0.75	.42		.24								
Total	6.69	2.17		2.19	1.25	1.00	.75	1.75	.50	1.17	2.02	2.94



54/Bg/1.2

Levels of nitrogen (in addition to N in basal dressing):

N cwt per acre as nitrochalk

Potatoes	None; 0.5
Sugar beet	None; 0.4
Barley	None; 0.2
Cut grass	0.15; 0.3 (in spring and after each cut)

Basal dressings: cwt per acre

As compound fertilizer

	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	Salt	Dung
Potatoes	0.5	0.5	0.75	-	15 tons
Sugar beet	0.4	0.4	0.6	5	-
Barley	0.2	0.2	0.3	-	-
Cut grass (yearly)	None	0.6	1.2	-	-

Cultivations, etc.:

Potatoes. Ploughed: Aug 25 and again Oct 21, 1953. F.Y.M. applied: Mar 23, 1954. Ploughed: Mar 23. Fertilizers applied: Mar 30. Potatoes planted by machine: Apr 6. Earthed up: June 15. Sprayed with copper fungicide, medium volume, 5 lb in 40 gallons per acre: July 30. At low volume, 5 lb in 10 gallons: Aug 16 and again Aug 27. Sprayed with 20% sulphuric acid: Sept 23. Lifted: Oct 7. Variety: Majestic.

Sugar beet. Ploughed: Mar 2. Fertilizers applied: Mar 30. Seed drilled at 18 lb per acre: Apr 5. Singled: May 28. Sprayed with Parathion, low volume: June 18. Lifted: Oct 27. Variety: Klein E.

Barley. Ploughed: Dec 24, 1953. Fertilizers applied: Mar 15, 1954. Seed drilled at 3 bushels per acre: Mar 16. Harvested: Aug 16. Variety: Herta.

Cut grass. Ploughed: Nov 24, 1953. Basal fertilizer and nitrochalk applied: Apr 6, 1954. Grass seed sown at 28 lb per acre: Apr 7. Sprayed MCPA, low volume, 2 pints per acre: June 19. Cut: June 16, July 15 (high N plots), July 23 (low N plots), July 30 (high N plots), Aug 17, Sept 10, Nov 4. Nitrochalk applied after each cut except the last. Variety: Cocksfoot S37.



54/Bg/1.3

Standard errors per plot:

Potatoes,	Total tubers,	whole plot:	0.731 tons per acre or 4.7%	(7 d.f.)
		sub plot:	1.069 tons per acre or 6.9%	(9 d.f.)
Sugar beet,	Total sugar,	whole plot:	4.83 cwt per acre or 10.9%	(8 d.f.)
		sub plot:	4.25 cwt per acre or 9.6%	(10 d.f.)
	Tops,	whole plot:	1.21 tons per acre or 12.5%	(8 d.f.)
		sub plot:	1.16 tons per acre or 12.0%	(10 d.f.)
Barley,	Grain,	whole plot:	1.08 cwt per acre or 3.1%	(6 d.f.)
		sub plot:	1.75 cwt per acre or 4.9%	(8 d.f.)
Cut grass	Hay (85% D.M.),	whole plot:	6.89 cwt per acre or 14.3%	(6 d.f.)
(total of 6 cuts)		sub plot:	5.01 cwt per acre or 10.4%	(8 d.f.)

Summary of Results

cwt N per acre	O & B	Irrigation A	C	Mean
Potatoes, total tubers: tons per acre				
	(±0.429) <sup>**</sup>	(±0.607) <sup>**</sup>		
0.0	14.06	13.42	13.25	13.70
0.5	17.88	16.94	16.95	17.41
Mean	(±0.422) 15.97 <sup>(1)</sup>	15.18	15.10	15.55
Difference	(±0.873) 3.82 <sup>(2)</sup>	3.52	3.70	3.71 (±0.436)
Potatoes, percentage ware				
0.0	87.0	83.7	85.5	85.8
0.5	91.2	86.2	88.4	89.8
Mean	89.1	86.0	87.0	87.8
Difference	4.2	4.5	2.9	4.0

(1) ±0.298

(2) ±0.617

<sup>\*\*</sup> for use in comparisons other than vertical.



54/Bg/1.4

cwt N per acre	Irrigation				Mean
	0	A	B	C	
Barley, grain: cwt per acre					
( $\pm 0.949$ ) <sup>‡</sup>					
0.0	31.3	35.7	35.6	33.1	33.9
0.2	36.8	36.3	38.6	36.0	36.9
Mean ( $\pm 0.625$ )	34.0	36.0	37.1	34.6	35.4
Difference ( $\pm 1.428$ )	5.5	0.6	3.0	2.9	3.0 ( $\pm 0.714$ )

Barley, straw: cwt per acre					
0.0	25.5	29.3	28.4	28.3	27.9
0.2	35.7	35.0	33.8	36.3	35.2
Mean	30.6	32.1	31.1	32.3	31.5
Difference	10.2	5.7	5.4	8.0	7.3

Cut grass, hay (at 85% D.M.) 6 cuts: cwt per acre					
( $\pm 4.47$ ) <sup>‡</sup>					
0.15	37.9	37.5	35.0	38.4	37.2
0.30	61.6	56.0	59.5	60.8	59.5
Mean ( $\pm 3.98$ )	49.8	46.7	47.2	49.6	48.3
Difference ( $\pm 4.09$ )	23.7	18.5	24.5	22.4	22.3 ( $\pm 2.04$ )

cwt N per acre	Irrigation		Mean
	C & B	A & C	
Sugar beet, roots (washed): tons per acre			
0.0	12.26	12.46	12.36
0.4	13.56	13.30	13.43
Mean	12.91	12.88	12.90
Difference	+1.30	+0.84	+1.07

Sugar beet, sugar percentage			
0.0	17.1	17.2	17.1
0.4	17.2	17.3	17.2
Mean	17.1	17.2	17.2
Difference	0.1	0.1	0.1

<sup>‡</sup> for use in comparisons other than vertical



54/Bg/1.5

cwt N per acre	Irrigation		Mean
	C & B	A & C	
Sugar beet, total sugar: cwt per acre			
	( $\pm 2.32$ ) <sup>*</sup>		
0.0	41.9	42.8	42.4
0.4	46.6	46.1	46.4
Mean	( $\pm 1.96$ )	44.3	44.5
Difference	( $\pm 2.46$ )	-4.7	3.3
			-4.0 ( $\pm 1.74$ )

Sugar beet, tops: tons per acre			
	( $\pm 0.597$ ) <sup>*</sup>		
0.0	9.11	9.23	9.17
0.4	10.07	10.28	10.17
Mean	( $\pm 0.494$ )	9.59	9.75
Difference	( $\pm 0.672$ )	-0.96	1.05
			-1.00 ( $\pm 0.475$ )

Sugar beet, noxious nitrogen: mg per 100 g.			
0.0	25	25	25
0.4	30	25	25
Mean	25	25	25
Difference	5	0	0

<sup>\*</sup>for use in comparisons other than vertical.