


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1954

[Full Table of Content](#)



54/R/BA/3 and 54/W/BA/3 Six-course Rotation

Rothamsted Research

Rothamsted Research (1955) *54/R/BA/3 and 54/W/BA/3 Six-course Rotation* ; Yields Of The Field Experiments 1954, pp 23 - 25 - DOI: <https://doi.org/10.23637/ERADOC-1-184>

54/Ba/3.1

SIX COURSE ROTATION EXPERIMENT

The 25th year

Seasonal effects of fertilizers - Rothamsted Long Hoos IV and Woburn Stackyard 1954.

For details of treatments, rotation etc., see "Results of the Field Experiments 1939-47" Vol.I, Section Ba/4.

Area of each plot: Rothamsted - 0.0250 acre. Woburn - 0.0266 acre.

Cultivations, etc.:

Rothamsted

Sugar beet.

Ploughed: Aug 21, 1953, Oct 20 and Dec 1. Fertilizers applied: Mar 26, 1954. Seed drilled at 18 lb per acre: Mar 30. Singled: May 31. Lifted: Nov 17. Variety: Klein E.

Barley.

Ploughed: Nov 27, 1953. Ground chalk applied at 22 cwt per acre: Dec 4. Fertilizers applied, seed drilled at 3 bushels per acre: Mar 15, 1954. Harvested: Aug 25. Variety: Plumage Archer.

Clover.

Seed undersown in barley at 40 lb per acre: Apr 23, 1953. Autumn fertilizers applied: Nov 2. Sulphate of ammonia applied: Apr 30, 1954. Cut: July 13. Variety: Late flowering Montgomery Red.

Wheat.

Ploughed: July 10, 1953 and again Sept 17. Autumn fertilizers applied: Oct 9. Seed drilled at 3 bushels per acre: Oct 17. Sulphate of ammonia applied: Apr 30, 1954. Harvested: Aug 26. Variety: Yeoman.

Potatoes.

Ploughed: Aug 29, 1953, Oct 20 and Nov 30. Ridged: Apr 13, 1954. Fertilizers applied, potatoes planted: Apr 14. Earthed up: July 6. Sprayed (low volume) with copper fungicide paste $\frac{1}{2}$ gallon per acre: July 27 and again Aug 18. Sprayed with sulphuric acid 15% B.O.V: Sept 28. Lifted: Oct 2. Variety: Majestic.

Rye.

Ploughed: Sept 30, 1953. Ground chalk applied at 22 cwt per acre: Oct 3. Autumn fertilizers applied: Oct 10. Seed drilled at 3 bushels per acre: Oct 17. Sulphate of ammonia applied: Apr 30, 1954. Harvested: Aug 25. Variety: King II.

Note:- Clover: 6 plots were carted off in error when green. No summary of results can be shown.

54/Ba/3.2

Woburn

Sugar beet.

Ploughed: Aug 20, 1953, Nov 3 and Jan 20, 1954. Fertilizers applied: Mar 29. Seed drilled at 18 lb per acre: Mar 30. Dusted with D.D.T; May 12. Singled: May 31. Sprayed with parathion $\frac{1}{2}$ pint per acre: June 21. Lifted: Nov 2. Variety: Klein E. Previous crop: Rye. N.B. The crop was attacked by flea beetle and mangold fly.

Barley.

Ploughed: Nov 4, 1953. 2nd ploughing: Jan 20, 1954. Fertilizers applied: Mar 11. Seed drilled at $2\frac{1}{2}$ bushels per acre: Mar 17. Harvested: Aug 28. Variety: Plumage Archer. Previous crop: Sugar beet.

Clover.

Seed undersown in barley at 40 lb per acre: Mar 26, 1953. Autumn fertilizers applied: Oct 23. Sulphate of ammonia applied: Apr 27, 1954. Cut: June 30. Variety: Montgomery Red. Previous crop: Barley. N.B. The crop was damaged by Sclerotinia and was very weedy.

Wheat.

Ploughed: July 3, 1953. 2nd ploughing: Sept 16. Autumn fertilizers applied: Oct 22. Drilled at $2\frac{1}{2}$ bushels per acre: Oct 23. Sprayed with D.N.O.C. at 6 lb per acre in 80 gallons: Apr 27. Sulphate of ammonia applied: Apr 29. Harvested: Sept 7. Variety: Squareheads Master $13/4$. Previous crop: Clover. N.B. The crop was very weedy.

Potatoes.

Ploughed: Sept 15, 1953, Nov 3 and Jan 20, 1954. Ridged, fertilizers applied, seed planted: Apr 18. Sprayed with copper fungicide: July 30. Sprayed with 15% sulphuric acid: Sept 23. Lifted: Oct 8. Variety: Majestic. Previous crop: Wheat.

Rye.

Ploughed: Oct 1, 1953. Autumn fertilizer applied, drilled seed at $2\frac{1}{2}$ bushels per acre: Oct 22. Sulphate of ammonia applied: Apr 29. Harvested: Aug 28. Variety: King II. Previous crop: Potatoes.

54/Ba/3.3

Summary of Results

Mean yields per acre and responses in yield per cwt of N, P₂O₅ and K₂O

	Rothamsted	Woburn	Rothamsted	Woburn	
		Sugar Beet, roots (washed): tons per acre		Barley, grain: cwt per acre	
Mean	8.55	4.57	31.7	29.1	
Response to: N	+2.88	+1.14	+4.6	+25.6	
P	+3.05	+4.46	+0.4	-1.9	
K	-0.08	-0.07	-1.2	-1.0	
		Sugar Beet, sugar percentage		Barley, straw: cwt per acre	
Mean	16.30	16.29	36.0	29.9	
Response to: N	+0.22	+0.56	+12.8	+40.3	
P	-1.21	+1.03	+1.1	-7.1	
K	+0.28	+0.58	-1.9	+2.8	
		Sugar Beet, total sugar: cwt per acre		Clover, hay: dry matter cwt per acre	
Mean	27.9	14.9		37.6	
Response to: N	+9.9	+4.2		-29.0	
P	+8.1	+15.7		+11.3	
K	+0.2	+0.2		+8.6	
		Sugar Beet, tops: tons per acre		Wheat, grain: cwt per acre	
Mean	10.93	4.88	26.5	14.8	
Response to: N	+5.88	+0.69	+8.0	+19.3	
P	+1.93	+5.21	-10.7	-9.6	
K	+0.02	-0.55	+5.2	+10.1	
		Sugar Beet, plant number: thousands per acre		Wheat, straw: cwt per acre	
Mean	26.0		48.2	18.6	
Response to: N	-1.1	†	+22.1	+22.7	
P	+3.1		-2.1	-9.8	
K	+1.7		-0.8	+12.7	
		Potatoes, total tubers: tons per acre		Rye, grain: cwt per acre	
Mean	6.08	4.21	32.8	26.4	
Response to: N	+3.07	+4.49	+32.2	+24.6	
P	+0.57	+1.39	-3.7	+3.3	
K	+1.32	+0.30	+0.1	-1.1	
		Potatoes, percentage ware		Rye, straw: cwt per acre	
Mean	78.9	80.5	40.2	30.9	
Response to: N	-10.7	+10.8	+23.6	+23.5	
P	-10.7	+6.6	-3.9	+8.9	
K	+6.5	+12.7	+2.0	-0.5	

* corrected to 85% Dry Matter. Mean Dry Matter %: 83.6 † not recorded.