


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1954

[Full Table of Content](#)



### 54/R/CA/2 Wheat - Eyespot, Varieties, Seed Rates and N 1st Year

#### Rothamsted Research

Rothamsted Research (1955) *54/R/CA/2 Wheat - Eyespot, Varieties, Seed Rates and N 1st Year ; Yields Of The Field Experiments 1954*, pp 65 - 66 - DOI: <https://doi.org/10.23637/ERADOC-1-184>

54/Ca/2.1

WHEAT

The effects of Crop sequences, Varieties, Seed rates and Nitrogen on the incidence of Eyespot (*Cercospora herpotrichoides*) - Long Hoos 1, 2 and 3, 1954. The 1st preliminary year.

System of replication: 8 randomized blocks of 4 plots each, the interaction VRN being confounded with block differences.

Area of each plot: 0.0212 acre. Area harvested: 0.0148 acre.

Treatments: All combinations of:-

Varieties: Holdfast; Cappelle.

Seed rates:  $1\frac{1}{2}$ ; 3 bushels per acre.

Nitrogen: 0.46; 0.93 cwt N per acre applied as nitrochalk, half in March, half in May.

Crop sequences:

During the preliminary years 1954 and 1955 four cropping systems will be set up, each on four blocks; in 1956 wheat will be taken over all 16 blocks. The crops will be:-

1954	1955	1956
Wheat	Wheat	Wheat
Wheat	Potatoes	Wheat
Potatoes	Wheat	Wheat
Beans	Potatoes	Wheat

In 1954 the 4 blocks of potatoes (Majestic) received 10 tons dung and 12 cwt compound fertilizer, (7% N, 7% P<sub>2</sub>O<sub>5</sub>, 10 $\frac{1}{2}$ % K<sub>2</sub>O) the 4 blocks of beans received 4 cwt compound fertilizer, (16% P<sub>2</sub>O<sub>5</sub>, 16% K<sub>2</sub>O).

Basal dressing, per acre: 1 cwt compound granular fertilizer (12% N, 12% P<sub>2</sub>O<sub>5</sub>, 15% K<sub>2</sub>O) combine drilled with seed. To correct some acid areas 25 cwt ground chalk and 23 cwt hydrated lime was applied to certain blocks.

Cultivations, etc.: Ploughed: Oct 6, 1953. Seed combine drilled: Oct 26. Ground chalk applied: Dec 29. Hydrated lime applied: Jan 18 - Feb 24. 1st application of nitrochalk: Mar 9. 2nd application: May 10. Sprayed with M.C.P.A., medium volume, 2 pints per acre: May 27. Combine harvested: Sept 6-7. Varieties: Holdfast and Cappelle. Previous crop: Potatoes.

Standard error per plot:

Grain (at 85% D.M.): 2.71 cwt per acre or 6.6% (18 d.f.)



54/Ga/2.2

Summary of Results

Response to	Mean	Responses to treatments					
		Variety		Seed rate		Nitrogen:	
		Hold- fast	Capp- elle	bushels per acre		cwt per acre	
				1½	3	0.46	0.93

Grain (at 85% Dry Matter): Mean yield 41.0 cwt per acre

(±0.96)

(±1.36)

Variety (Cappelle - Holdfast)	+3.4	-	-	-1.3	+8.1	+3.4	+3.4
Seed rate (3-1½)	+4.8	+0.1	+9.5	-	-	+4.7	+4.9
Nitrogen (0.93 - 0.46)	+3.2	+3.2	+3.2	+3.1	+3.3	-	-

Mean Dry Matter % as harvested: 78.0

Records of incidence of disease (Eyespot and Take-All) and counts of plant, shoot, and straw numbers were made.