


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1954

[Full Table of Content](#)



## 54/R/CD/3 Potatoes - Control of Blight

### Rothamsted Research

Rothamsted Research (1955) *54/R/CD/3 Potatoes - Control of Blight* ; Yields Of The Field Experiments 1954, pp 87 - 87 - DOI: <https://doi.org/10.23637/ERADOC-1-184>

POTATOES

Control of Blight - Great Field I 1954.

System of replication: 4 x 4 Latin Square, plots being split into 2 for determination of tractor damage.

Area of each sub plot: 0.0140 acre.

Treatments:

Whole plots: No Spray; Copper fungicide 5 lb in 40 gallons per acre sprayed twice; 100 gallons sulphuric acid, 20% B.O.V. sprayed to destroy haulms; Copper fungicide and sulphuric acid sprayed as above. The tractor used for spraying was driven over all the plots on each occasion.

Sub plots: 4 rows damaged by three passages of the tractor were compared with 4 undamaged rows.

Basal dressing, per acre: 10 tons F.Y.M. and 10 cwt compound granular fertilizer (7% N, 7% P<sub>2</sub>O<sub>5</sub>, 10½% K<sub>2</sub>O).

Cultivations, etc.: F.Y.M. applied: Jan 18. Ploughed: Jan 20 - Feb 24. Fertilizer applied: Apr 21. Ridged: Apr 22. Planted: Apr 23. Cultivated with mechanical weeder: June 1, 25, 29. Ridged: June 3 and again July 14. Fungicide treatment applied: July 30 and again Aug 16. Sulphuric acid treatment applied: Sept 13. Lifted: Oct 23. Variety: Majestic S.S. Previous crop: Wheat.

Standard errors per plot: Total tubers.

Whole plot: 0.594 tons per acre or 5.8% (6 d.f.)

Sub plot: 0.969 tons per acre or 9.5% (12 d.f.)

Note: Blight counts were made and are available.

Summary of Results

	Spray			Copper fungicide and Sulphuric Acid	Mean
	None	Copper fungicide	Sulphuric Acid		
Total tubers: tons per acre (±0.454)*					
Undamaged rows	9.30	12.56	9.56	12.52	10.99
Damaged rows	8.01	9.99	9.12	10.91	9.51
Mean (±0.297)	8.66	11.27	9.34	11.71	10.25
Difference (±0.685)	-1.29	-2.57	-0.44	-1.61	-1.48 (±0.343)
Percentage ware					
Undamaged rows	86.2	87.0	86.8	82.4	85.6

\*for use in comparisons other than vertical.