

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1955

[Full Table of Content](#)



Default Title

Rothamsted Research

Rothamsted Research (1956) *Default Title* ; Yields Of The Field Experiments 1955, pp 1 - 106 - **DOI:** <https://doi.org/10.23637/ERADOC-1-175>

Rothamsted Experimental Station
Harpenden
LAWES AGRICULTURAL TRUST

RESULTS
OF THE
FIELD
EXPERIMENTS
1955

Rothamsted Experimental Station

Harpenden

Lawes Agricultural Trust

RESULTS

of the

FIELD

EXPERIMENTS

1955

The summaries given in this report are similar to those contained in the appendices to the Annual Reports of the Station before the war. This year's report includes only experiments conducted at Rothamsted and Woburn. The design and supervision of these experiments are the responsibility of the Field Plots Committee (present members: F.Yates (Chairman), H.V.Garner (Secretary), F.C.Bawden, G.W.Cooke, H.H.Mann, J.R. Moffatt, R.G.Warren, D.J.Watson).

Price 5/-

INDEX 1955

Classical Experiments*

| | | |
|---------------------------|-------------------------|-----|
| Broadbalk | Wheat | A/1 |
| Hoosfield | Barley | A/2 |
| Hoosfield | Wheat after fallow | A/3 |
| Agdell | Rotation | A/3 |
| Barnfield | Mangolds and sugar beet | A/4 |
| Park Grass | Hay | A/5 |
| Hoosfield Exhaustion Land | Barley | A/6 |
| Stackyard Woburn | Wheat | A/7 |
| Stackyard Woburn | Barley | A/8 |

Long Term Experiments

| | | |
|---------------------------|-----------------------|------|
| 3-Course Rotation | Rothamsted | Ba/1 |
| 4-Course Rotation | Rothamsted | Ba/2 |
| 6-Course Rotation | Rothamsted and Woburn | Ba/3 |
| Deep Cultivation Rotation | Rothamsted | Bb/1 |
| Ley and Arable Rotations | Rothamsted | Bc/1 |
| Green Manuring | Woburn | Bd/1 |
| Ley and Arable Rotations | Woburn | Be/1 |
| Market Garden Soil | Woburn | Bf/1 |
| Irrigation | Woburn | Bg/1 |

Short Term Experiments*

| | | |
|--------------|--|------|
| Winter Wheat | Eyespot Rotation, Varieties, Seed rates and N - 2nd year. | Ca/1 |
| Winter Wheat | Control of Bulb Fly (Insecticides). | Ca/2 |
| Winter Wheat | Control of Bulb Fly (Seed rates). | Ca/3 |
| Winter Wheat | Varieties, Seed rates, levels and times of N - Woburn. | Ca/4 |
| Spring Wheat | Residuals of Dung, N, P and K. | Ca/5 |
| Spring Wheat | Levels and times of N - Rothamsted and Woburn. | Ca/6 |
| Spring Wheat | Varieties and N. | Ca/7 |
| Barley | Seed rates and levels of N. | Cb/1 |
| Barley | Levels and times of N - Rothamsted and Woburn. | Cb/2 |
| Barley | Varieties and N (Growth Study). | Cb/3 |
| Barley | Residuals of P and K. | Cb/4 |
| Barley | Varieties and N. | Cb/5 |
| Spring Beans | Control of aphids (spraying and times of sowing). | Cc/1 |
| Spring Beans | Control of weeds (spraying and cultivations). | Cc/2 |
| Spring Beans | Setting of flowers (Hormone sprays). | Cc/3 |

*At Rothamsted unless otherwise stated.

55/A/1.1

WHEAT - BROADBALK 1955

The 112th year

For history, details of treatments etc. see "Results of the Field Experiments" 1939-47, Vol. I, Section A/1.

Commencing in autumn 1954 a scheme of chalking was introduced as follows:- the half of section V nearest the drain received a single application of 5 tons of calcium carbonate per acre as ground chalk for this year only. On all plots receiving either ammonium sulphate or castor meal an annual compensating dressing of ground chalk will be applied at the following rates: 100 lb of calcium carbonate for each 14 lb of N applied as ammonium sulphate; 50 lb of calcium carbonate for each 14 lb of N applied as castor meal. In autumn 1954 this compensating dressing was given at double rate; henceforward it will be given at single rate on the stubble.

No compensating dressing is given to the section under fallow or to plots 17 and 18 in their 'no nitrogen' years.

In 1955 Section I was harvested in two separate sub-sections IA and IB. Starting in 1956 IA (nearest the wilderness) will carry wheat without fallow; IB will continue in the following cycle.

Cultivations, etc.:

Cropped sections. Ground chalk applied to lower half of Section V: Sept 28-Oct 21, 1954. Dung applied, ploughed all plots: Oct 1. Ground chalk applied to ammonium sulphate and castor meal plots: Oct 20. Autumn fertilizers applied: Nov 5. Seed drilled at 3 bushels per acre: Dec 6. Spring fertilizers applied: May 16, 1955. Second dressing of nitrate of soda applied to plot 16: June 1. Harvested: Aug 19. Variety: Squareheads Master 13/4.

Fallow section. Ploughed: Oct 1, 1954, April 21 and July 25, 1955.

55/A/1.2

Summary of Results

| Section Years after fallow | Grain (at 85% dry matter): cwt per acre | | | | | | Mean |
|-------------------------------------|---|--------------|------------|------|------|------|------|
| | IV | VA | VB | II | IA | IB | |
| | 1 | Unlimed 2 | Limed 2 | 3 | 4 | 4 | |
| 2A | 36.5 | 27.6 | 30.9 | 25.8 | 23.1 | 17.2 | 28.0 |
| 2B | 36.7 | 31.7 | 25.3 | 21.3 | 18.8 | 25.7 | 27.6 |
| 3 | 19.7 | 11.1 | 10.4 | 11.2 | 9.5 | 10.7 | 13.1 |
| 5 | 20.9 | 8.0 | 9.9 | 12.7 | 16.5 | 11.4 | 13.9 |
| 6 | 28.2 | 12.9 | 13.1 | 16.4 | 15.1 | 14.5 | 18.2 |
| 7 | 31.5 | 19.4 | 17.5 | 19.5 | 18.8 | 20.5 | 22.4 |
| 8 | 33.3 | 29.4 | 26.1 | 23.8 | 22.2 | 26.0 | 27.5 |
| 9 | 25.3 | 19.3 | 17.0 | 18.9 | 8.9 | 15.6 | 19.1 |
| 10 | 17.1 | 10.6 | 19.1 | 15.0 | 15.2 | 17.0 | 15.8 |
| 11 | 23.1 | 19.9 | 20.6 | 17.0 | 19.4 | 17.6 | 19.7 |
| 12 | 26.9 | 16.9 | 19.7 | 20.9 | 18.7 | 19.3 | 21.3 |
| 13 | 28.3 | 19.0 | 15.0 | 20.9 | 19.0 | 20.3 | 21.5 |
| 14 | 25.7 | 20.7 | 12.9 | 18.9 | 18.9 | 18.2 | 20.0 |
| 15 | 24.0 | 12.6 | 9.5 | 16.2 | 17.8 | 14.5 | 16.7 |
| 16 | 29.1 | 27.4 | 21.2 | 25.3 | 25.2 | 24.0 | 25.8 |
| 17 | 18.0 | 11.2 | 10.0 | 10.1 | 6.7 | 5.7 | 11.4 |
| 18 | 29.0 | 16.0 | 16.4 | 22.6 | 21.1 | 22.6 | 22.4 |
| 19 | 23.5 | 15.6 | 11.7 | 14.5 | 14.8 | 15.4 | 16.7 |
| 20 | - | - | - | 17.4 | 15.7 | 18.1 | 17.3 |

Mean dry matter % as threshed: 84.8

55/A/1.3

| Section Years after fallow | Straw (at 85% dry matter): cwt per acre | | | | | | Mean |
|-------------------------------------|---|--------------|------------|------|------|------|------|
| | IV | VA | VB | II | IA | IB | |
| | 1 | Unlimed 2 | Limed 2 | 3 | 4 | 4 | |
| 2A | 52.6 | 43.0 | 40.3 | 40.4 | 39.3 | 30.8 | 42.4 |
| 2B | 52.1 | 45.8 | 43.8 | 45.5 | 40.5 | 41.8 | 46.1 |
| 3 | 24.7 | 15.2 | 17.8 | 17.9 | 22.4 | 18.9 | 19.7 |
| 5 | 34.6 | 13.4 | 14.8 | 25.5 | 40.7 | 21.4 | 25.3 |
| 6 | 44.1 | 18.5 | 21.3 | 27.2 | 34.5 | 26.9 | 30.1 |
| 7 | 49.2 | 33.9 | 30.4 | 34.8 | 31.9 | 32.3 | 37.2 |
| 8 | 55.0 | 44.8 | 42.5 | 41.7 | 39.1 | 38.6 | 45.0 |
| 9 | 41.0 | 26.5 | 27.0 | 28.0 | 27.2 | 29.7 | 31.2 |
| 10 | 25.2 | 19.7 | 28.4 | 21.0 | 22.9 | 22.7 | 23.3 |
| 11 | 34.3 | 32.2 | 28.6 | 26.2 | 32.2 | 25.7 | 29.8 |
| 12 | 40.8 | 29.2 | 26.9 | 30.2 | 36.0 | 28.6 | 32.5 |
| 13 | 47.8 | 31.6 | 26.9 | 33.0 | 31.8 | 34.0 | 35.9 |
| 14 | 40.9 | 28.0 | 22.8 | 27.3 | 31.8 | 26.8 | 30.5 |
| 15 | 43.9 | 19.7 | 17.9 | 26.9 | 33.8 | 30.4 | 30.1 |
| 16 | 46.0 | 40.5 | 35.8 | 37.0 | 39.1 | 35.6 | 39.6 |
| 17 | 27.8 | 13.3 | 16.2 | 13.7 | 14.6 | 11.3 | 17.3 |
| 18 | 43.8 | 28.4 | 32.3 | 36.3 | 39.8 | 35.1 | 36.7 |
| 19 | 36.8 | 27.4 | 26.6 | 22.7 | 26.8 | 24.3 | 28.0 |
| 20 | - | - | - | 23.7 | 28.4 | 29.9 | 26.9 |

Mean dry matter % as threshed: 85.2

55/A/2.1

BARLEY - HOOSFIELD 1955

The 104th year

For history, details of treatments, etc., see "Results of the Field Experiments" 1939-47, Vol. I, Section A/2.

In the autumn of 1954 a system of chalking was started as follows:- strips 3 and 4, including plots 50 and 5A, receive a dressing of 5 tons of calcium carbonate per acre as ground chalk in 1954-55 only. Plots receiving ammonium sulphate or castor meal have a compensating dressing of ground chalk at the following rates: 100 lb calcium carbonate for each 14 lb of N applied as ammonium sulphate; 50 lb calcium carbonate for each 14 lb of N given as castor meal. The compensating dressings, at five times the annual rate, are given once every five years commencing 1955.

Cultivations, etc.: Ploughed: Sept 27, 1954. Part of ground chalk applied to strips 3 and 4: Nov 6-18. Dung applied: Nov 15. Ploughed: Dec 27. Remainder of ground chalk applied to strips 3 and 4, and compensating dressing applied to series A and C: Mar 17, 1955. Fertilizers applied: Apr 5. Seed drilled at 3 bushels per acre: Apr 7. Sprayed with 8 lb D.N.C. (active substance) in 80 gallons per acre: May 24. Cut and discarded areas of plots in series N, AA, AAS, C and plots 5A and 50, leaving 16 rows per plot: July 25. Harvested: Aug 18. Variety: Plumage Archer.

55/A/2.2

Summary of Results

| Plot | Grain (at 85% dry matter): cwt per acre | Straw (at 85% dry matter): cwt per acre |
|--------------------|--|--|
| 1 O | 4.4 | 4.5 |
| 2 O | 9.1 | 7.6 |
| 3 O | 4.2 | 4.8 |
| 4 O | 7.3 | 6.9 |
| 5 O | 4.0 | 4.8 |
| 1 A | 7.0 | 7.9 |
| 2 A | 21.5 | 19.4 |
| 3 A | 8.4 | 10.1 |
| 4 A | 16.0 | 15.0 |
| 5 A | 19.0 | 19.9 |
| 1 AA | 9.9 | 10.2 |
| 2 AA | 25.0 | 23.8 |
| 3 AA | 7.6 | 8.5 |
| 4 AA | 22.4 | 22.6 |
| 1 AAS | 10.5 | 10.6 |
| 2 AAS | 25.0 | 22.6 |
| 3 AAS | 15.6 | 16.9 |
| 4 AAS | 26.7 | 24.3 |
| 1 C | 15.2 | 14.5 |
| 2 C | 20.1 | 18.1 |
| 3 C | 11.8 | 10.5 |
| 4 C | 17.7 | 16.0 |
| 7 - 1 | 10.1 | 9.2 |
| 7 - 2 | 27.7 | 25.4 |
| 6 - 1 | 4.6 | 5.0 |
| 6 - 2 | 8.0 | 7.5 |
| 1 N | 10.6 | 11.1 |
| 2 N | 17.1 | 14.5 |
| Mean dry matter %: | 84.9 | 84.3 |

55/A/3

WHEAT AFTER FALLOW - HOOSFIELD 1955

Without manure 1851 and since

For history, details of treatments etc. see "Results of the Field Experiments" 1939-47 Vol.I, Section A/3.

Cultivations, etc.:

Cropped plots. Ploughed: Sept 25, 1954. Seed drilled at 3 bushels per acre: Dec 7. Harvested: Aug 20, 1955. Variety: Squareheads Master 13/4.

Fallowed plots. Ploughed: Sept 25, 1954 and Sept 28, 1955.

Summary of Results

Mean yields: cwt per acre

| Plot | B3 | B4 | B1 | |
|------------------------|------|------|------|------|
| No. of years of fallow | 1 | 1 | 3 | Mean |
| Grain | 9.4 | 10.2 | 11.2 | 10.3 |
| Straw | 12.9 | 12.3 | 15.3 | 13.5 |

Mean dry matter % as threshed, Grain: 85.3
Straw: 86.2

SPRING WHEAT - AGDELL 1955

For history, details of treatments etc., see "Results of the Field Experiments" 1939-47 Vol.I, Section A/4, and 1954, page 54/A/3. In 1955 spring wheat was grown with uniform treatment, manured with nitrogen only. Yields were not recorded. Detailed sampling both of crops and soils was carried out.

Cultivations, etc.:

Ploughed: Nov 4, 1954. Seed drilled at 2 bushels per acre: Apr 4, 1955. Basal dressing of nitrochalk, 4 cwt per acre applied: Apr 5. Sprayed with MCPA at 2½ pts. in 18 gallons per acre: May 23. Combine harvested: Sept 2. Variety: Koga II.

55/A/4.1

MANGOLDS AND SUGAR BEET - BARNFIELD 1955

The 80th and 10th years

For history, details of treatments etc., see "Results of the Field Experiments 1939-47" Vol. 1, Section A/5.

Because of a serious infestation of couch grass (Agropyron repens) it was decided to apply trichloroacetic acid to the following plots: all the O series, 1N. In 1955 half of each plot was sprayed at half the standard rate. No damage to the crops was noted, and the halves were not harvested separately.

Cultivations, etc.: Dung applied: Dec 30, 1954. Ploughed: Jan 3, 1955. Certain plots sprayed with trichloroacetic acid, sodium salt: at 20 lb. in 40 gallons per acre: Apr 9. Fertilizers applied: Apr 22. Seed drilled, Mangolds - 9 lb; Sugar beet - 16 lb per acre. Calcium chloride applied to 4NB only: June 2. Sprayed with miscible D.D.T. 3 pints per acre: June 11. Singling started: July 19. Top dressings applied; nitrate of soda - July 20, sulphate of ammonia - July 22. Lifting started: Mangolds, - Nov 13, Sugar beet - Nov 16. Varieties Mangolds - Yellow Globe, Sugar beet - Klein E.

Erratum to "Results of the Field Experiments" 1951, p. 51/A/4.2.
Sugar Beet: Tops. Cross Dressing - 0. Yields for strips 4-8 should read "1.65, 1.49, 1.57, 1.63, 1.79", NOT "6.61, ..., 7.14"

Summary of Results

| Strip | Cross Dressing | | | | |
|--|----------------|-----------------------|-------|------|-------|
| | O | N | A | AC | C |
| Mangolds, Roots: tons per acre | | | | | |
| 1 | 3.55 | 11.93 | 10.20 | 7.44 | 10.34 |
| 2 | 5.04 | 9.95 | 7.98 | 7.95 | 10.26 |
| 4 | 1.75 | (a) 7.34 (b) 7.29* | 5.38 | 7.95 | 8.04 |
| 5 | 1.03 | 6.01 | 2.50 | 3.99 | 6.07 |
| 6 | 1.01 | 4.30 | 4.67 | 7.98 | 8.72 |
| 7 | 0.89 | 7.22 | 6.97 | 8.18 | 9.38 |
| 8 | 0.44 | 2.22 | 2.47 | 4.32 | 4.05 |
| 9 | 5.66 | | | | |
| Mangolds, Leaves: tons per acre | | | | | |
| 1 | 1.47 | 3.64 | 3.27 | 3.00 | 2.69 |
| 2 | 1.71 | 3.00 | 3.10 | 3.42 | 2.20 |
| 4 | 0.94 | (a) 3.40 (b) 3.13* | 2.05 | 2.86 | 2.00 |
| 5 | 0.74 | 3.44 | 1.81 | 2.03 | 2.25 |
| 6 | 0.81 | 2.81 | 1.88 | 2.54 | 1.81 |
| 7 | 0.72 | 3.96 | 2.49 | 3.69 | 2.54 |
| 8 | 0.60 | 1.66 | 2.34 | 2.91 | 2.64 |
| 9 | 3.42 | | | | |
| Mangolds, Plant Number: thousands per acre | | | | | |
| 1 | 19.8 | 23.2 | 21.4 | 16.2 | 22.0 |
| 2 | 22.2 | 21.5 | 19.8 | 17.6 | 21.4 |
| 4 | 22.8 | (a) 24.7 (b) 23.5* | 19.0 | 17.8 | 21.3 |
| 5 | 22.6 | 24.1 | 20.3 | 16.8 | 22.0 |
| 6 | 22.1 | 22.6 | 21.5 | 19.2 | 22.5 |
| 7 | 21.3 | 23.1 | 21.5 | 18.9 | 21.9 |
| 8 | 21.0 | 21.2 | 20.2 | 20.9 | 23.0 |
| 9 | 22.5 | | | | |

*No nitrate of soda. Nitrogen applied as calcium and potassium nitrates.

55/A/4.3

| Strip | Cross Dressing | | | | |
|--|----------------|-----------|------|------|------|
| | O | N | A | AC | C |
| Sugar Beet, Roots (washed): tons per acre | | | | | |
| 1 | 2.67 | 7.83 | 7.96 | 6.78 | 8.46 |
| 2 | 3.03 | 4.78 | 3.02 | 2.86 | 6.92 |
| 4 | 1.32 | (b) 4.67* | 3.33 | 6.56 | 6.59 |
| 5 | 1.31 | 3.55 | 2.82 | 4.37 | 6.07 |
| 6 | 1.27 | 3.10 | 3.29 | 6.14 | 6.66 |
| 7 | 1.04 | 4.46 | 4.24 | 6.61 | 5.14 |
| 8 | 0.90 | 3.04 | 2.58 | 4.76 | 4.04 |
| 9 | 2.94 | | | | |
| Sugar Beet, Tops: tons per acre | | | | | |
| 1 | 2.39 | 9.28 | 8.94 | 5.96 | 4.93 |
| 2 | 3.32 | 7.03 | 4.54 | 3.91 | 5.42 |
| 4 | 1.15 | (b) 5.67* | 4.40 | 5.52 | 4.69 |
| 5 | 1.06 | 6.94 | 3.61 | 6.30 | 5.52 |
| 6 | 1.27 | 4.93 | 4.25 | 6.55 | 5.42 |
| 7 | 1.05 | 6.74 | 4.20 | 8.40 | 4.25 |
| 8 | 0.93 | 4.54 | 3.27 | 6.74 | 5.52 |
| 9 | 3.52 | | | | |
| Sugar Beet, Plant Number: thousands per acre | | | | | |
| 1 | 19.6 | 22.3 | 21.5 | 19.6 | 21.9 |
| 2 | 22.4 | 22.8 | 23.0 | 17.5 | 22.0 |
| 4 | 23.1 | (b) 21.4* | 22.1 | 20.8 | 20.5 |
| 5 | 23.9 | 24.3 | 22.4 | 22.0 | 23.4 |
| 6 | 23.1 | 23.1 | 21.4 | 20.8 | 23.5 |
| 7 | 24.2 | 23.5 | 23.4 | 21.5 | 22.8 |
| 8 | 22.8 | 24.2 | 24.2 | 22.8 | 24.0 |
| 9 | 23.2 | | | | |
| Sugar Beet, Sugar Percentage | | | | | |
| 1 | 16.1 | 14.9 | 15.3 | 16.0 | 16.2 |
| 2 | 15.5 | 13.9 | 14.0 | 14.1 | 15.8 |
| 4 | 15.8 | (b) 14.3* | 14.8 | 15.7 | 16.2 |
| 5 | 16.0 | 14.2 | 15.5 | 15.6 | 16.4 |
| 6 | 15.4 | 14.6 | 15.3 | 15.7 | 16.1 |
| 7 | 15.9 | 15.6 | 16.0 | 15.6 | 16.1 |
| 8 | 15.3 | 14.9 | 15.9 | 14.9 | 15.5 |
| 9 | 15.4 | | | | |

*No nitrate of soda. Nitrogen applied as calcium and potassium nitrates.

HAY - THE PARK GRASS PLOTS 1955

The 100th year

For history, details of treatments etc. see "Results of the Field Experiments 1939-47" Vol.I, Section A/6.

Cultivations, etc.: Mineral fertilizers applied: Dec 27, 1954.
 Nitrogenous fertilizers applied: 1st dressing - Mar 16, 1955,
 2nd dressing - May 7. Cut: 1st - June 28, 2nd - Sept 16.

Summary of Results

Yield of Hay: cwt per acre

| Plot | Not limed | | | Limed | | |
|------|-----------|----------|-------|-------------------|------------------|-------------------|
| | 1st Crop | 2nd Crop | Total | 1st Crop | 2nd Crop | Total |
| 1 | 1.8 | 1.8 | 3.6 | 22.0 | 2.6 | 24.6 |
| 2 | 15.1 | 2.2 | 17.3 | 20.0 | 3.0 | 23.0 |
| 3 | 11.0 | 1.2 | 12.2 | 18.8 | 2.5 | 21.3 |
| 4-1 | 21.8 | 2.0 | 23.8 | 23.6 | 3.8 | 27.4 |
| 4-2 | 3.1 | 1.2 | 4.3 | 28.7 | 3.4 | 32.1 |
| 5-1 | 11.3 | 1.8 | 13.1 | | | |
| 5-2 | 26.6 | 5.6 | 32.2 | | | |
| 6 | 33.6 | 6.4 | 40.0 | | | |
| 7 | 33.1 | 3.9 | 37.0 | 52.6 | 7.5 | 60.1 |
| 8 | 28.1 | 2.1 | 30.2 | 28.7 | 3.1 | 31.8 |
| 9 | 18.0 | 8.5 | 26.5 | 54.1 | 6.4 | 60.5 |
| 10 | 10.2 | 8.2 | 18.4 | 39.1 | 3.1 | 42.2 |
| 11-1 | 9.0 | 19.5 | 28.5 | 54.1 | 7.8 | 61.9 |
| 11-2 | 20.3 | 18.2 | 38.5 | 62.4 | 13.4 | 75.8 |
| 12 | 16.7 | 2.5 | 19.2 | | | |
| 13 | 41.8 | 6.4 | 48.2 | 38.7 | 8.2 | 46.9 |
| 14 | 45.3 | 5.8 | 51.1 | 48.8 | 6.5 | 55.3 |
| 15 | 25.8 | 4.5 | 30.3 | 35.8 | 6.1 | 41.9 |
| 16 | 35.7 | 6.2 | 41.9 | 39.8 | 7.6 | 47.4 |
| 17 | 21.0 | 4.4 | 25.4 | 25.6 | 5.0 | 30.6 |
| 18 | 16.8 | 4.4 | 21.2 | 31.5 [*] | 3.9 [*] | 35.4 [*] |
| | | | | 31.7 ⁺ | 5.0 ⁺ | 36.7 ⁺ |
| 19 | 29.9 | 3.1 | 33.0 | 30.0 [*] | 2.9 [*] | 32.9 [*] |
| | | | | 34.6 ⁺ | 3.6 ⁺ | 38.2 ⁺ |
| 20 | 38.5 | 4.0 | 42.5 | 39.1 [*] | 3.1 [*] | 42.2 [*] |
| | | | | 35.7 ⁺ | 4.2 ⁺ | 39.9 ⁺ |

*Heavy liming

+Light liming

Note: The second crop was carted green; hay yields were estimated from the dry matter.

55/A/6

BARLEY - EXHAUSTION LAND HOOSFIELD 1955

For history, details of treatments etc., see "Results of the Field Experiments 1952", section A/6.

Cultivations, etc.: Ploughed: Sept 28, 1954. 1st dressing of ground chalk applied to acid areas: Dec 31. Ploughed: Jan 10, 1955. 2nd dressing of ground chalk applied: Mar 18. Nitrochalk applied at $3\frac{1}{2}$ cwt per acre: Mar 21. Seed drilled at 3 bushels per acre: Mar 22. Sprayed with M.C.P.A. at $2\frac{1}{2}$ pints in 40 gallons per acre: Mar 31. Parts of certain plots infested with wild oats (*Avena fatua*) cut and discarded: July 28. Harvested: Aug 15. Variety: Plumage Archer.

Note: Wild oats were pulled by hand during August from the harvested areas.

Summary of Results

| Manuring to Potatoes 1876-1901* | Yields: cwt per acre | |
|---|----------------------|--------------------|
| | Grain | Straw ⁺ |
| 1 Unmanured | 9.5 | 12.5 |
| 2 Unmanured after 6 years dung | 12.4 | 16.6 |
| 3 Dung | 27.1 | 28.5 |
| 4 Dung | 25.3 | 27.0 |
| 5 Ammonium salts | 11.8 | 14.1 |
| 6 Nitrate of soda | 8.0 | 11.4 |
| 7 Ammonium salts and complete minerals | 24.6 | 27.3 |
| 8 Nitrate of soda and complete minerals | 22.8 | 24.9 |
| 9 Superphosphate | 20.9 | 22.5 |
| 10 Complete minerals | 23.9 | 26.7 |

*For certain changes see history.

⁺At 85% dry matter

Mean dry matter % as threshed, Grain: 85.4
Straw: 85.6

55/A/7

WHEAT - WOBURN STACKYARD 1955

For history, details of treatments etc., see "Results of the Field Experiments" 1939-47, Vol. I, Section A/7.

In 1955 dressings of calcium carbonate were applied ranging from 20 to 50 cwt per acre according to the acidity of the plots, which were bare fallowed.

Cultivations, etc.: Ploughed: Nov 2, 1954. First dressing of ground chalk applied: Feb 15, 1955. Second dressing of ground chalk applied: Mar 8. Ploughed: Mar 17. Third dressing of ground chalk applied: Apr 16. Ploughed: May 6; Aug 2; Sept 26.

55/A/8

BARLEY - WOBURN STACKYARD 1955

For history, details of treatments etc., see "Results of the Field Experiments" 1939-47, Vol. I, Section A/7 and 1953 Section A/8. In 1955 dressings of calcium carbonate were applied ranging from 20 to 50 cwt per acre according to the acidity of the plots.

Cultivations, etc.: Ploughed: Nov 3, 1954. First dressing of chalk applied: Feb 15, 1955. Second dressing of chalk applied: Mar 8. Ploughed: Mar 18. Third dressing of chalk applied: Apr 6. Seed drilled at 3 bushels per acre: Apr 7. 'Nitrochalk' applied: Apr 12. Sprayed with MCPA amine at $2\frac{1}{2}$ pints per acre in low volume: May 22. Harvested: Aug 24. Variety: Plumage Archer.

Summary of Results

| Plot | 'Nitrochalk' dressing: cwt per acre | Grain: cwt per acre | Straw: cwt per acre |
|---------|-------------------------------------|---------------------|---------------------|
| 1 | 2 | 5.6 | 7.1 |
| 3 | 4 | 8.0 | 10.7 |
| 7 | 6 | 8.4 | 12.1 |
| 6 | 2 | 12.2 | 11.5 |
| 9 | 4 | 18.7 | 19.2 |
| 4 | 6 | 11.9 | 15.0 |
| 11b (3) | 2 | 11.7 | 12.7 |
| 11b (1) | 4 | 19.8 | 22.4 |
| 11b (2) | 6 | 22.4 | 25.9 |
| 10b | 2 | 6.1 | 7.1 |
| 11a | 4 | 9.8 | 11.9 |
| 10a | 6 | 4.4 | 7.8 |
| 5 | 2 | 6.9 | 8.1 |
| 8 | 4 | 9.6 | 11.3 |
| 2 | 6 | 5.6 | 7.9 |

55/Ba/1.1

THREE COURSE ROTATION EXPERIMENT

4th year of revised scheme

For details of treatments and rotation see "Results of the Field Experiments 1952", Section Ba/1.

Area of each plot: Potatoes (sub-plot), 0.0093 acre;
Barley, 0.0200 acre; Sugar beet, 0.0205 acre.

Cultivations, etc.:

Potatoes. Straw applied, all plots ploughed: Feb 3, 1955.
Fertilizers applied, potatoes planted: May 3. Earthed up:
July 5. Sprayed with sulphuric acid, 15% B.O.V: Sept 26.
Lifted: Oct 1. Variety: Majestic. Note: The crop yellowed
off and died down very early in the season.

Barley. Straw applied, all plots ploughed: Feb 3, 1955.
Ground chalk applied at 20 cwt per acre: Mar 19. Seed drilled
at 3 bushels per acre: Mar 22. Fertilizers applied: Mar 28.
Harvested: Aug 11. Variety: Plumage Archer.

Sugar beet. Straw applied, all plots ploughed: Feb 3, 1955.
Fertilizers applied: Apr 15. Seed drilled at 18 lb per acre:
Apr 16. Sprayed with D.D.T. emulsion, 3 pints in 40 gallons
per acre: June 13. Singled: June 18. Lifted: Nov 17.
Variety: Klein E.

Treatment symbols:-

Old Scheme

- Ar Complete artificials only
- St1 Straw ploughed in in autumn, artificials applied in spring
- St2 Straw ploughed in in autumn, artificials applied half in autumn,
half in spring
- Ad Adco ploughed in in autumn with supplementary artificials.

Revised Scheme

- St $5\frac{1}{3}$ cwt cut straw per acre in autumn
- Nitrogen dressing: 0.2; 0.4; 0.6 cwt N per acre as sulphate of
ammonia
- K_s Muriate of potash equivalent to K₂O in straw
- K 0.5 cwt K₂O per acre as muriate of potash.

Summary of Results

Potatoes

| Treatments applied: | 1953 and 1955 | 1955 | 0 | 0.4N | St + 0.2N | St + 0.6N | K _s | K + 0.4N |
|---------------------|---------------|-----------------------|------|------|-----------|-----------|----------------|----------|
| 1950 | 1952 & 1954 | 1955 | - | - | - | - | - | - |
| Ar | Ar | 0 | 3.67 | 3.78 | 4.00 | 4.56 | 4.23 | 5.26 |
| Ar | Ar | 0.4N | 3.66 | 4.32 | 4.08 | 4.47 | 4.00 | 4.38 |
| Ar | Ar | 0 | 3.43 | 4.40 | 4.00 | 4.47 | 4.00 | 4.38 |
| Ar | Ar | 0.4N | 3.72 | 4.05 | 4.08 | 4.47 | 4.00 | 4.38 |
| St1 St2 | St1 St2 | 0 | 4.35 | 4.12 | 4.08 | 4.47 | 4.00 | 4.38 |
| St1 St2 | St1 St2 | 0.4N | 4.06 | 4.15 | 4.08 | 4.47 | 4.00 | 4.38 |
| St1 St2 | St1 St2 | 0 | 4.09 | 4.19 | 4.08 | 4.47 | 4.00 | 4.38 |
| St1 St2 | St1 St2 | 0.4N | 4.19 | 4.15 | 4.08 | 4.47 | 4.00 | 4.38 |
| St1 St2 | St1 St2 | St+ 0.2N | 4.07 | 4.52 | 4.08 | 4.47 | 4.00 | 4.38 |
| St1 St2 | St1 St2 | St+ 0.6N | 4.38 | 4.06 | 4.08 | 4.47 | 4.00 | 4.38 |
| St1 St2 | St1 St2 | K _s | 3.54 | 4.06 | 4.08 | 4.47 | 4.00 | 4.38 |
| St1 St2 | St1 St2 | K _s +0.4N | 3.45 | 4.06 | 4.08 | 4.47 | 4.00 | 4.38 |
| St1 St2 | St1 St2 | 0 | 3.40 | 4.46 | 4.08 | 4.47 | 4.00 | 4.38 |
| Ad | Ad | 0.4N | 3.13 | 4.46 | 4.08 | 4.47 | 4.00 | 4.38 |
| Ad | Ad | St+ 0.6N | 4.46 | 4.46 | 4.08 | 4.47 | 4.00 | 4.38 |
| Ad | Ad | K _s + 0.4N | 3.94 | 4.46 | 4.08 | 4.47 | 4.00 | 4.38 |

Total tubers: tons per acre

| Treatments applied: | | Potatoes | | | | | | | |
|---------------------|---------|-----------------------|------|------|-----------|-----------|----------------|-----------------------|------|
| 1950 | 1951 | 1953 and 1955 | 0 | 0.4N | St + 0.2N | St + 0.6N | K _s | K _s + 0.4N | K |
| | | 1955 | - | - | - | - | - | - | - |
| | | 1952 & 1954 | - | - | - | - | - | - | - |
| | Ar | 0 | 64.7 | 69.4 | 69.8 | 73.0 | | | |
| | | 0.4N | | | | | | | |
| Ar | | 0 | 63.4 | 68.6 | 75.1 | 73.7 | | | |
| | | 0.4N | | | | | | | |
| St1 St2 | St1 St2 | 0 | 70.2 | 69.6 | 69.1 | 71.7 | 69.4 | 74.5 | 77.5 |
| | | 0.4N | | | | | | | |
| St1 St2 | | 0 | 76.2 | 67.7 | 68.1 | 71.9 | 68.6 | 69.8 | 72.8 |
| | | 0.4N | | | | | | | |
| | | St+ 0.2N | 78.2 | 71.7 | 65.1 | 75.6 | | | |
| | | St+ 0.6N | | | | | | | |
| | | K _s + 0.4N | 66.5 | 73.3 | 76.1 | 66.5 | | | |
| | Ad | 0 | | | 68.8 | 74.1 | 77.7 | 80.3 | 72.6 |
| Ad | | 0.4N | 73.2 | 71.9 | | | | | 71.1 |
| | | St+ 0.6N | 77.6 | 81.0 | | | | | |
| | | K _s + 0.4N | 73.2 | 75.8 | | | | | |

Percentage Ware (1 1/2" riddle)

55/Ba/1.4

| Treatments applied: | | | Barley | | | | | | |
|---|---------|-----------------------|--------|------|-----------|-----------|----------------|-----------------------|--|
| 1953 and 1955 | | | 0 | 0.4N | St + 0.2N | St + 0.6N | K _s | K _s + 0.4N | |
| 1950 | 1951 | 1952 & 1954 | | | | | | | |
| Grain (at 85% dry matter): cwt per acre | | | | | | | | | |
| | Ar | 0 | | 38.2 | | | | | |
| | | 0.4N | 32.8 | | | | | | |
| Ar | | 0 | | 40.0 | | | | | |
| | | 0.4N | 34.4 | | | | | | |
| | St1 St2 | 0 | | 40.3 | | 34.4 | | 38.2 | |
| | | 0.4N | 32.9 | | 29.5 | | 33.1 | | |
| St1 St2 | | 0 | | 40.6 | | | | | |
| | | 0.4N | 32.8 | | | | | | |
| | | St+ 0.2N | | 34.9 | | | | | |
| | | St+ 0.6N | 29.5 | | | | | | |
| | | K _s | | 40.3 | | | | | |
| | | K _s +0.4N | 35.7 | | | | | | |
| | Ad | 0 | | 39.4 | | 37.0 | | 38.1 | |
| Ad | | 0.4N | 39.9 | | | | | | |
| | | St+ 0.6N | 41.1 | | | | | | |
| | | K _s + 0.4N | 30.6 | | | | | | |
| Straw (at 85% dry matter): cwt per acre | | | | | | | | | |
| | Ar | 0 | | 39.8 | | | | | |
| | | 0.4N | 32.6 | | | | | | |
| Ar | | 0 | | 43.0 | | | | | |
| | | 0.4N | 34.8 | | | | | | |
| | St1 St2 | 0 | | 42.8 | | 37.6 | | 39.0 | |
| | | 0.4N | 29.9 | | 36.1 | | 31.0 | | |
| St1 St2 | | 0 | | 45.0 | | | | | |
| | | 0.4N | 29.7 | | | | | | |
| | | St+ 0.2N | | 34.5 | | | | | |
| | | St+ 0.6N | 29.2 | | | | | | |
| | | K _s | | 43.9 | | | | | |
| | | K _s +0.4N | 33.6 | | | | | | |
| | Ad | 0 | | 43.0 | | 37.9 | | 39.6 | |
| Ad | | 0.4N | 36.9 | | | | | | |
| | | St+ 0.6N | 38.9 | | | | | | |
| | | K _s + 0.4N | 27.8 | | | | | | |

Mean dry matter % as harvested Grain: 83.7
Straw: 80.8

55/Ba/1.5

| Treatments applied: 1953 and 1955 | | | Sugar Beet | | | | | |
|-----------------------------------|---------|-----------------------|-------------------------------|------|-----------|-----------|----------------|-----------------------|
| 1950 | 1951 | 1952 & 1954 | 0 | 0.4N | St + 0.2N | St + 0.6N | K _s | K _s + 0.4N |
| | | | Roots (washed): tons per acre | | | | | |
| | Ar | 0 | | 9.56 | | | | |
| | | 0.4N | 6.74 | | | | | |
| Ar | | 0 | | 8.18 | | | | |
| | | 0.4N | 6.72 | | | | | |
| | St1 St2 | 0 | | 7.40 | | 8.45 | | 7.55 |
| | | 0.4N | 7.05 | | 7.07 | | 8.47 | |
| St1 St2 | | 0 | | 7.70 | | | | |
| | | 0.4N | 7.07 | | | | | |
| | | St+ 0.2N | | 7.31 | | | | |
| | | St+ 0.6N | 7.99 | | | | | |
| | | K _s | | 8.01 | | | | |
| | | K _s + 0.4N | 8.18 | | | | | |
| | Ad | 0 | | 9.54 | | 7.44 | | 7.77 |
| Ad | | 0.4N | 6.29 | | | | | |
| | | St+ 0.6N | 5.98 | | | | | |
| | | K _s + 0.4N | 9.49 | | | | | |
| | | | Sugar Percentage | | | | | |
| | Ar | 0 | | 17.1 | | | | |
| | | 0.4N | 17.1 | | | | | |
| Ar | | 0 | | 16.9 | | | | |
| | | 0.4N | 16.7 | | | | | |
| | St1 St2 | 0 | | 16.8 | | 16.5 | | 17.0 |
| | | 0.4N | 16.8 | | 17.3 | | 16.9 | |
| St1 St2 | | 0 | | 16.8 | | | | |
| | | 0.4N | 17.0 | | | | | |
| | | St+ 0.2N | | 17.0 | | | | |
| | | St+ 0.6N | 17.5 | | | | | |
| | | K _s | | 16.8 | | | | |
| | | K _s + 0.4N | 17.6 | | | | | |
| | Ad | 0 | | 17.2 | | 16.6 | | 16.1 |
| Ad | | 0.4N | 16.8 | | | | | |
| | | St+ 0.6N | 16.6 | | | | | |
| | | K _s + 0.4N | 17.6 | | | | | |

55/Ba/1.6

Sugar Beet

| Treatments applied: 1953 and 1955 | | | 0 | 0.4N | St + 0.2N | St + 0.6N | K _s | K _s + 0.4N |
|-----------------------------------|---------|-----------------------|---------------------------|------|-----------|-----------|----------------|-----------------------|
| 1950 | 1951 | 1952 & 1954 | Total sugar: cwt per acre | | | | | |
| | Ar | 0 | | 32.6 | | | | |
| | | 0.4N | 23.1 | | | | | |
| Ar | | 0 | | 27.7 | | | | |
| | | 0.4N | 22.4 | | | | | |
| | St1 St2 | 0 | | 24.8 | | 27.9 | | 25.8 |
| | | 0.4N | 23.7 | | 24.5 | | 28.7 | |
| St1 St2 | | 0 | | 25.9 | | | | |
| | | 0.4N | 24.1 | | | | | |
| | | St+ 0.2N | | 24.8 | | | | |
| | | St+ 0.6N | 28.0 | | | | | |
| | | K | | 27.0 | | | | |
| | | K _s + 0.4N | 28.8 | | | | | |
| | Ad | 0 | | 32.9 | | 24.8 | | 25.1 |
| Ad | | 0.4N | 21.1 | | | | | |
| | | St+ 0.6N | 19.9 | | | | | |
| | | K _s + 0.4N | 33.5 | | | | | |
| | | | Tops: tons per acre | | | | | |
| | Ar | 0 | | 6.87 | | | | |
| | | 0.4N | 3.91 | | | | | |
| Ar | | 0 | | 6.18 | | | | |
| | | 0.4N | 4.12 | | | | | |
| | St1 St2 | 0 | | 5.09 | | 6.83 | | 5.85 |
| | | 0.4N | 4.34 | | 4.69 | | 5.28 | |
| St1 St2 | | 0 | | 5.65 | | | | |
| | | 0.4N | 4.54 | | | | | |
| | | St+ 0.2N | | 5.28 | | | | |
| | | St+ 0.6N | 4.82 | | | | | |
| | | K | | 5.70 | | | | |
| | | K _s + 0.4N | 4.74 | | | | | |
| | Ad | 0 | | 6.59 | | 5.78 | | 6.59 |
| Ad | | 0.4N | 3.78 | | | | | |
| | | St+ 0.6N | 3.95 | | | | | |
| | | K _s + 0.4N | 6.37 | | | | | |

| Treatments applied: | | | Sugar Beet | | | | | |
|---------------------|---------|-----------------------|----------------------------------|------|-----------|-----------|----------------|-----------------------|
| 1953 and 1955 | | | 0 | 0.4N | St + 0.2N | St + 0.6N | K _s | K _s + 0.4N |
| 1950 | 1951 | 1952 & 1954 | | | | | | |
| | | | Plant number: thousands per acre | | | | | |
| | Ar | 0 | | 27.7 | | | | |
| | | 0.4N | 31.1 | | | | | |
| Ar | | 0 | | 27.8 | | | | |
| | | 0.4N | 29.6 | | | | | |
| | St1 St2 | 0 | | 26.7 | | 28.2 | | 26.9 |
| | | 0.4N | 27.7 | | 29.5 | | 28.0 | |
| St1 St2 | | 0 | | 27.8 | | | | |
| | | 0.4N | 29.7 | | | | | |
| | | St+ 0.2N | | 25.8 | | | | |
| | | St+ 0.6N | 30.6 | | | | | |
| | | K _s | | 27.6 | | | | |
| | | K _s + 0.4N | 28.7 | | | | | |
| | Ad | 0 | | 27.8 | | 26.7 | | 29.5 |
| Ad | | 0.4N | 28.3 | | | | | |
| | | St+ 0.6N | 25.8 | | | | | |
| | | K _s + 0.4N | 26.9 | | | | | |

55/Ba/2.1

FOUR COURSE ROTATION EXPERIMENT

1st year of revised scheme

The experiment in its original form ended at the harvest of 1954. A summary of the results and details of the original scheme appear in the Station annual report for 1954 (p.153).

The cropping rotation has been modified by introducing beans (autumn sown when possible) instead of ryegrass ley, the present rotation being: potatoes, barley, beans, wheat.

The applications of dung, straw, straw compost and rock phosphate have been discontinued. The plots of the original dung, straw and superphosphate series now receive an annual dressing of 0.24 cwt P_2O_5 per acre applied as superphosphate, while the old compost plots receive 0.12 cwt P_2O_5 annually as superphosphate. The rock phosphate plots receive no phosphate. All plots receive a basal dressing of 0.6 cwt K_2O annually as muriate of potash (but see below for the beans of 1955 and the wheat of 1956).

Each plot of wheat, barley and potatoes is split into 2 for the application of nitrogen:-

| | |
|-------------------|---|
| wheat and barley: | none; 0.4 cwt N per acre applied as sulphate of ammonia. |
| potatoes: | 0.2; 0.6 cwt N per acre applied as sulphate of ammonia |

The arrangement of the levels of nitrogen is randomized afresh each season. The beans do not receive nitrogen.

The phosphate and potash fertilizers are applied in autumn for beans and wheat, half-plots of wheat receiving a single top dressing of nitrogen in spring. All fertilizers for barley are applied to the seedbed. All fertilizers for potatoes are broadcast on the flat before planting, which is by machine.

In 1955 the plots of beans were split into 3 for the application of potash:-

none; 0.8; 1.6 cwt K_2O per acre applied as muriate of potash.

The wheat following these beans will receive equalizing amounts of potash:- 1.6 cwt K_2O following none; 0.8 following 0.8 and none following 1.6. It is not intended to repeat this test.

55/Ba/2.2

Area of each sub plot: Potatoes, wheat: 0.0129 acre. Barley: 0.0120
acre. Beans: 0.0081 acre.

Area harvested: Potatoes: 0.0106 acre. Wheat: 0.0057 acre.
Barley: 0.0052 acre. Beans: 0.0038 acre.

Cultivations, etc.:

Potatoes.

Ploughed: Sept 22, 1954. Fertilizers applied broadcast on flat:
May 3, 1955. Machine planted: May 4. Earthed up: July 6.
Sprayed with sulphuric acid, 20% B.O.V.: Oct 4. Lifted: Oct 10.
Variety: Majestic.

Barley.

Ploughed: Oct 12, 1954. Seed drilled at 3 bushels per acre:
Mar 22, 1955. Fertilizers applied: Mar 31. Sprayed with M.C.P.A.,
medium volume, $2\frac{1}{2}$ pints per acre: June 8. Combine harvested:
Aug 19. Variety: Plumage Archer.

Beans.

Ploughed: Sept 8, 1954. Potash dressings applied: Oct 26.
Phosphate dressings applied: Mar 18, 1955. Seed drilled at 200 lb.
per acre: Mar 21. Harvested: Aug 22. Variety: Gartons Spring Tick.

Wheat.

Ploughed: July 23 and again Aug 26, 1954. Seed sown at $2\frac{3}{4}$ bushels
per acre: Oct 22. Basal potash fertilizer applied: Oct 26.
Nitrogen and phosphate fertilizer treatments applied: May 11, 1955.
Sprayed with M.C.P.A., medium volume, $2\frac{1}{2}$ pints per acre: May 31.
Combine harvested: Aug 24. Variety: Yeoman.

55/Ba/2.3

Summary of Results

| Previous Treatment | Year applied | P2O5: cwt per acre applied 1955 | Potatoes | | | Percentage Ware | | | Barley (at 85% Dry matter): cwt per acre | | | |
|----------------------|--------------|---------------------------------|-----------------------------|-------------|-------|-----------------|------|-------|--|------|-------|-------|
| | | | Total tubers: tons per acre | N: per acre | Diff. | N: per acre | Mean | Diff. | N: per acre | Mean | Diff. | |
| Manure | 1954 | | 4.28 | 5.03 | 4.66 | +0.75 | 82.1 | 82.1 | 27.2 | 34.0 | 30.6 | +6.8 |
| | 1953 | | 4.28 | 4.91 | 4.60 | +0.63 | 81.6 | 80.8 | 19.4 | 33.8 | 26.6 | +14.4 |
| | 1952 | 0.24 | 5.08 | 6.71 | 5.90 | +1.63 | 81.6 | 85.6 | 18.5 | 28.9 | 23.7 | +10.4 |
| | 1951 | | 3.69 | 4.70 | 4.20 | +1.01 | 81.8 | 82.0 | 25.7 | 26.9 | 26.3 | +1.2 |
| | 1950 | | 5.24 | 5.33 | 5.28 | +0.09 | 82.0 | 82.9 | 18.4 | 25.0 | 21.7 | +6.6 |
| Adco (straw compost) | 1954 | | 4.41 | 4.83 | 4.62 | +0.42 | 84.1 | 83.8 | 28.9 | 25.9 | 27.4 | -3.0 |
| | 1953 | | 4.66 | 4.70 | 4.68 | +0.04 | 87.1 | 84.8 | 17.9 | 21.1 | 19.5 | +3.2 |
| | 1952 | 0.12 | 3.52 | 4.74 | 4.13 | +1.22 | 84.3 | 85.3 | 11.4 | 20.6 | 16.0 | +9.2 |
| | 1951 | | 4.66 | 4.62 | 4.64 | -0.04 | 81.9 | 82.6 | 10.0 | 16.8 | 13.4 | +6.8 |
| | 1950 | | 3.99 | 4.03 | 4.01 | +0.04 | 84.6 | 86.0 | 22.6 | 23.1 | 22.8 | +0.5 |
| Straw | 1954 | | 4.78 | 5.20 | 4.99 | +0.42 | 82.0 | 83.9 | 22.4 | 34.9 | 28.6 | +12.5 |
| | 1953 | | 4.78 | 5.33 | 5.06 | +0.55 | 83.0 | 84.6 | 23.5 | 23.3 | 23.4 | -0.2 |
| | 1952 | 0.24 | 5.12 | 5.75 | 5.44 | +0.63 | 85.9 | 85.4 | 23.0 | 26.7 | 24.8 | +3.7 |
| | 1951 | | 3.73 | 5.12 | 4.42 | +1.39 | 78.9 | 79.6 | 18.5 | 33.7 | 26.1 | +15.2 |
| | 1950 | | 4.95 | 5.24 | 5.10 | +0.29 | 87.7 | 86.0 | 23.1 | 35.4 | 29.2 | +12.3 |
| Super-phosphate | 1954 | | 4.28 | 4.91 | 4.60 | +0.63 | 83.3 | 84.6 | 26.5 | 24.7 | 25.6 | -1.8 |
| | 1953 | | 4.99 | 6.04 | 5.52 | +1.05 | 80.9 | 87.7 | 17.3 | 27.6 | 22.4 | +10.3 |
| | 1952 | 0.24 | 4.70 | 4.78 | 4.74 | +0.08 | 81.0 | 81.6 | 23.6 | 35.2 | 29.4 | +11.6 |
| | 1951 | | 3.78 | 4.41 | 4.10 | +0.63 | 79.5 | 81.7 | 15.0 | 23.6 | 19.3 | +8.6 |
| | 1950 | | 4.49 | 5.20 | 4.84 | +0.71 | 85.0 | 84.4 | 13.1 | 23.0 | 18.0 | +9.9 |
| Rock phosphate | 1954 | | 4.03 | 4.74 | 4.38 | +0.71 | 80.5 | 87.6 | 11.6 | 16.0 | 13.8 | +4.4 |
| | 1953 | | 3.52 | 4.57 | 4.04 | +1.05 | 85.4 | 86.7 | 18.9 | 21.3 | 20.1 | +2.4 |
| | 1952 | None | 3.57 | 4.15 | 3.86 | +0.58 | 76.5 | 84.8 | 19.9 | 23.5 | 21.7 | +3.6 |
| | 1951 | | 3.52 | 4.45 | 3.98 | +0.93 | 83.4 | 85.3 | 10.9 | 13.8 | 12.4 | +2.9 |
| | 1950 | | 4.49 | 4.45 | 4.47 | -0.04 | 86.7 | 86.0 | 13.6 | 19.6 | 16.6 | +6.0 |

Mean dry matter % as harvested:

55/Ba/2.4

| Previous Treatment | Year applied | P2O5: cwt per acre applied 1955 | Beans (at 85% dry matter): | | | Wheat | | | | |
|---------------------------------|--------------|---------------------------------|--|--------------|------|-------------|---------------------|------|-------|-------|
| | | | Grain (at 85% dry matter): K ₂ O: cwt per acre | cwt per acre | Mean | N: per acre | Grain: cwt per acre | Mean | Diff. | |
| Manure | | | None | 0.8 | 1.6 | None | 0.4 | | | |
| Dung | 1954 | | 11.4 | 11.7 | 17.1 | 13.4 | 13.8 | 24.7 | 19.2 | +10.9 |
| | 1953 | | 15.5 | 15.2 | 13.8 | 14.8 | 9.4 | 17.2 | 13.3 | +7.8 |
| | 1952 | 0.24 | 14.3 | 16.2 | 20.2 | 16.9 | 12.8 | 23.2 | 18.0 | +10.4 |
| | 1951 | | 14.8 | 12.1 | 12.1 | 13.0 | 9.7 | 20.4 | 15.0 | +10.7 |
| | 1950 | | 17.9 | 19.8 | 14.5 | 17.4 | 10.7 | 18.5 | 14.6 | +7.8 |
| Adco (straw compost) | 1954 | | 18.1 | 14.8 | 15.0 | 16.0 | 12.2 | 21.6 | 16.9 | +9.4 |
| | 1953 | | 16.7 | 13.1 | 12.6 | 14.1 | 11.7 | 19.1 | 15.4 | +7.4 |
| | 1952 | 0.12 | 11.4 | 15.2 | 11.2 | 12.6 | 11.3 | 19.6 | 15.4 | +8.3 |
| | 1951 | | 18.1 | 16.7 | 14.3 | 16.4 | 9.1 | 16.0 | 12.6 | +6.9 |
| | 1950 | | 12.4 | 15.7 | 9.3 | 12.5 | 11.7 | 20.1 | 15.9 | +8.4 |
| Straw | 1954 | | 16.4 | 18.8 | 15.0 | 16.7 | 13.3 | 19.6 | 16.4 | +6.3 |
| | 1953 | | 16.4 | 13.1 | 15.7 | 15.1 | 11.0 | 20.1 | 15.6 | +9.1 |
| | 1952 | 0.24 | 17.1 | 16.0 | 16.7 | 16.6 | 14.1 | 16.0 | 15.0 | +1.9 |
| | 1951 | | 13.8 | 12.1 | 10.5 | 12.1 | 11.0 | 20.4 | 15.7 | +9.4 |
| | 1950 | | 12.6 | 13.6 | 15.2 | 13.8 | 11.3 | 23.5 | 17.4 | +12.2 |
| Super-phosphate | 1954 | | 15.0 | 11.4 | 14.8 | 13.7 | 8.6 | 16.4 | 12.5 | +7.8 |
| | 1953 | | 15.0 | 6.9 | 13.3 | 11.7 | 9.4 | 18.5 | 14.0 | +9.1 |
| | 1952 | 0.24 | 19.8 | 17.4 | 15.5 | 17.6 | 11.0 | 16.4 | 13.7 | +5.4 |
| | 1951 | | 10.7 | 13.3 | 12.9 | 12.3 | 9.1 | 16.9 | 13.0 | +7.8 |
| | 1950 | | 11.2 | 17.4 | 9.8 | 12.8 | 8.6 | 16.9 | 12.8 | +8.3 |
| Rock phosphate | 1954 | | 12.6 | 15.2 | 13.1 | 13.6 | 8.6 | 18.5 | 13.6 | +9.9 |
| | 1953 | | 12.6 | 12.4 | 11.4 | 12.1 | 11.7 | 17.2 | 14.4 | +5.5 |
| | 1952 | None | 12.4 | 11.2 | 10.0 | 11.2 | 8.1 | 16.0 | 12.0 | +7.9 |
| | 1951 | | 12.1 | 11.2 | 13.3 | 12.2 | 10.7 | 17.2 | 14.0 | +6.5 |
| | 1950 | | 10.2 | 9.8 | 11.4 | 10.5 | 7.5 | 13.3 | 10.4 | +5.8 |
| Mean dry matter % as harvested: | | | | | | 87.0 | | | 84.9 | |

55/Ba/3.1

SIX COURSE ROTATION EXPERIMENT

The 26th year

Seasonal effects of fertilizers - Rothamsted Long Hoos IV and Woburn Stackyard 1955.

For details of treatments, rotation etc., see "Results of the Field Experiments 1939-47" Vol.I, Section Ba/4.

Area of each plot: Rothamsted, 0.0250 acre. Woburn, 0.0266 acre.

Cultivations, etc.:

Rothamsted

Sugar beet.

Ploughed: Sept 6, 1954 and Nov 5. Fertilizers applied: Apr 15, 1955. Seed drilled at 18 lb per acre: Apr 16. Sprayed with D.D.T. emulsion, 3 pints in 40 gallons: June 13. Singled: June 20. Lifted: Nov 16. Variety: Klein E.

Barley.

Ploughed: Dec 27, 1954. Ground chalk applied at 20 cwt per acre: Mar 19, 1955. Seed drilled at 3 bushels per acre: Mar 22. Fertilizers applied: Mar 29. Harvested: Aug 8. Variety: Plumage Archer. Mean D.M.%: Grain, 81.8; Straw, 74.9.

Clover.

Seed undersown in barley at 40 lb per acre: Apr 20, 1954. Autumn fertilizers applied: Dec 1. Sulphate of ammonia applied: Apr 25, 1955. Cut: July 12. Variety: Late flowering Montgomery Red.

Wheat.

Ploughed: Aug 5, 1954. Seed drilled at $2\frac{3}{4}$ bushels per acre: Oct 22. Autumn fertilizers applied: Oct 25. Sulphate of ammonia applied: May 12, 1955. Harvested: Aug 8. Variety: Yeoman. Mean D.M.%: Grain, 81.6; Straw, 82.0.

Potatoes.

Ploughed: Sept 6, 1954 and Nov 4. Ridged, fertilizers applied: Apr 26, 1955. Potatoes planted: Apr 27. Earthed up: July 4. Sprayed with sulphuric acid, 15% B.O.V: Sept 26. Lifted: Oct 1. Variety: Majestic.

Rye.

Cultivated: Oct 4, 1954. Ground chalk applied at 20 cwt per acre: Oct 9. Seed drilled at 3 bushels per acre: Oct 22. Autumn fertilizers applied: Oct 25. Sulphate of ammonia applied: May 12, 1955. Harvested: Aug 15. Variety: King II. Mean D.M.%: Grain, 85.9; Straw, 87.5.

55/Ba/3.2

Woburn

Sugar beet.

Ploughed: Sept 29, 1954 and Feb 2, 1955. Fertilizers applied: Apr 18.
Seed drilled at 12 lb per acre: Apr 19. Sprayed with parathion, $\frac{1}{2}$
pint per acre in 40 gallons: June 10. Singled: June 28. Lifted:
Oct 27. Variety: Klein E.

Barley.

Ploughed: Nov 5, 1954 and Feb 1, 1955. Ground chalk applied at
20 cwt per acre: Mar 14. Fertilizers applied, seed drilled at $3\frac{1}{3}$
bushels per acre: Mar 17. Harvested: Aug 5. Variety:
Plumage Archer.

Clover.

Seed undersown in barley at 40 lb per acre: Apr 26, 1954. Autumn
fertilizers applied: Dec 2. Sulphate of ammonia applied:
Apr 29, 1955. Cut: July 6. Variety: Late flowering Montgomery
Red. N.B. The crop was severely and irregularly damaged by
Sclerotinia.

Wheat.

Ploughed: July 5, 1954 and Oct 1. Autumn fertilizers applied, seed
drilled at 3 bushels per acre: Oct 28. Sulphate of ammonia applied:
May 6, 1955. Sprayed with M.C.P.A. low volume, at $2\frac{1}{2}$ pints per acre:
May 20. Harvested: Aug 15. Variety: Squareheads Master 13/4.

Potatoes.

Ploughed: Sept 29, 1954 and Feb 1, 1955. Ridged, fertilizers
applied: Apr 21. Potatoes planted: Apr 22. Earthed up: June 28.
Sprayed with copper fungicide, 5 lb in 40 gallons per acre:
Aug 19. Sprayed with arsenious compound, 1 gallon in 40 gallons
per acre: Sept 27. Lifted: Oct 3. Variety: Majestic.

Rye.

Ploughed: Oct 23, 1954. Ground chalk applied at 20 cwt per acre,
autumn fertilizer applied: Oct 28. Seed drilled at 3 bushels per
acre: Oct 29. Sulphate of ammonia applied: May 6, 1955.
Harvested: Aug 15. Variety: King II.

Erratum to "Results of the Field Experiments" 1954!

Page 54/Ba/3.3:- After "clover, hay" read "(at 85% Dry Matter)" not
"dry matter".

Erratum to "Results of the Field Experiments" 1939-47, Vol. I.

Page Ba/4.4:- Woburn. Clover, Hay Dry Matter: cwt per acre.
Response to N in 1945 should read "-12.5" NOT "12.5".

55/Ba/3.3

Summary of Results

Mean yields per acre and responses in yield per cwt of N, P₂O₅ and K₂O

| | Rothamsted | Woburn | Rothamsted | Woburn |
|----------------|---|---------------------|---|--------|
| | Sugar Beet, roots (washed): tons per acre | | Barley, grain: cwt per acre | |
| Mean | 7.58 | 8.46 | 34.3 | 27.2 |
| Response to: N | +5.95 | +4.07 | +23.3 | +20.1 |
| P | +0.11 | -1.74 | +1.4 | +2.2 |
| K | -0.12 | -0.58 | -5.3 | +3.7 |
| | Sugar Beet, sugar percentage | | Barley, straw: cwt per acre | |
| Mean | 16.7 | 18.4 | 39.0 | 34.7 |
| Response to: N | +0.7 | -0.1 | +34.5 | +39.9 |
| P | +0.3 | +0.1 | +2.3 | +3.3 |
| K | +0.3 | +0.9 | -7.3 | +2.2 |
| | Sugar Beet, total sugar: cwt per acre | | Clover, hay: dry matter cwt per acre | |
| Mean | 25.4 | 31.1 | 54.0 | 46.0 |
| Response to: N | +21.1 | +14.6 | +13.3 | -9.5 |
| P | +0.8 | -6.2 | +7.9 | +2.5 |
| K | +0.1 | -0.6 | -2.0 | +12.8 |
| | Sugar Beet, tops: tons per acre | | Wheat, grain: cwt per acre | |
| Mean | 4.63 | 5.55 | 38.0 | 8.8 |
| Response to: N | +4.05 | +3.19 | +8.1 | +10.0 |
| P | -0.16 | -1.27 | -5.3 | +2.1 |
| K | -0.25 | +0.36 | +0.4 | +1.5 |
| | Sugar Beet, plant number: thousands per acre | | Wheat, straw: cwt per acre | |
| Mean | 28.3 | + | 56.1 | 12.6 |
| Response to: N | +0.9 | | +14.3 | +20.1 |
| P | -0.1 | | -7.0 | +2.5 |
| K | +0.9 | | +3.5 | +0.8 |
| | Potatoes, total tubers: tons per acre | | Rye, grain: cwt per acre | |
| Mean | 4.99 | 5.33 | 32.7 | 25.4 |
| Response to: N | +1.24 | +0.83 | +28.2 | +27.7 |
| P | +0.93 | +0.67 | -6.5 | -2.4 |
| K | +0.50 | -0.23 | -1.1 | -0.5 |
| | Potatoes, percentage ware | | Rye, straw: cwt per acre | |
| Mean | 83.1 ⁽¹⁾ | 85.4 ⁽²⁾ | 37.2 | 30.9 |
| Response to: N | -2.2 | -1.4 | +28.0 | +33.3 |
| P | -10.9 | -4.8 | -2.3 | +0.5 |
| K | +6.5 | +3.1 | -3.3 | -1.8 |

* (at 85% dry matter) + not recorded.

Riddle: (1) 1½"; (2) 1⅝"

DEEP CULTIVATION ROTATION EXPERIMENT

The 12th year

Deep ploughing, fertilizers and dung - Long Hoos I and II 1955.

For details of rotation and treatments etc. see "Results of the Field Experiments 1939-47" Vol.I, Section Bc/1.

Area of each plot: 0.0312 acre. Area harvested: Sugar beet
($\frac{1}{2}$ plot) 0.0119 acre; barley, spring oats, 0.0265 acre;
ley, 0.0275 acre; wheat, 0.0188 acre; potatoes ($\frac{1}{2}$ plot), 0.0107 acre.

Cultivations, etc.:

Sugar beet

Fertilizers for ploughing in "deep" applied: Sept 27, 1954.
Dung to "deep" plots applied, ploughed all "deep" plots: Oct 22.
Dung and fertilizers for ploughing in "shallow" applied: Oct 22.
"Shallow" plots ploughed: Oct 29. Fertilizers for surface application applied: Apr 16, 1955. Seed drilled at 18 lb per acre: Apr 16. Sprayed with D.D.T. emulsion at 3 pints in 40 gallons per acre: June 13. Singled: June 21. Lifted: Nov 22. Variety: Klein E.

Barley

Ploughed: Jan 1, 1955. Ground chalk at 20 cwt per acre applied: Mar 18. Basic slag and sulphate of ammonia applied: Mar 21. Seed drilled at 3 bushels per acre: Mar 22. Harvested: Aug 8. Variety: Plumage Archer.

Ley

Seeds undersown in barley: Apr 21, 1954. Cut: July 11, 1955. Seeds mixture (per acre): 18 lb S.24 perennial ryegrass, 8 lb Montgomery red clover, 2 lb American Alsike clover.

Wheat

"Deep" plots ploughed: Oct 21, 1954. "Shallow" plots ploughed: Oct 27. Seed drilled at $2\frac{3}{4}$ bushels per acre: Dec 22. Sulphate of ammonia applied: May 13, 1955. Sprayed with M.C.P.A., medium volume, $2\frac{1}{2}$ pints per acre: May 31. Combine harvested: Aug 24. Variety: Yeoman.

Potatoes

Fertilizers for ploughing in "deep" applied: Sept 27, 1954. Dung to "deep" plots applied, ploughed all "deep" plots: Oct 22. Dung and fertilizers for ploughing in "shallow" applied: Oct 22. "Shallow" plots ploughed: Oct 29. Ridged: Apr 22, 1955. Fertilizers for surface application applied: Apr 27. Potatoes planted: Apr 28. Earthed up: July 4. Sprayed with sodium arsenite, 1 gallon in 80 gallons: Oct 10. Lifted: Oct 18. Variety: Majestic.

55/Bb/1.2

Spring oats

Ploughed: Nov 4, 1954. Ground chalk at 20 cwt per acre applied:
Mar 18, 1955. Sulphate of ammonia applied, seed drilled at
4 bushels per acre: Mar 21. Sprayed with M.C.P.A., high volume,
2 pints in 80 gallons: June 2. Harvested: Aug 4. Variety: Star.

Standard errors per plot:

| | | | |
|-----------------------------------|-------------|------------------------------|----------|
| Sugar beet, Total sugar. | Whole plot: | 1.99 cwt per acre or 6.1% | (4 d.f.) |
| | Sub plot: | 2.31 cwt per acre or 7.1% | (7 d.f.) |
| Tops. | Whole plot: | 0.387 tons per acre or 4.9% | (4 d.f.) |
| | Sub plot: | 0.553 tons per acre or 7.0% | (7 d.f.) |
| Barley, Grain: | | 2.27 cwt per acre or 7.1% | (4 d.f.) |
| Ley, Hay: | | 1.88 cwt per acre or 2.8% | (4 d.f.) |
| Wheat, Grain: | | 1.51 cwt per acre or 4.0% | (4 d.f.) |
| Potatoes, Total tubers. | Whole plot: | 0.927 tons per acre or 13.3% | (4 d.f.) |
| | Sub plot: | 0.389 tons per acre or 5.6% | (7 d.f.) |
| Spring Oats, Grain (at 85% D.M.): | | 2.21 cwt per acre or 5.8% | (4 d.f.) |

Erratum to "Results of the Field Experiments" 1949, page 49/Bb/1.5.

Roots (washed): tons per acre, "Mean of Potash-None" should read
"8.93 not 8.03".

Summary of Results

Sugar Beet

Responses to treatments

| Response to | Mean | Ploughing | | Dung | | Phosphate | | Potash | |
|-------------|------|-----------|------|------|-------|-----------|-------|--------|-------|
| | | Shallow | Deep | Abs. | Pres. | Abs. | Pres. | Abs. | Pres. |

Roots (washed): Mean yield 10.05 tons per acre

| | | | | | | | | | |
|------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Ploughing deep-shallow | +0.89 | - | - | +1.07 | +0.71 | +0.31 | +1.47 | +0.77 | +1.01 |
| Dung | +5.18 | +5.36 | +5.00 | - | - | +5.90 | +4.46 | +5.51 | +4.85 |
| Phosphate | +0.67 | +0.09 | +1.25 | +1.39 | -0.05 | - | - | +0.71 | +0.63 |
| Potash | +0.53 | +0.41 | +0.65 | +0.86 | +0.20 | +0.57 | +0.49 | - | - |

Sugar Percentage: Mean 16.1

| | | | | | | | | | |
|------------------------|------|------|------|------|------|------|------|------|------|
| Ploughing deep-shallow | -0.1 | - | - | -0.3 | +0.1 | -0.3 | +0.1 | -0.1 | -0.1 |
| Dung | +0.6 | +0.4 | +0.8 | - | - | +0.7 | +0.5 | +0.9 | +0.3 |
| Phosphate | +0.3 | +0.1 | +0.5 | +0.4 | +0.2 | - | - | +0.3 | +0.3 |
| Potash | 0.0 | 0.0 | 0.0 | +0.3 | -0.3 | 0.0 | 0.0 | - | - |

Total Sugar: Mean yield 32.6 cwt per acre

(± 1.00)

(± 1.41)

| | | | | | | | | | |
|------------------------|-------|-------|-------|------|------|-------|-------|-------|-------|
| Ploughing deep-shallow | +2.9 | - | - | +3.1 | +2.7 | +0.4 | +5.4 | +2.3 | +3.5 |
| Dung | +17.9 | +18.1 | +17.7 | - | - | +20.2 | +15.6 | +19.6 | +16.2 |
| Phosphate | +2.7 | +0.2 | +5.2 | +5.0 | +0.4 | - | - | +3.0 | +2.4 |
| Potash | +1.6 | +1.0 | +2.2 | +3.3 | -0.1 | +1.9 | +1.3 | - | - |

Tops: Mean yield 7.86 tons per acre

(± 0.193)

(± 0.274)

| | | | | | | | | | |
|------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Ploughing deep-shallow | +0.41 | - | - | +0.26 | +0.56 | +0.25 | +0.57 | +0.55 | +0.27 |
| Dung | +2.92 | +2.77 | +3.07 | - | - | +2.98 | +2.86 | +2.81 | +3.03 |
| Phosphate | +0.66 | +0.50 | +0.82 | +0.72 | +0.60 | - | - | +0.91 | +0.41 |
| Potash | +0.43 | +0.57 | +0.29 | +0.32 | +0.54 | +0.68 | +0.18 | - | - |

Plant Number: Mean 28.7 thousands per acre

| | | | | | | | | | |
|------------------------|------|------|------|------|------|------|------|------|------|
| Ploughing deep-shallow | +0.4 | - | - | +0.9 | -0.1 | +0.1 | +0.7 | +0.5 | +0.3 |
| Dung | +0.4 | +0.9 | -0.1 | - | - | +0.5 | +0.3 | 0.0 | +0.8 |
| Phosphate | -0.2 | -0.5 | +0.1 | -0.1 | -0.3 | - | - | -0.3 | -0.1 |
| Potash | +0.1 | +0.2 | 0.0 | -0.3 | +0.5 | 0.0 | +0.2 | - | - |

55/Bb/1.4

Sugar Beet

| | Phosphate | | | Potash | | | Mean |
|-------------------------------|-----------|-------------|-------------|--------|-------------|-------------|-------|
| | None | Ploughed in | In seed bed | None | Ploughed in | In seed bed | |
| Roots (washed): tons per acre | | | | | | | |
| Shallow | 9.56 | 9.96 | 9.34 | 9.40 | 9.77 | 9.86 | 9.61 |
| Deep | 9.87 | 11.49 | 10.75 | 10.17 | 10.91 | 10.74 | 10.49 |
| No dung | 6.77 | 8.78 | 7.53 | 7.03 | 8.05 | 7.73 | 7.46 |
| Dung | 12.67 | 12.67 | 12.56 | 12.54 | 12.62 | 12.86 | 12.64 |
| Mean | 9.72 | 10.72 | 10.04 | 9.78 | 10.34 | 10.30 | 10.05 |

Sugar Percentage

| | | | | | | | |
|---------|------|------|------|------|------|------|------|
| Shallow | 16.1 | 16.2 | 16.1 | 16.2 | 16.0 | 16.3 | 16.2 |
| Deep | 15.8 | 16.3 | 16.4 | 16.1 | 16.0 | 16.2 | 16.1 |
| No dung | 15.6 | 16.1 | 15.9 | 15.6 | 16.0 | 16.0 | 15.8 |
| Dung | 16.3 | 16.4 | 16.6 | 16.6 | 16.1 | 16.5 | 16.4 |
| Mean | 16.0 | 16.3 | 16.3 | 16.1 | 16.0 | 16.2 | 16.1 |

Total Sugar: cwt per acre

| | (a) | (b) and (c) | (a) | (b) and (c) | |
|---------|------|-------------|------|-------------|------|
| Shallow | 31.1 | 32.3 | 30.2 | 30.7 | 31.2 |
| Deep | 31.5 | 37.8 | 35.6 | 33.0 | 35.2 |
| No dung | 21.2 | 28.3 | 23.9 | 22.0 | 25.9 |
| Dung | 41.4 | 41.7 | 41.9 | 41.7 | 40.6 |
| Mean | 31.3 | 35.0 | 32.9 | 31.8 | 33.2 |

Tops: tons per acre

| | (a) | (b) and (c) | (a) | (b) and (c) | |
|---------|------|-------------|------|-------------|------|
| Shallow | 7.41 | 8.08 | 7.72 | 7.37 | 7.98 |
| Deep | 7.65 | 8.72 | 8.23 | 7.92 | 8.32 |
| No dung | 6.04 | 7.11 | 6.40 | 6.24 | 6.54 |
| Dung | 9.02 | 9.69 | 9.55 | 9.05 | 9.76 |
| Mean | 7.53 | 8.40 | 7.97 | 7.64 | 8.15 |

Plant Number: thousands per acre

| | | | | | | | |
|---------|------|------|------|------|------|------|------|
| Shallow | 28.8 | 28.2 | 28.4 | 28.4 | 28.8 | 28.6 | 28.6 |
| Deep | 28.9 | 29.0 | 28.9 | 28.9 | 28.8 | 29.0 | 28.9 |
| No dung | 28.6 | 28.5 | 28.6 | 28.7 | 28.4 | 28.4 | 28.6 |
| Dung | 29.1 | 28.7 | 28.7 | 28.6 | 29.2 | 29.2 | 28.9 |
| Mean | 28.9 | 28.6 | 28.7 | 28.7 | 28.8 | 28.8 | 28.7 |

Total Sugar Tops

| | | | |
|-----|-------|--------|--|
| (a) | ±1.00 | ±0.193 | for use in comparisons other than horizontal |
| (b) | ±1.16 | ±0.276 | for use in horizontal comparisons |
| (c) | ±1.29 | ±0.275 | as (a). |

Responses to treatments

Barley^{**}

| Response to | Mean | Ploughing | | Dung | | Phosphate | | Potash | |
|-------------|------|-----------|------|------|-------|-----------|-------|--------|-------|
| | | Shallow | Deep | Abs. | Pres. | Abs. | Pres. | Abs. | Pres. |

Grain: Mean yield 32.2 cwt per acre
(±1.13) (±1.60)

| | | | | | | | | | |
|------------------------|------|------|------|------|------|------|------|------|------|
| Ploughing deep-shallow | +0.5 | - | - | +0.2 | +0.8 | +1.7 | -0.7 | +0.3 | +0.7 |
| Dung | +3.3 | +3.0 | +3.6 | - | - | +3.1 | +3.5 | +3.2 | +3.4 |
| Phosphate | +2.9 | +4.1 | +1.7 | +2.7 | +3.1 | - | - | +2.2 | +3.6 |
| Potash | +1.5 | +1.3 | +1.7 | +1.4 | +1.6 | +0.8 | +2.2 | - | - |

Straw: Mean yield 39.5 cwt per acre

| | | | | | | | | | |
|------------------------|------|------|------|------|------|------|------|------|------|
| Ploughing deep-shallow | +2.6 | - | - | +0.5 | +4.7 | +3.1 | +2.1 | +2.5 | +2.7 |
| Dung | +6.6 | +4.5 | +8.7 | - | - | +6.1 | +7.1 | +6.8 | +6.4 |
| Phosphate | +3.0 | +3.5 | +2.5 | +2.5 | +3.5 | - | - | +2.3 | +3.7 |
| Potash | +3.7 | +3.6 | +3.8 | +3.9 | +3.5 | +3.0 | +4.4 | - | - |

Ley^{**}

Hay: Mean yield 68.3 cwt per acre

(±0.94) (±1.33)

| | | | | | | | | | |
|------------------------|------|------|------|------|------|------|------|-------|------|
| Ploughing deep-shallow | +2.6 | - | - | +2.4 | +2.8 | +3.9 | +1.3 | +1.7 | +3.5 |
| Dung | +6.5 | +6.3 | +6.7 | - | - | +9.4 | +3.6 | +10.9 | +2.1 |
| Phosphate | +0.4 | +1.7 | -0.9 | +3.3 | -2.5 | - | - | +0.6 | +0.2 |
| Potash | +0.4 | -0.5 | +1.3 | +4.8 | -4.0 | +0.6 | +0.2 | - | - |

Wheat^{*}

Grain: Mean yield 37.3 cwt per acre

(±0.75) (±1.07)

| | | | | | | | | | |
|------------------------|------|------|------|------|------|------|------|------|------|
| Ploughing deep-shallow | +3.2 | - | - | +2.8 | +3.6 | +1.5 | +4.9 | +2.9 | +3.5 |
| Dung | +1.5 | +1.1 | +1.9 | - | - | +1.7 | +1.3 | +0.8 | +2.2 |
| Phosphate | -0.6 | -2.3 | +1.1 | -0.4 | -0.8 | - | - | 0.0 | -1.2 |
| Potash | +0.6 | +0.3 | +0.9 | -0.1 | +1.3 | +1.2 | 0.0 | - | - |

Potatoes

Total tubers: Mean yield 6.99 tons per acre

(±0.464) (±0.656)

| | | | | | | | | | |
|------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Ploughing deep-shallow | +0.62 | - | - | +0.50 | +0.74 | +0.43 | +0.81 | +0.96 | +0.28 |
| Dung | +3.10 | +2.98 | +3.22 | - | - | +3.26 | +2.94 | +4.33 | +1.87 |
| Phosphate | +0.67 | +0.48 | +0.86 | +0.83 | +0.51 | - | - | +0.29 | +1.05 |
| Potash | +1.73 | +2.07 | +1.39 | +2.96 | +0.50 | +1.35 | +2.11 | - | - |

Percentage ware (1½" riddle): Mean 82.8

| | | | | | | | | | |
|------------------------|------|------|------|-------|------|------|------|-------|------|
| Ploughing deep-shallow | +1.9 | - | - | +3.2 | +0.6 | +1.5 | +2.3 | +4.5 | -0.7 |
| Dung | +7.2 | +8.5 | +5.9 | - | - | +6.2 | +8.2 | +12.6 | +1.8 |
| Phosphate | -1.7 | -2.1 | -1.3 | -2.7 | -0.7 | - | - | -2.9 | -0.5 |
| Potash | +4.8 | +7.4 | +2.2 | +10.2 | -0.6 | +3.6 | +6.0 | - | - |

*Cultivation treatments direct to wheat, manures to previous sugar beet.

**Treatments to previous Sugar beet.

55/Bb/1.6

Potatoes

| | Phosphate | | | Potash | | | Mean |
|-----------------------------|-----------|-------------|-----------|--------|-------------|-----------|------|
| | None | Ploughed in | In ridges | None | Ploughed in | In ridges | |
| Total tubers: tons per acre | | | | | | | |
| | (a) | (b) and (c) | | (a) | (b) and (c) | | |
| Shallow | 6.44 | 6.75 | 7.08 | 5.64 | 7.49 | 7.94 | 6.68 |
| Deep | 6.86 | 7.52 | 7.94 | 6.60 | 7.57 | 8.41 | 7.29 |
| No dung | 5.02 | 5.62 | 6.07 | 3.96 | 6.42 | 7.40 | 5.43 |
| Dung | 8.28 | 8.65 | 8.94 | 8.29 | 8.64 | 8.94 | 8.54 |
| Mean | 6.65 | 7.14 | 7.51 | 6.12 | 7.53 | 8.17 | 6.99 |

Percentage ware (1½" riddle)

| | | | | | | | |
|---------|------|------|------|------|------|------|------|
| Shallow | 82.9 | 80.9 | 80.7 | 78.1 | 84.7 | 86.4 | 81.8 |
| Deep | 84.4 | 84.6 | 81.6 | 82.6 | 86.0 | 83.8 | 83.8 |
| No dung | 80.5 | 78.7 | 77.0 | 74.1 | 84.0 | 84.6 | 79.2 |
| Dung | 86.7 | 86.9 | 85.2 | 86.7 | 86.6 | 85.6 | 86.4 |
| Mean | 83.6 | 82.8 | 81.1 | 80.4 | 85.3 | 85.1 | 82.8 |

Spring Oats

Responses to treatments to previous potatoes

| Response to | Mean | Ploughing | | Dung | | Phosphate | | Potash | |
|-------------|------|-----------|------|------|-------|-----------|-------|--------|-------|
| | | Shallow | Deep | Abs. | Pres. | Abs. | Pres. | Abs. | Pres. |

Grain (at 85% D.M.): Mean yield 38.4 cwt per acre
(±1.11) (±1.57)

| | | | | | | | | | |
|------------------------|------|------|------|------|------|------|------|------|------|
| Ploughing deep-shallow | -2.3 | - | - | -4.5 | -0.1 | -3.0 | -1.6 | -0.2 | -4.4 |
| Dung | +0.2 | -2.0 | +2.4 | - | - | +0.3 | +0.1 | -1.5 | +1.9 |
| Phosphate | -0.4 | -1.1 | +0.3 | -0.3 | -0.5 | - | - | +1.2 | -2.0 |
| Potash | +0.4 | +2.5 | -1.7 | -1.3 | +2.1 | +2.0 | -1.2 | - | - |

Total tubers

- (a) ±0.464 for use in comparisons other than horizontal
- (b) ±0.194 for use in horizontal comparisons
- (c) ±0.484 as (a).

Mean dry matter % as harvested

Barley, grain: Not recorded
Wheat, grain: 86.4
Spring oats, grain: 84.4

LEY AND ARABLE ROTATIONS

Highfield and Fosters Field 1955 - the 7th year.

For details of treatments, rotations, etc., see "Results of the Field Experiments" 1952, Section Bc/1, with the exception that the following alterations to the original scheme were introduced in the 1955 season:-

1. The third treatment crop of the arable rotation is now spring oats instead of barley.
2. On Highfield only, the rates of application of "nitro-chalk" to barley and oats are now: nil; 0.2 cwt N per acre. On Fosters the manuring of barley is unaltered (0.2; 0.4 cwt N) and oats receive these levels also.
3. Reseeded and old permanent grass are now grazed for 5 years; one cut of hay (followed by grazing of the aftermath) is taken in the sixth. In 1956 and later years the hay cut will be taken from plots in the blocks that are in the 1st treatment year; in 1955 however (in order to avoid hay cuts in successive years on certain plots) the hay cut was taken from plots in blocks in the 3rd treatment year. Rates of application of fertilizers are at the higher level (given under "Reseeded and Old Permanent Grass 3rd year" in "Results" 1952, on page 52/Bc/1.2) in hay years only.
4. On blocks in 2 out of the 6 phases viz:- the 1st and 2nd treatment years corrective dressings of muriate of potash were applied in order to compensate for different rates of withdrawal of potash by previous crops. Further dressings will be applied in future years, commencing as each block enters the 1st treatment year.

Rates of Application of Corrective Potash (K_2O : cwt per acre)

| Crop | Year of cycle | Rate |
|------------------------|-------------------------|----------------------------------|
| Cut grass | "1st treatment" | 2.4 (3 years previous cutting) |
| | "2nd treatment" | 1.2 (1 year previous cutting) |
| Grazed Ley and Arable | "1st and 2nd treatment" | Nil |
| | "1st treatment" | 2.4 (3 years previous Lucerne) |
| Lucerne | "2nd treatment" | 0.6 (1 year previous Lucerne) |
| | "1st treatment" | 2.4 (2 previous hay crops taken) |
| Permanent and Reseeded | "1st test" | None |
| | "2nd treatment" | 1.2 (1 previous hay crop taken) |
| | "2nd test" | None |

5. Each sub-plot of test crop potatoes is split into two for the application (in addition to the basal dressing) of all combinations of:

Phosphate: None; 0.9 cwt P_2O_5 per acre as superphosphate.

Potash: None; 0.9 cwt K_2O per acre as muriate of potash.

The PK interaction is confounded with block differences except on the Lucerne rotation where it is confounded with quarter plot differences.

For the succeeding barley crop similar dressings will be applied, but to different sub-plots, so that the totals of P_2O_5 and K_2O for the 2 crops will be equal. The barley will be harvested as hitherto by quarter plots.

6. Lucerne is now sown in rows 18" apart instead of 12".

Cultivations, etc.:

HIGHFIELD

1st year Treatment Crops

Cut grass: Ploughed: Oct 15, 1954. 1st dressing of supplementary K and nitrochalk applied: Apr 22, 1955. Basal PK applied: Apr 23. Seed sown at 33 lb per acre: Apr 25. 2nd dressing of supplementary K applied: July 9. Cut 3 times: June 30, July 26, Nov 21. Nitrochalk applied after each cut except the last.

Grazed ley: Ploughed: Oct 15, 1954. Nitrochalk applied: Apr 22, 1955. Basal PK applied: Apr 23. Seed sown at 44 lb per acre: Apr 25. Nitrochalk applied: July 16. Grazed: 5 circuits, June 22-Oct 6.

Lucerne: Ploughed: Oct 15, 1954. 1st dressing of supplementary K: Apr 22, 1955. Basal PK applied: Apr 23. Seed drilled 18" drills at 28 lb per acre: Apr 25. 2nd dressing of supplementary K applied: July 28. Cut 3 times: July 26, Sept 1, Nov 21. Variety: Du Puits.

Hay: Seeds undersown in barley at 28 lb per acre: Apr 22, 1954. Basal PK applied: Dec 30. Nitrochalk applied: Apr 15, 1955. Cut: June 10.

2nd year Treatment Crops

Cut grass: Basal PK applied: Dec 21, 1954. Supplementary K applied: Mar 11, 1955. Nitrochalk applied: Apr 18 and after each cut except the last. Cut 5 times: May 10, June 6, June 30, July 26, Nov 21.

Grazed ley: Basal PK applied: Dec 21, 1954. Nitrochalk applied: Apr 18 and July 9, 1955. Grazed: 8 circuits, Apr 27-Oct 3.

Lucerne: Basal PK applied: Dec 21, 1954. Supplementary K applied: Mar 11, 1955. Cut 4 times: June 24, July 26, Sept 1, Nov 21.

Potatoes: Ploughed: June 29, Aug 17 and Nov 16, 1954. Ridged: Apr 26, 1955. Basal PK, sulphate of ammonia, and dung applied, potatoes planted: Apr 29. For later cultivations see Potato Test Crop.

3rd year Treatment Crops

Cut grass: Basal PK applied: Dec 21, 1954. Nitrochalk applied: Apr 18, 1955 and after each cut except the last. Cut 5 times: May 19, June 13, June 30, July 26, Oct 14.

Grazed ley: Basal PK applied: Dec 21, 1954. Nitrochalk applied: Apr 18, 1955 and July 9. Grazed: 8 circuits, May 1-Oct 8.

Lucerne: Basal PK applied: Dec 21, 1954. Cut 4 times: June 14, July 26, Sept 1, Oct 12, 1955.

Oats: Ploughed: Nov 18, 1954. Nitrochalk applied: Mar 28, 1955. Seed drilled at $3\frac{1}{2}$ bushels per acre with basal PK: Mar 30. Combine harvested: Aug 12. Variety: Sun II.

1st Test Crop, Wheat

Ploughed leys: Oct 14, 1954. Ploughed after barley: Oct 15. Seed drilled at $2\frac{3}{4}$ bushels per acre with basal PK: Oct 20. Nitrochalk applied: May 10, 1955. Combine harvested: Aug 16. Variety: Yeoman.

55/Bc/1.3

2nd Test Crop, Potatoes

Ploughed: Oct 2, 1954. Ridged: Apr 26, 1955. Basal PK and additional P and K applied: Apr 28. Sulphate of ammonia and dung applied and potatoes planted: Apr 29. Earthed up: July 5. Sprayed with 20% sulphuric acid in 100 gallons per acre: Oct 4. Lifted: Oct 11. Variety: Majestic.

3rd Test Crop, Barley

Ploughed: Nov 18, 1954. Ground chalk applied to blocks 5 and 8: Dec 30. Nitrochalk applied: Mar 28, 1955. Seed drilled at 2 bushels per acre with basal PK: Mar 30. Combine harvested: Aug 12. Variety: Proctor.

Permanent Grasses. Basal PK applied to all plots: Dec 21, 1954. 5th year Reseeded, 5th experimental year of permanent grass, Blocks 9-12. Supplementary K applied to blocks 10 and 11: Mar 11, 1955. Nitrochalk applied: Apr 18 and July 9. Grazed: Permanent grass blocks 9 and 12 - 5 circuits, May 1-Oct 12; Remainder - 6 circuits Apr 27-Oct 15.

6th year Reseeded, 6th experimental year of permanent grass, Blocks 5-8. Blocks 5 and 8: Ground chalk applied: Dec 30, 1954. Nitrochalk applied: Apr 18 and July 16, 1955. Grazed: 5 circuits, May 5-Oct 7. Blocks 6 and 7: Nitrochalk applied: Apr 15, 1955. Cut for hay: June 27. Nitrochalk applied: June 28. Grazed: 2 circuits, July 19-Oct 7.

7th year Reseeded, 7th experimental year of permanent grass, Blocks 1-4. Nitrochalk applied: Apr 22 and July 2, 1955. Supplementary K applied to blocks 2 and 3: Mar 11 and July 20. Grazed: 5 circuits, May 9-Sept 29.

FOSTERS

1st year Treatment Crops

Cut grass: Ploughed: Oct 13, 1954. 1st dressing of supplementary K and nitrochalk applied: Apr 22, 1955. Basal PK applied and seed sown at 33 lb per acre: Apr 23. 2nd dressing of supplementary K applied: July 8. Cut 3 times: July 1, July 27, Nov 19. Nitrochalk applied after each cut except the last.

Grazed ley: Ploughed: Oct 13, 1954. Nitrochalk applied: Apr 22, 1955. Basal PK applied and seed sown at 44 lb per acre: Apr 23. Nitrochalk applied: July 22. Grazed: 2 circuits, July 17-Sept 26.

Lucerne: Ploughed: Oct 13, 1954. 1st dressing of supplementary K applied: Apr 22, 1955. Basal PK applied and seed sown at 28 lb per acre: Apr 23. 2nd dressing of supplementary K applied: July 28. Cut 3 times: July 27, Aug 31, Nov 19. Variety: Du Puits.

Hay: Seeds undersown in barley at 28 lb per acre: Apr 21, 1954. Basal PK applied: Dec 23. Nitrochalk applied: Apr 15, 1955. Cut: June 6.

55/Bc/1.4

2nd year Treatment Crops

Cut grass: Basal PK applied: Dec 22, 1954. Supplementary K applied: Mar 11, 1955. Nitrochalk applied: Apr 18 and after each cut except the last. Cut 5 times: May 10, June 6, July 1, July 27, Nov 19.

Grazed ley: Basal PK applied: Dec 22, 1954. Nitrochalk applied: Apr 18 and July 22, 1955. Grazed: 6 circuits, Apr 28-Oct 4.

Lucerne: Basal PK applied: Dec 22, 1954. Supplementary K applied: Mar 11, 1955. Cut 4 times: June 23, July 27, Aug 31, Nov 19.

Potatoes: Ploughed: June 28, Aug 17 and Nov 4, 1954. Basal PK applied: Apr 28, 1955. Sulphate of ammonia and dung applied: May 2. For later cultivations see Potato Test Crop.

3rd year Treatment Crops

Cut grass: Basal PK applied: Dec 22, 1954. Nitrochalk applied: Apr 19, 1955 and after each cut except the last. Cut 5 times: May 19, June 13, July 1, July 27, Oct 14.

Grazed ley: Basal PK applied: Dec 22, 1954. Nitrochalk applied: Apr 19 and July 22, 1955. Grazed: 6 circuits, May 2-Sept 30.

Lucerne: Basal PK applied: Dec 22, 1954. Cut 4 times: June 14, July 27, Aug 31, Oct 12.

Oats: Ploughed: Oct 18, 1954. Nitrochalk applied: Mar 28. Seed drilled at $3\frac{1}{2}$ bushels per acre with basal PK: Mar 30. Combine harvested: Aug 9. Variety: Sun II.

1st Test Crop, Wheat

Ploughed: Oct 12, 1954. Seed drilled at $2\frac{3}{4}$ bushels with basal PK: Oct 20, 1954. Nitrochalk applied: May 11, 1955. Combine harvested: Aug 16. Variety: Yeoman.

2nd Test Crop, Potatoes

Ploughed: Oct 1, 1954. Ridged: Apr 27, 1955. Basal PK applied: Apr 28. Additional P and K, dung and sulphate of ammonia: May 2. Potatoes planted: May 3. Earthed up: June 29. Sprayed with 20% sulphuric acid in 100 gallons: Sept 30. Lifted: Oct 11. Variety: Majestic.

3rd Test Crop, Barley

Ploughed: Oct 18, 1954. Nitrochalk applied: Mar 28, 1955. Seed drilled at 2 bushels per acre with basal PK: Mar 30. Sprayed with MCPA, $2\frac{1}{2}$ pints in 40 gallons per acre: June 8. Combine harvested: Aug 9. Variety: Proctor.

Permanent grasses. Basal PK applied to all plots: Dec 22, 1954.

5th year reseeded grass, Blocks 6, 10, 11, 12.

Supplementary K applied to blocks 10 and 12: Mar 11, 1955. Nitrochalk applied: Apr 4 and July 15. Grazed: Blocks 6 and 10, 5 circuits, Apr 28-Oct 14; Blocks 11 and 12, 4 circuits, May 2-Oct 10.

55/Bc/1.5

6th year reseeded grass, Blocks 5,7,8,9.

Blocks 5 and 7: Nitrochalk applied: Apr 18 and July 7, 1955.

Grazed: 5 circuits, May 16-Oct 14.

Blocks 8 and 9: Nitrochalk applied: Apr 18 and June 27, 1955. Cut for hay: June 27. Aftermath grazed: 1 circuit, Oct 2-Oct 6.

7th year reseeded grass, Blocks 1, 2, 3, 4.

Supplementary K applied to blocks 2 and 4: Mar 11, 1955 and

July 15, 1955. Nitrochalk: Apr 18 and July 15. Grazed: 5 circuits, May 10-Sept 30.

Standard errors per $\frac{1}{4}$ plot. Test crops.

Wheat, grain (at 85% dry matter). Highfield: 2.10 cwt per acre or 4.6% (13 d.f.)

Fosters: 2.70 cwt per acre or 7.7% (13 d.f.)

Potatoes, total tubers: Highfield $\frac{1}{4}$ plot: 0.636 tons per acre or 6.2% (14 d.f.)

Highfield $\frac{1}{8}$ plot: 0.702 tons per acre or 6.9% (20 d.f.)

Fosters $\frac{1}{4}$ plot: 0.465 tons per acre or 5.8% (14 d.f.)

Fosters $\frac{1}{8}$ plot: 0.594 tons per acre or 7.4% (20 d.f.)

Barley, grain (at 85% D.M.). Highfield: 2.55 cwt per acre or 5.2% (15 d.f.)

Fosters: 2.08 cwt per acre or 4.5% (15 d.f.)

55/Bc/1.6

Summary of Results

Wheat 1st test crop

| N: cwt per acre | Treatment crops 1952-1954 | | | | Mean |
|---|---------------------------|------|--------------|--------------------|------------------------|
| | Lucerne | Ley | Cut Grass | Arable with hay | |
| Grain (at 85% Dry Matter): cwt per acre | | | | | |
| <u>Highfield</u> | | | | | |
| Mean | 48.0 | 48.9 | 38.9 | 47.9 | 45.9 |
| To test crop | | | | | |
| 0.3 | 45.2 | 48.9 | 38.1 | 47.4 | 44.9 |
| 0.6 | 50.8 | 48.9 | 39.7 | 48.3 | 46.9 |
| Difference (± 1.49) | +5.6 | 0.0 | +1.6 | +0.9 | +2.0 (± 0.74) |
| To treatment crops | | | | | |
| Single rate | | 48.0 | 40.9 | 47.5 | 45.5 |
| Double rate | | 49.9 | 36.9 | 48.2 | 45.0 |
| Difference (± 1.49) | | +1.9 | -4.0 | +0.7 | -0.5 (± 0.86) |
| <u>Fosters</u> | | | | | |
| Mean | 38.2 | 34.8 | 34.6 | 32.9 | 35.1 |
| To test crop | | | | | |
| 0.3 | 36.3 | 33.0 | 34.9 | 30.6 | 33.7 |
| 0.6 | 40.1 | 36.5 | 34.3 | 35.2 | 36.5 |
| Difference (± 1.91) | +3.8 | +3.5 | -0.6 | +4.6 | +2.8 (± 0.95) |
| To treatment crops | | | | | |
| Single rate | | 35.6 | 35.2 | 33.1 | 34.6 |
| Double rate | | 33.9 | 34.0 | 32.7 | 33.5 |
| Difference (± 1.91) | | -1.7 | -1.2 | -0.4 | -1.1 (± 1.10) |

55/Bc/1.7

Wheat 1st test crop

| N: cwt per acre | Excluding Lucerne N to previous treatment crop | | | Arable with hay only Dung to potatoes 1953: tons per acre | | |
|-----------------|--|----------------|------|--|----|------|
| | Single rate | Double rate | Mean | None | 12 | Mean |

Grain (at 85% Dry Matter): cwt per acre

Highfield

| | | | | | | |
|--------------------------------|--------------|------|--------------|--------------|------|--------------|
| To test crop | (± 0.86) | | (± 0.61) | (± 1.49) | | (± 1.05) |
| 0.3 | 45.5 | 44.1 | 44.8 | 45.3 | 49.5 | 47.4 |
| 0.6 | 45.5 | 45.9 | 45.7 | 49.0 | 47.7 | 48.3 |
| Mean | 45.5 | 45.0 | 45.2 | 47.1 | 48.6 | 47.9 |
| | (± 0.61) | | | (± 1.05) | | |
| To previous treatment crops | | | | (± 1.49) | | (± 1.05) |
| Single rate | | | | 45.8 | 49.3 | 47.5 |
| Double rate | | | | 48.5 | 48.0 | 48.2 |
| Mean | | | | 47.1 | 48.6 | 47.9 |
| | | | | (± 1.05) | | |

Mean dry matter % as harvested: 83.7

Fosters

| | | | | | | |
|--------------------------------|--------------|------|--------------|--------------|------|--------------|
| To test crop | (± 1.10) | | (± 0.78) | (± 1.91) | | (± 1.35) |
| 0.3 | 32.8 | 32.9 | 32.8 | 29.0 | 32.2 | 30.6 |
| 0.6 | 36.5 | 34.2 | 35.4 | 35.2 | 35.2 | 35.2 |
| Mean | 34.6 | 33.5 | 34.1 | 32.1 | 33.7 | 32.9 |
| | (± 0.78) | | | (± 1.35) | | |
| To previous treatment crops | | | | (± 1.91) | | (± 1.35) |
| Single rate | | | | 31.4 | 34.9 | 33.1 |
| Double rate | | | | 32.9 | 32.5 | 32.7 |
| Mean | | | | 32.1 | 33.7 | 32.9 |
| | | | | (± 1.35) | | |

Mean dry matter % as harvested: 81.9

55/Bc/1.8

Potatoes 2nd test crop. Total tubers: tons per acre

| | Treatment crops 1951-1953 | | | | Mean |
|---|---------------------------|-------|-----------|-----------------|--------------------------|
| | Lucerne | Ley | Cut Grass | Arable with hay | |
| <u>Highfield</u> | | | | | |
| Mean | 10.05 | 10.48 | 9.78 | 10.53 | 10.21 |
| N: cwt per acre | | | | | |
| 0.5 | 9.50 | 10.18 | 9.78 | 10.28 | 9.94 |
| 1.0 | 10.60 | 10.77 | 9.78 | 10.78 | 10.48 |
| Difference (± 0.450) | +1.10 | +0.59 | 0.00 | +0.50 | +0.54 (± 0.225) |
| Dung: tons per acre | | | | | |
| None | 8.76 | 10.09 | 8.88 | 9.74 | 9.37 |
| 12 | 11.34 | 10.86 | 10.69 | 11.32 | 11.05 |
| Difference (± 0.450) | +2.58 | +0.77 | +1.81 | +1.58 | +1.68 (± 0.225) |
| P ₂ O ₅ : cwt per acre [‡] | | | | | |
| 0.9 | 9.77 | 10.53 | 9.72 | 10.40 | 10.11 |
| 1.8 | 10.33 | 10.42 | 9.85 | 10.66 | 10.31 |
| Difference (± 0.351) | +0.56 | -0.11 | +0.13 | +0.26 | +0.20 (± 0.175) |
| K ₂ O: cwt per acre [‡] | | | | | |
| 0.9 | 9.38 | 9.94 | 8.95 | 10.12 | 9.60 |
| 1.8 | 10.72 | 11.01 | 10.62 | 10.94 | 10.82 |
| Difference (± 0.351) | +1.34 | +1.07 | +1.67 | +0.82 | +1.22 (± 0.175) |
| <u>Fosters</u> | | | | | |
| Mean | 8.62 | 8.29 | 7.35 | 7.65 | 7.97 |
| N: cwt per acre | | | | | |
| 0.5 | 8.73 | 8.13 | 7.64 | 7.58 | 8.02 |
| 1.0 | 8.50 | 8.44 | 7.05 | 7.71 | 7.93 |
| Difference (± 0.329) | -0.23 | +0.31 | -0.59 | +0.13 | -0.09 (± 0.164) |
| Dung: tons per acre | | | | | |
| None | 7.81 | 7.97 | 6.22 | 7.26 | 7.32 |
| 12 | 9.42 | 8.60 | 8.48 | 8.03 | 8.63 |
| Difference (± 0.329) | +1.61 | +0.63 | +2.26 | +0.77 | +1.31 (± 0.164) |
| P ₂ O ₅ : cwt per acre [‡] | | | | | |
| 0.9 | 8.39 | 8.04 | 7.08 | 7.59 | 7.78 |
| 1.8 | 8.84 | 8.53 | 7.62 | 7.71 | 8.17 |
| Difference (± 0.297) | +0.45 | +0.49 | +0.54 | +0.12 | +0.39 (± 0.148) |
| K ₂ O: cwt per acre [‡] | | | | | |
| 0.9 | 8.29 | 8.12 | 6.87 | 7.52 | 7.70 |
| 1.8 | 8.94 | 8.46 | 7.82 | 7.77 | 8.25 |
| Difference (± 0.297) | +0.65 | +0.34 | +0.95 | +0.25 | +0.55 (± 0.148) |

[‡]Including basal dressing.

55/Bc/1.9

Potatoes 2nd test crop. Total tubers: tons per acre

| | Dung: tons per acre | | P ₂ O ₅ : cwt per acre [‡] | | K ₂ O: cwt per acre [‡] | |
|---|---------------------|-------|---|-------|---|-------|
| | None | 12 | 0.9 | 1.8 | 0.9 | 1.8 |
| <u>Highfield</u> | | | | | | |
| | (± 0.225) | | (1) and (2) | | (1) and (2) | |
| N: cwt per acre | | | | | | |
| 0.5 | 9.19 | 10.68 | 9.94 | 9.93 | 9.18 | 10.69 |
| 1.0 | 9.55 | 11.42 | 10.27 | 10.69 | 10.01 | 10.96 |
| | | | (1) and (2) | | (1) and (2) | |
| Dung: tons per acre | | | | | | |
| None | | | 9.37 | 9.37 | 8.49 | 10.25 |
| 12 | | | 10.84 | 11.26 | 10.71 | 11.40 |
| <u>Lucerne rotation only</u> | | | | | | |
| | | | K ₂ O: cwt per acre [‡] | | | |
| | | | 0.9 | 1.8 | Mean | |
| <u>P₂O₅: cwt per acre[‡]</u> | | | | | | |
| | | | (3) and (4) | | | |
| 0.9 | | | 9.12 | 10.42 | 9.77 | |
| 1.8 | | | 9.65 | 11.01 | 10.33 | |
| Mean | | | 9.38 | 10.72 | 10.05 | |
| <u>Dung: tons per acre</u> | | | | | | |
| | | | | | | |
| None | | | 0.9 | 1.8 | 0.9 | 1.8 |
| 12 | | | | | | |
| <u>Fosters</u> | | | | | | |
| | (± 0.164) | | (1) and (2) | | (1) and (2) | |
| N: cwt per acre | | | | | | |
| 0.5 | 7.20 | 8.84 | 7.86 | 8.18 | 7.83 | 8.21 |
| 1.0 | 7.43 | 8.43 | 7.69 | 8.16 | 7.57 | 8.28 |
| | | | (1) and (2) | | (1) and (2) | |
| Dung: tons per acre | | | | | | |
| None | | | 7.31 | 7.32 | 7.00 | 7.63 |
| 12 | | | 8.24 | 9.02 | 8.40 | 8.86 |
| <u>Lucerne rotation only</u> | | | | | | |
| | | | K ₂ O: cwt per acre [‡] | | | |
| | | | 0.9 | 1.8 | Mean | |
| <u>P₂O₅: cwt per acre[‡]</u> | | | | | | |
| | | | (3) and (4) | | | |
| 0.9 | | | 7.96 | 8.83 | 8.39 | |
| 1.8 | | | 8.63 | 9.05 | 8.84 | |
| Mean | | | 8.29 | 8.94 | 8.62 | |

[‡]Including basal dressing

Highfield

Fosters

- | | | |
|-----------------|-----------------|---|
| (1) ± 0.175 | (1) ± 0.148 | for use in horizontal and interaction comparisons |
| (2) ± 0.202 | (2) ± 0.157 | for use in all others |
| (3) ± 0.450 | (3) ± 0.329 | for use only in testing the PK interaction |
| (4) ± 0.403 | (4) ± 0.313 | for use in all other comparisons. |

Potatoes 2nd test crop. Percentage ware (1½" riddle)

| | Treatment crops 1951-1953 | | | | Mean |
|---|---------------------------|------|--------------|--------------------|------|
| | Lucerne | Ley | Cut Grass | Arable with hay | |
| <u>Highfield</u> | | | | | |
| Mean | 82.5 | 82.7 | 81.0 | 82.3 | 82.1 |
| N: cwt per acre | | | | | |
| 0.5 | 82.2 | 80.8 | 81.2 | 82.3 | 81.6 |
| 1.0 | 82.8 | 84.6 | 80.7 | 82.4 | 82.6 |
| Difference | +0.6 | +3.8 | -0.5 | +0.1 | +1.0 |
| Dung: tons per acre | | | | | |
| None | 81.4 | 81.3 | 79.1 | 81.4 | 80.8 |
| 12 | 83.5 | 84.1 | 82.8 | 83.2 | 83.4 |
| Difference | +2.1 | +2.8 | +3.7 | +1.8 | +2.6 |
| P ₂ O ₅ : cwt per acre [*] | | | | | |
| 0.9 | 82.5 | 82.2 | 81.8 | 83.3 | 82.4 |
| 1.8 | 82.4 | 83.2 | 80.1 | 81.4 | 81.8 |
| Difference | -0.1 | +1.0 | -1.7 | -1.9 | -0.6 |
| K ₂ O: cwt per acre [*] | | | | | |
| 0.9 | 81.2 | 82.8 | 78.9 | 81.3 | 81.0 |
| 1.8 | 83.7 | 82.6 | 83.0 | 83.4 | 83.2 |
| Difference | +2.5 | -0.2 | +4.1 | +2.1 | +2.2 |
| <u>Fosters</u> | | | | | |
| Mean | 83.0 | 84.7 | 83.6 | 83.2 | 83.6 |
| N: cwt per acre | | | | | |
| 0.5 | 84.9 | 84.5 | 84.9 | 84.2 | 84.6 |
| 1.0 | 81.0 | 85.0 | 82.2 | 82.1 | 82.6 |
| Difference | -3.9 | +0.5 | -2.7 | -2.1 | -2.0 |
| Dung: tons per acre | | | | | |
| None | 82.5 | 85.1 | 80.9 | 82.7 | 82.8 |
| 12 | 83.4 | 84.4 | 86.2 | 83.7 | 84.4 |
| Difference | +0.9 | -0.7 | +5.3 | +1.0 | +1.6 |
| P ₂ O ₅ : cwt per acre [*] | | | | | |
| 0.9 | 82.7 | 84.7 | 83.6 | 83.3 | 83.6 |
| 1.8 | 83.2 | 84.8 | 83.6 | 83.1 | 83.7 |
| Difference | +0.5 | +0.1 | 0.0 | -0.2 | +0.1 |
| K ₂ O: cwt per acre [*] | | | | | |
| 0.9 | 83.6 | 85.0 | 82.6 | 82.4 | 83.4 |
| 1.8 | 82.3 | 84.5 | 84.6 | 84.0 | 83.8 |
| Difference | -1.3 | -0.5 | +2.0 | +1.6 | +0.4 |

*Including basal dressing.

Potatoes 2nd test crop. Percentage ware (1½" riddle)

| Dung: tons per acre | | P ₂ O ₅ : cwt per acre [‡] | | K ₂ O: cwt per acre [‡] | |
|---------------------|----|---|-----|---|-----|
| None | 12 | 0.9 | 1.8 | 0.9 | 1.8 |

Highfield

| | | | | | | |
|---------------------|------|------|------|------|------|------|
| N: cwt per acre | | | | | | |
| 0.5 | 80.7 | 82.5 | 82.2 | 81.0 | 80.4 | 82.8 |
| 1.0 | 80.9 | 84.3 | 82.6 | 82.6 | 81.7 | 83.5 |
| Dung: tons per acre | | | | | | |
| None | | | 81.6 | 80.0 | 79.2 | 82.4 |
| 12 | | | 83.2 | 83.6 | 82.8 | 84.0 |

| <u>Lucerne rotation only</u> | K ₂ O: cwt per acre [‡] | | Mean |
|---|---|------|------|
| | 0.9 | 1.8 | |
| P ₂ O ₅ : cwt per acre [‡] | | | |
| 0.9 | 81.4 | 83.6 | 82.5 |
| 1.8 | 81.1 | 83.8 | 82.4 |
| Mean | 81.2 | 83.7 | 82.5 |

| Dung: tons per acre | | P ₂ O ₅ : cwt per acre [‡] | | K ₂ O: cwt per acre [‡] | |
|---------------------|----|---|-----|---|-----|
| None | 12 | 0.9 | 1.8 | 0.9 | 1.8 |

Fosters

| | | | | | | |
|---------------------|------|------|------|------|------|------|
| N: cwt per acre | | | | | | |
| 0.5 | 83.8 | 85.5 | 84.5 | 84.8 | 85.3 | 84.0 |
| 1.0 | 81.8 | 83.4 | 82.6 | 82.5 | 81.4 | 83.7 |
| Dung: tons per acre | | | | | | |
| None | | | 83.1 | 82.5 | 82.5 | 83.1 |
| 12 | | | 84.0 | 84.8 | 84.3 | 84.6 |

| <u>Lucerne rotation only</u> | K ₂ O: cwt per acre [‡] | | Mean |
|---|---|------|------|
| | 0.9 | 1.8 | |
| P ₂ O ₅ : cwt per acre [‡] | | | |
| 0.9 | 81.8 | 83.6 | 82.7 |
| 1.8 | 85.4 | 81.0 | 83.2 |
| Mean | 83.6 | 82.3 | 83.0 |

[‡]Including basal dressing.

Barley 3rd test crop. Grain (at 85% Dry Matter): cwt per acre

| | Treatment crops 1950-1952 | | | | Mean |
|---|---------------------------|------|-----------|-----------------|------------------------|
| | Lucerne | Ley | Cut Grass | Arable with hay | |
| <u>Highfield</u> | | | | | |
| Mean | 50.6 | 48.6 | 48.6 | 49.0 | 49.2 |
| N: cwt per acre | | | | | |
| None | 49.7 | 48.5 | 50.6 | 46.8 | 48.9 |
| 0.2 | 51.5 | 48.7 | 46.6 | 51.2 | 49.5 |
| Difference (± 1.80) | +1.8 | +0.2 | -4.0 | +4.4 | +0.6 (± 0.90) |
| Dung to potatoes 1954: tons per acre | | | | | |
| None | 50.8 | 48.2 | 47.9 | 47.9 | 48.7 |
| 12 | 50.4 | 49.1 | 49.3 | 50.1 | 49.7 |
| Difference (± 1.80) | -0.4 | +0.9 | +1.4 | +2.2 | +1.0 (± 0.90) |

| | N: cwt per acre | |
|---|-----------------|------|
| | None | 0.2 |
| Dung to potatoes 1954: tons per acre | | |
| None | 48.4 | 49.0 |
| 12 | 49.5 | 50.0 |
| | (± 0.90) | |

| | Treatment crops 1950-1952 | | | | Mean |
|---|---------------------------|------|-----------|-----------------|------------------------|
| | Lucerne | Ley | Cut Grass | Arable with hay | |
| <u>Fosters</u> | | | | | |
| Mean | 48.4 | 46.5 | 45.1 | 45.5 | 46.4 |
| N: cwt per acre | | | | | |
| 0.2 | 47.3 | 44.5 | 43.6 | 44.7 | 45.0 |
| 0.4 | 49.5 | 48.5 | 46.7 | 46.4 | 47.8 |
| Difference (± 1.47) | +2.2 | +4.0 | +3.1 | +1.7 | +2.8 (± 0.73) |
| Dung to potatoes 1954: tons per acre | | | | | |
| None | 48.5 | 45.9 | 44.8 | 44.9 | 46.0 |
| 12 | 48.3 | 47.1 | 45.5 | 46.2 | 46.8 |
| Difference (± 1.47) | -0.2 | +1.2 | +0.7 | +1.3 | +0.8 (± 0.73) |

| | N: cwt per acre | |
|---|-----------------|------|
| | 0.2 | 0.4 |
| Dung to potatoes 1954: tons per acre | | |
| None | 43.8 | 48.2 |
| 12 | 46.2 | 47.4 |
| | (± 0.73) | |

Mean dry matter % as harvested: Highfield: 78.7, Fosters: 79.6

55/Bc/1.13

Treatment crops Arable and Hay rotation
(values based on Mean of 2 sub plots only)

| | Highfield | | | Fosters | | |
|--|---------------------------------|-------------|------|---------------------------------|-------------|------|
| | N: cwt per acre applied in 1955 | | | N: cwt per acre applied in 1955 | | |
| | Single rate | Double rate | Mean | Single rate | Double rate | Mean |
| Hay (dry matter): cwt per acre | | | | | | |
| No dung | 51.4 | 53.1 | 52.3 | 51.4 | 55.7 | 53.5 |
| Dung in 1953 | 48.5 | 45.7 | 47.1 | 52.7 | 57.0 | 54.9 |
| Mean | 50.0 | 49.4 | 49.7 | 52.0 | 56.4 | 54.2 |
| Potatoes, total tubers: tons per acre | | | | | | |
| No dung | 8.83 | 8.61 | 8.72 | 6.96 | 7.34 | 7.15 |
| Dung in 1955 | 10.29 | 9.68 | 9.99 | 9.29 | 9.28 | 9.29 |
| Mean | 9.56 | 9.15 | 9.35 | 8.13 | 8.31 | 8.22 |
| Potatoes, percentage ware (1½" riddle) | | | | | | |
| No dung | 81.2 | 78.4 | 79.8 | 79.3 | 80.8 | 80.0 |
| Dung in 1955 | 83.9 | 83.5 | 83.7 | 87.2 | 86.0 | 86.6 |
| Mean | 82.6 | 81.0 | 81.8 | 83.2 | 83.4 | 83.3 |
| Oats, grain: cwt per acre | | | | | | |
| | None | 0.2 | | 0.2 | 0.4 | |
| | | | | (at 85% Dry Matter) | | |
| No dung | 50.6 | 52.9 | 51.8 | 39.7 | 45.9 | 42.8 |
| Dung in 1954 | 50.2 | 49.7 | 49.9 | 40.6 | 45.1 | 42.9 |
| Mean | 50.4 | 51.3 | 50.8 | 40.2 | 45.5 | 42.8 |

Highfield, Oats, Mean dry matter % as harvested: 83.2
Fosters, Oats, Mean dry matter % as harvested: 81.9

55/Bc/1.14

Cut grass. Dry Matter: cwt per acre

| Corrective dressing of K ₂ O: cwt per acre | 2.4 | Highfield | | | | | Fosters | | | | |
|---|-----|-----------------------------|-------------|-------------------------------------|----|------|-----------------------------|-------------|-------------------------------------|----|------|
| | | N: to previous 3 test crops | | Dung to potatoes 1953 tons per acre | | Mean | N: to previous 3 test crops | | Dung to potatoes 1953 tons per acre | | Mean |
| | | Single rate | Double rate | None | 12 | | Single rate | Double rate | None | 12 | |

| | | | | | | | | | | | |
|----------------------------|------|------|------|------|------|------|------|------|------|------|--|
| N(1) to cut grass (3 cuts) | | | | | | | | | | | |
| Single rate | 30.3 | 31.2 | 32.9 | 28.5 | 30.7 | 7.9 | 9.8 | 9.1 | 8.6 | 8.8 | |
| Double rate | 38.0 | 36.9 | 38.7 | 36.2 | 37.4 | 13.8 | 10.4 | 13.2 | 11.0 | 12.1 | |
| N to test crops | | | | | | | | | | | |
| Single rate | | | 37.4 | 30.8 | 34.1 | | | 11.8 | 9.9 | 10.8 | |
| Double rate | | | 34.2 | 33.8 | 34.0 | | | 10.5 | 9.7 | 10.1 | |
| Mean | | | 35.8 | 32.3 | 34.1 | | | 11.1 | 9.8 | 10.5 | |

| | | Highfield | | | Fosters | | |
|--------------------------|------|--------------------|-------------|------|--------------------|-------------|------|
| | | N to cut grass (1) | | | N to cut grass (1) | | |
| | | Single rate | Double rate | Mean | Single rate | Double rate | Mean |
| <u>2nd year</u> (5 cuts) | 1.2 | 46.8 | 60.7 | 53.7 | 37.8 | 42.6 | 40.2 |
| <u>3rd year</u> (5 cuts) | None | 32.2 | 45.7 | 39.0 | 31.9 | 36.5 | 34.2 |

(1) 0.15 v. 0.3 cwt N as Nitrochalk for every cut.

Lucerne. Dry Matter: cwt per acre

| Corrective dressing of K ₂ O: cwt per acre | 2.4 | Highfield | | | Fosters | | |
|---|-----|----------------------------|-------------|------|----------------------------|-------------|------|
| | | N to 3 previous test crops | | | N to 3 previous test crops | | |
| | | Single rate | Double rate | Mean | Single rate | Double rate | Mean |

| | | | | | | | |
|-----------------------|--|------|------|------|------|------|------|
| Dung to potatoes 1953 | | | | | | | |
| None | | 24.4 | 23.2 | 23.8 | 21.6 | 23.5 | 22.6 |
| 12 tons | | 25.9 | 24.4 | 25.2 | 24.4 | 23.7 | 24.0 |
| Mean | | 25.2 | 23.8 | 24.5 | 23.0 | 23.6 | 23.3 |

| | | | | | | | |
|--------------------------|------|--|--|-------|--|--|-------|
| <u>2nd year</u> (4 cuts) | 0.6 | | | 109.8 | | | 111.9 |
| <u>3rd year</u> (4 cuts) | None | | | 79.6 | | | 106.9 |

55/Bo/1.15

Grazed Ley. Dry Matter: cwt per acre (estimated from sample cuts)

| | Highfield | | | Fosters | | |
|----------|--------------------------|------|------|--------------------------|------|------|
| | N: cwt per acre (yearly) | | | N: cwt per acre (yearly) | | |
| | 0.15 | 0.30 | Mean | 0.15 | 0.30 | Mean |
| 1st year | 26.9 | 27.2 | 27.1 | 10.4 | 10.3 | 10.4 |
| 2nd year | 56.5 | 57.5 | 57.0 | 38.7 | 36.5 | 37.6 |
| 3rd year | 55.8 | 66.5 | 61.2 | 40.7 | 42.3 | 41.5 |

Reseeded Grass. Dry Matter: cwt per acre

| Corrective dressing of K ₂ O: cwt per acre | Cut for hay | | | Grazed Estimated from sampling cuts | | |
|---|-------------|-------------|------|-------------------------------------|-------------|------|
| | N | | | N | | |
| | Single rate | Double rate | Mean | Single rate | Double rate | Mean |

Highfield

| | | | | | | |
|--------------|------|------|------|-------------------|-------------------|-------------------|
| 5th year | | | | | | |
| None | | | | 49.5 | 55.7 | 52.6 |
| 1.2 | | | | 50.4 | 52.0 | 51.2 |
| Mean | | | | 49.9 | 53.9 | 51.9 |
| 6th year | | | | | | |
| Blocks 5 & 8 | | | | 42.1 | 42.1 | 42.1 |
| Blocks 6 & 7 | 64.5 | 65.6 | 65.0 | 23.7 [‡] | 36.4 [‡] | 30.1 [‡] |
| 7th year | | | | | | |
| None | | | | 27.1 | 38.1 | 32.6 |
| 2.4 | | | | 34.0 | 33.9 | 33.9 |
| Mean | | | | 30.6 | 36.0 | 33.3 |

Fosters

| | | | | | | |
|--------------|------|------|------|-------------------|-------------------|-------------------|
| 5th year | | | | | | |
| None | | | | 40.9 | 34.9 | 37.9 |
| 1.2 | | | | 38.2 | 44.0 | 41.1 |
| Mean | | | | 39.5 | 39.4 | 39.5 |
| 6th year | | | | | | |
| Blocks 5 & 7 | | | | 27.7 [‡] | 29.1 [‡] | 28.4 [‡] |
| Blocks 8 & 9 | 66.7 | 68.5 | 67.6 | 21.7 [‡] | 17.5 [‡] | 19.6 [‡] |
| 7th year | | | | | | |
| None | | | | 30.3 | 35.1 | 32.7 |
| 2.4 | | | | 38.2 | 30.4 | 34.3 |
| Mean | | | | 34.3 | 32.7 | 33.5 |

[‡]Aftermath grazing.

55/Bc/1.16

Permanent Grass. Dry Matter: cwt per acre

| Corrective dressing of K ₂ O: cwt per acre | <u>Highfield</u> | | | Grazed | | |
|--|------------------|----------------|------|--------------------------------|-------------------|-------------------|
| | Cut for hay | | Mean | Estimated from sample cuts: | | |
| | Single rate | Double rate | | Single rate | Double rate | Mean |
| 5th experimental year Blocks 9-12 | | | | | | |
| None | | | | 36.6 | 39.3 | 38.0 |
| 1.2 | | | | 42.7 | 45.1 | 43.9 |
| Mean | | | | 39.7 | 42.2 | 40.9 |
| 6th experimental year Blocks 5&8 | | | | 32.4 | 38.2 | 35.3 |
| Blocks 6&7 | 51.0 | 60.7 | 55.9 | 25.3 [*] | 25.2 [*] | 25.2 [*] |
| 7th experimental year Blocks 1-4 | | | | 37.3 | 34.8 | 36.1 |
| None | | | | 30.9 | 34.0 | 32.4 |
| 2.4 | | | | | | |
| Mean | | | | 34.1 | 34.4 | 34.3 |

^{*}Aftermath grazing.

55/Bd/1.1

GREEN MANURING EXPERIMENT

Woburn Stackyard - 1955, the 2nd year of revised scheme.

For details of treatments etc. see "Results of the Field Experiments" 1954, Section Bd/1.

Area of each plot: 0.0395 acre. Area harvested: Barley, 0.0395; Potatoes, 0.0237 acre.

Cultivations, etc.:

Green manures after potatoes: Trefoil at 30 lb per acre, ryegrass at 40 lb per acre, sown: Aug 3, 1954. Varieties: Trefoil - English; Ryegrass - Western Wolths.

Barley: "Fallow" and "early" green manure plots ploughed: Dec 22, 1954. "Late" green manure plots ploughed: Mar 14, 1955. Ground chalk at 20 cwt per acre applied: Mar 15. Nitrochalk applied, seed drilled at 2½ bushels per acre: Mar 18. Trefoil and Italian ryegrass undersown: Apr 26. Harvested: Aug 5. Variety: Herta.

Early potatoes: Straw applied, fallow plots ploughed: Sept 29, 1954. Trefoil plots patched: Nov 3, 1954. Trefoil and ryegrass plots ploughed: Mar 14, 1955. Nitrochalk and basal fertilizers applied: Apr 6. Potatoes mechanically planted: Apr 12. Earthed up: June 28. Lifted: July 26. Variety: Ulster Chieftain.

Standard errors per plot:

Barley, Grain: 2.43 cwt per acre or 7.0% (20 d.f.)
Potatoes, Total tubers: 0.368 tons per acre or 13.1% (18 d.f.)

Estimates of produce (roots and tops) of green manure crops:
cwt per acre

| | Green manure | Ploughed in | Dry matter | Nitrogen |
|---------------------|--------------|-------------|------------|----------|
| <u>For Barley</u> | Trefoil | Early | 27.6 | 0.923 |
| | Ryegrass | Early | 29.6 | 0.519 |
| | Trefoil | Late | 15.6 | 0.514 |
| | Ryegrass | Late | 26.7 | 0.533 |
| <u>For Potatoes</u> | Trefoil | | 20.6 | 0.594 |
| | Ryegrass | | 33.9 | 0.505 |

55/Ba/1.4

| Barley, straw: cwt per acre | | | | | | | | | |
|---|-------------------|------|------------------------|-----------|-----------------------------------|------|------------------------|------|-------|
| | Green manures | | In barley for potatoes | | N: cwt per acre (including basal) | | Dung to cabbages 1952: | | Mean* |
| | Ploughed in Early | Late | None | Undersown | 0.23 | 0.46 | None | 10 | |
| Excluding plots fallow under old scheme | | | | | | | | | |
| Green manures after potatoes for barley | | | | | | | | | |
| Trefoil | 45.6 | 44.8 | 45.6 | 44.7 | 42.2 | 48.1 | 43.1 | 47.2 | 45.2 |
| Ryegrass | 34.7 | 35.0 | 35.1 | 34.6 | 27.1 | 42.6 | 32.5 | 37.2 | 34.9 |
| Green manures ploughed in | | | | | | | | | |
| Early | | | 41.0 | 39.2 | 36.2 | 44.1 | 38.0 | 42.3 | 40.1 |
| Late | | | 39.7 | 40.1 | 33.1 | 46.7 | 37.6 | 42.2 | 39.9 |
| Green manures in barley for potatoes | | | | | | | | | |
| None | | | | | 36.2 | 44.5 | 37.5 | 43.2 | 40.4 |
| Undersown | | | | | 33.1 | 46.2 | 38.1 | 41.2 | 39.7 |
| N: cwt per acre (including basal) | | | | | | | | | |
| 0.23 | | | | | | | 33.0 | 36.3 | 34.7 |
| 0.46 | | | | | | | 42.6 | 48.1 | 45.4 |
| Mean | | | | | | | 37.8 | 42.2 | 40.0 |
| Plots fallow under old scheme | | | | | | | | | |
| Green manures after potatoes for barley | | | | | | | | | |
| None | | | | | | | 14.8 | 23.1 | 19.0 |
| Fallow | | | | | | | 28.4 | 32.5 | 30.4 |
| Old scheme | 24.7 | 45.2 | 34.9 | Mean | | | 21.6 | 27.8 | 24.7 |

55/Be/1.1

LEY AND ARABLE ROTATIONS

Woburn Stackyard - 1955, the 18th year.

For details of rotations and treatments etc., see "Results of the Field Experiments" 1939-47, Vol. 1, section Bf/1 with the following exceptions:-

In 1949 and subsequently rye replaced wheat.

In 1954 and 1955 the seeds hay plots were split into two after the first crop, for testing 0.15 v. 0.30 cwt N per acre applied as nitrochalk.

In 1955 each of the 16 plots of test potatoes was split into four for the application (in addition to the basal fertilizer) of all combinations of

Nitrogen: None; 0.56 cwt N per acre as sulphate of ammonia.

Potash: None; 0.84 cwt K₂O per acre as muriate of potash.

Cultivations, etc.:

Treatment crops

Ley rotations

Ley 1st year. Ploughed twice: Sept 30, 1954 and Feb 2, 1955.

Basal fertilizers applied: Apr 29. Seed sown: Apr 30.

Nitrochalk applied: July 5. Grazed 4 circuits: June 28 - July 2, July 18-26, Sept 26-29, Oct 15-26. Seeds mixture (sown at 40 lb per acre): 24 lb S24 Perennial Ryegrass, 12 lb S143 Cocksfoot, 6 lb S123 Late Flowering Red Clover, 3 lb S100 White Clover.

Ley 2nd year. Nitrochalk applied: May 9, 1955 and July 5.

Grazed 7 circuits: Apr 30 - May 9, May 19-27, June 8-20, July 2-10, July 26 - Aug 3, Sept 29 - Oct 7, Oct 26 - Nov 3.

Ley 3rd year. Nitrochalk applied: May 11, 1955 and July 21.

Grazed 6 circuits: May 9-17, May 31 - June 8, June 20-28, July 10-18, Sept 17-25, Oct 10-18.

Lucerne 1st year. Ploughed twice: Sept 30, 1954 and Feb 2, 1955.

Basal fertilizers applied: Apr 29. Seed sown at 25 lb per acre: Apr 30. Cut twice: July 27 and Sept 16. Variety: Du Puits.

Lucerne 2nd year. Cut three times: June 13, July 27, Sept 16.

Lucerne 3rd year. Cut three times: June 13, July 27, Sept 16.

Arable rotations

Potatoes 1st Course. Ploughed twice: Sept 30, 1954 and Feb 2, 1955. Basal fertilizers applied, ridged, potatoes planted with dropper: Apr 19. Earthed up: June 28. Lifted: Sept 2. Variety: Majestic.

Rye 2nd Course. Ploughed: Oct 26, 1954. Seed drilled at 3½ bushels per acre: Dec 20. Nitrochalk applied: Apr 29, 1955. Seeds hay mixture undersown on 4 plots: May 9. Harvested: Aug 25. Variety: King II.

55/Be/1.2

Seed Hay 3rd Course. Seeds undersown in Rye: May 7, 1954.
Basal nitrochalk applied: Apr 19, 1955. 1st cut: June 13.
Nitrochalk applied: June 14. 2nd cut: Sept 16. Seeds
mixture per acre: 27 lb S24 Perennial Ryegrass, 12 lb Montgomery
Red clover, 3 lb Canadian Alsike Clover.
Sugar beet 3rd Course. Ploughed twice: Oct 1, 1954 and
Feb 1, 1955. Rubbed seed drilled at 8 lb per acre: Apr 16.
Basal nitrate of soda applied: Apr 18. Sprayed with systemic
insecticide, $\frac{1}{2}$ pint in 40 gallons per acre: June 10. Singled:
June 22. Lifted: Oct 28. Variety: Klein E.

Test crops

Potatoes 1st test crop. Ploughed twice: Nov 5, 1954 and Feb 4,
1955. Ridged, dung, basal and treatment fertilizers applied:
Apr 21. Potatoes hand planted: Apr 22. Earthed up: June 28.
Sprayed with copper fungicide, 5 lb per acre: Aug 19. Sprayed
with arsenious compound, 1 gallon in 40 gallons per acre: Sept 27.
Lifted: Oct 3. Variety: Majestic.
Barley 2nd test crop. Ploughed twice: Oct 28, 1954 and Feb 3, 1955.
Ground chalk applied: Mar 14. Nitrochalk applied, seed drilled
at $3\frac{1}{3}$ bushels per acre: Mar 17. Harvested: Aug 15. Variety:
Plumage Archer.

Standard errors per plot, Test Crops.

| | |
|-------------------------|--|
| Potatoes, Total tubers. | Whole plot: 0.828 tons per acre or 10.8% |
| | (4 d. f.) |
| | $\frac{1}{2}$ plot: 0.497 tons per acre or 6.5% |
| | (4 d. f.) |
| | $\frac{1}{8}$ plot: 1.162 tons per acre or 15.2% |
| | (24 d. f.) |
| Barley, Grain. | Whole plot: 3.25 cwt per acre or 10.6% |
| | (4 d. f.) |
| | $\frac{1}{2}$ plot: 1.97 cwt per acre or 6.5% |
| | (4 d. f.) |

Note. Potato root eelworm was found in this experiment in 1955. On
the treatment crop potatoes, several plots were very badly affected
and the results are unreliable. The eelworm was present on some
of the test crop potatoes but the yields were probably only slightly
affected.

55/Be/1.3

Summary of Results

Treatment crops

Ley, Sheep days of grazing per acre

| 1st year | 2nd year | 3rd year |
|----------|----------|----------|
| 533 | 1575 | 1358 |

Lucerne, yield of hay (at 85% dry matter): cwt per acre

| | 1st crop | 2nd crop | 3rd crop | Total |
|------------------------|----------|----------|----------|-------|
| <u>1st year</u> | | | | |
| No dung | 8.8 | 7.0 | | 15.8 |
| Dung in 1953 | 10.8 | 8.8 | | 19.6 |
| Increase | +2.0 | +1.8 | | +3.8 |
| Previous Rotation | | | | |
| Lucerne | 8.1 | 8.8 | | 16.9 |
| Arable with Sugar beet | 11.5 | 7.0 | | 18.5 |
| Mean | 9.8 | 7.9 | | 17.7 |
| <u>2nd year</u> | | | | |
| No dung | 33.1 | 30.4 | 14.2 | 77.7 |
| Dung in 1952 | 40.8 | 38.2 | 15.8 | 94.8 |
| Increase | +7.7 | +7.8 | +1.6 | +17.1 |
| Previous Rotation | | | | |
| Lucerne | 34.5 | 32.9 | 15.0 | 82.4 |
| Arable with Hay | 39.4 | 35.7 | 15.0 | 90.1 |
| Mean | 37.0 | 34.3 | 15.0 | 86.3 |
| <u>3rd year</u> | | | | |
| No dung | 27.1 | 29.3 | 11.0 | 67.4 |
| Dung in 1951 | 33.5 | 32.6 | 10.2 | 76.3 |
| Increase | +6.4 | +3.3 | -0.8 | +8.9 |
| Previous Rotation | | | | |
| Lucerne | 26.6 | 27.9 | 8.9 | 63.4 |
| Arable with Sugar beet | 34.0 | 34.0 | 12.3 | 80.3 |
| Mean | 30.3 | 31.0 | 10.6 | 71.9 |

55/Be/1.4

Treatment crops

| | Potatoes | | Rye | |
|---------------------------|--------------------------------|--|------------------------|--------|
| | Total tubers: tons per acre | Percentage ware ($1\frac{5}{8}$ " riddle) | Grain: cwt per acre | Straw: |
| No dung | 3.89 | 43.4 | 33.3 | 32.5 |
| Dung* | 4.62 | 49.2 | 34.4 | 35.9 |
| Increase | 0.73 | 5.8 | 1.1 | 3.4 |
| Previous Rotation | | | | |
| Ley | 7.72 | 76.5 | 36.2 | 35.6 |
| Lucerne | 5.86 | 62.9 | 32.4 | 33.6 |
| Arable with Hay | 1.92 | 23.4 | 33.7 | 33.3 |
| Arable with Sugar beet | 1.51 | 22.6 | 33.1 | 34.3 |
| Mean | 4.25 | 46.3 | 33.9 | 34.2 |

Hay
Yield (at 85% D.M.): cwt per acre

| | 1st crop | 2nd crop | Total | 2nd crop Resp. to N |
|-------------------|----------|----------|-------|------------------------|
| No dung | 50.8 | 5.3 | 56.1 | -1.2 |
| Dung in 1951 | 56.9 | 6.8 | 63.7 | +0.4 |
| Increase | +6.1 | +1.5 | +7.6 | +1.6 |
| Previous Rotation | | | | |
| Ley | 58.1 | 7.4 | 65.5 | -1.1 |
| Arable with Hay | 49.6 | 4.8 | 54.4 | +0.3 |
| Mean | 53.8 | 6.1 | 59.9 | -0.4 |

Sugar beet

| | Roots (washed): tons per acre | Sugar percentage | Total sugar: cwt per acre | Tops: tons per acre |
|---------------------------|----------------------------------|---------------------|------------------------------|------------------------|
| No dung | 9.20 | 18.1 | 33.4 | 6.70 |
| Dung in 1951 | 12.58 | 18.6 | 46.7 | 8.35 |
| Increase | 3.38 | 0.5 | 13.3 | 1.65 |
| Previous Rotation | | | | |
| Lucerne | 10.91 | 18.2 | 39.6 | 7.28 |
| Arable with Sugar beet | 10.86 | 18.5 | 40.4 | 7.78 |
| Mean | 10.89 | 18.3 | 40.0 | 7.53 |

*Dung applied: Potatoes:- for test crop potatoes in 1953.
Rye:- for test crop potatoes in 1952.

55/Be/1.5

| | Test Crop | | | | Mean |
|--|-------------------|---------|-----------------|------------------------|------|
| | Previous Rotation | | | | |
| | Ley | Lucerne | Arable with hay | Arable with sugar beet | |

Potatoes, Total tubers: tons per acre

| | | | | | |
|---|-------|-------|-------|-------|--------------------------|
| Mean (± 0.586) | 8.86 | 8.46 | 7.20 | 6.09 | 7.65 |
| No dung (± 0.636) | 8.09 | 7.15 | 6.42 | 4.92 | 6.64 |
| Dung | 9.63 | 9.78 | 7.97 | 7.25 | 8.66 |
| Response to 15 tons dung per acre (± 0.497) | +1.54 | +2.63 | +1.55 | +2.33 | +2.02 (± 0.249) |
| Response to additional 0.56 cwt N per acre | | | | | (± 0.411) |
| No dung (± 0.822) | -1.89 | +1.69 | +1.52 | +0.53 | +0.47 |
| Dung | +1.41 | +0.69 | +0.88 | +0.25 | +0.81 |
| Response to additional 0.84 cwt K ₂ O per acre | | | | | (± 0.411) |
| No dung (± 0.822) | +0.15 | +1.11 | +0.38 | +1.44 | +0.78 |
| Dung | -0.61 | +2.41 | +0.06 | -0.02 | +0.46 |

Potatoes, Percentage ware ($1\frac{5}{8}$ " riddle)

| | | | | | |
|---|------|------|------|------|------|
| Mean | 87.7 | 87.4 | 86.0 | 80.2 | 85.3 |
| No dung | 88.0 | 86.0 | 85.6 | 80.5 | 85.0 |
| Dung | 87.5 | 88.7 | 86.5 | 80.0 | 85.7 |
| Response to 15 tons dung per acre | -0.5 | +2.7 | +0.9 | -0.5 | +0.7 |
| Response to additional 0.56 cwt N per acre | | | | | |
| No dung | +1.0 | -1.1 | +1.5 | +1.5 | +0.7 |
| Dung | +0.4 | +2.5 | +1.1 | +1.1 | +1.3 |
| Response to additional 0.84 cwt K ₂ O per acre | | | | | |
| No dung | -1.1 | +4.6 | +4.0 | +1.9 | +2.4 |
| Dung | -0.7 | -0.6 | -0.8 | +2.1 | 0.0 |

55/Be/1.6

Test Crop

Plots receiving no additional N or K

| | Previous Rotation | | | | Mean |
|--|-------------------|---------|-----------------|------------------------|-------|
| | Ley | Lucerne | Arable with hay | Arable with sugar beet | |
| Potatoes, Total tubers: tons per acre | | | | | |
| Mean (± 0.675) | 9.33 | 6.54 | 6.57 | 5.80 | 7.06 |
| No dung (± 0.955)* | 9.35 | 5.54 | 5.27 | 4.28 | 6.11 |
| Dung | 9.32 | 7.54 | 7.87 | 7.32 | 8.01 |
| Response to 15 tons dung per acre (± 1.122) | -0.03 | +2.00 | +2.60 | +3.04 | +1.90 |
| Potatoes, Percentage ware ($\frac{5}{16}$ " riddle) | | | | | |
| Mean | 88.0 | 86.1 | 84.8 | 78.4 | 84.3 |
| No dung | 88.0 | 84.0 | 82.8 | 78.5 | 83.3 |
| Dung | 88.0 | 88.2 | 86.6 | 78.4 | 85.3 |
| Response to 15 tons dung per acre | 0.0 | +4.2 | +3.8 | -0.1 | +2.0 |

*For use in comparisons other than vertical.

55/Be/1.7

| | Test Crop | | | | Mean |
|------------------------------|-----------|-------------------|-----------------|------------------------|-----------------------|
| | Ley | Previous Rotation | | Arable with sugar beet | |
| | | Lucerne | Arable with hay | | |
| Barley, Grain: cwt per acre | | | | | |
| No dung | 32.2 | 31.0 | 25.7 | 22.3 | 27.8 |
| Dung in 1954 (± 2.50)* | 37.5 | 37.2 | 30.8 | 27.6 | 33.3 |
| Mean (± 2.30) | 34.9 | 34.1 | 28.3 | 24.9 | 30.5 |
| Increase (± 1.97) | 5.3 | 6.2 | 5.1 | 5.3 | 5.5 (± 0.99) |
| Barley, Straw: cwt per acre | | | | | |
| No dung | 34.2 | 29.6 | 25.7 | 22.3 | 27.9 |
| Dung in 1954 | 38.0 | 38.2 | 33.3 | 28.5 | 34.5 |
| Mean | 36.1 | 33.9 | 29.5 | 25.4 | 31.2 |
| Increase | 3.8 | 8.6 | 7.6 | 6.2 | 6.6 |

*For use in comparisons other than vertical.

55/Bf/1.1

WOBURN MARKET GARDEN EXPERIMENT

Organic Manures and Nitrogen, Lansome 1955 the 14th year

The present cropping comprises two series, each carrying in turn the crops of a two course rotation: 1st year - Globe beet followed by Spring cabbages; 2nd year - Leeks.

Note: The results for the 1955-56 leeks will be included in the 1956 report.

Design (each series): 4 randomized blocks of 10 plots each, certain interactions being confounded with block differences.

Area of each plot: 0.0125 acre.

Treatments applied to each crop.

Organic manures: Dung; Sewage sludge compost; Sewage sludge (West Middlesex); Vegetable compost, each at 10 and 20 tons per acre. N (applied as 'Nitro-Chalk'): None; 0.3 cwt per acre on plots receiving organic manures. None; 0.3; 0.6; 0.9 cwt per acre on plots not receiving organic manure. The last two rates are applied in two equal dressings.

Basal dressing per acre to each crop: 0.3 cwt P_2O_5 ; 0.3 cwt K_2O applied as granular fertilizer (13% P_2O_5 , 13% K_2O).

Cultivations, etc.:

Spring cabbages 1954-55.

Organic manures spread and ploughed in: Sept 25, 1954. Ground chalk at 20 cwt per acre, Aldrin at $1\frac{1}{2}$ cwt per acre and basal fertilizer applied: Sept 27. Cabbages planted and watered in: Sept 29. First dressing of 'Nitro-Chalk' applied: Mar 14, 1955. Second dressing of 'Nitro-Chalk' applied: Apr 13. Cut 8 times: June 3 - July 8. Variety: Durham Early.

Note: The cabbages were attacked by pigeons in winter. The plots without organic manures or 'Nitro-Chalk' were particularly badly damaged.

Globe beet 1955.

Organic manures applied and ploughed in: Apr 29. Ground chalk applied at 20 cwt per acre: May 3. Basal fertilizer and first dressing of 'Nitro-Chalk' applied: May 11. Seed drilled at 14 lb per acre: May 16. Singled: July 1-8. Second dressing of 'Nitro-Chalk' applied: July 15. Harvested: Aug 18 - Sept 8. Variety: Detroit.

Standard errors per plot.

Spring cabbages 1954-55, weight of headed: 1.21 tons per acre or 23.3% (16 d.f.)*

Globe beet 1955, saleable bulbs: 1.30 tons per acre or 23.4% (17 d.f.)

*1 missing value.

55/Bf/1.2

Summary of Results

Spring cabbages 1954-55

| Organic manures | Level of manuring: tons per acre | N: cwt per acre | | | Mean | |
|---------------------------------|-------------------------------------|---------------------|-------------------|------|----------|-------|
| | | None | 0.3 | 0.6 | | 0.9 |
| Weight of headed: tons per acre | | | | | | |
| | | (±0.853) | | | (±0.603) | |
| None | | 0.22 ⁽¹⁾ | 2.80 | 5.01 | 4.03 | 1.51* |
| Dung | 10 | 2.06 ⁽²⁾ | 5.49 | | | 3.77 |
| | 20 | 5.60 | 8.05 | | | 6.82 |
| Sludge compost | 10 | 3.01 | 5.84 | | | 4.43 |
| | 20 | 4.45 | 8.56 | | | 6.51 |
| Sludge | 10 | 4.65 | 5.57 | | | 5.11 |
| | 20 | 5.84 | 10.45 | | | 8.14 |
| Vegetable compost | 10 | 2.86 | 5.55 | | | 4.20 |
| | 20 | 4.30 | 9.18 | | | 6.74 |
| Mean (±0.301) | | 4.10 ⁺ | 7.34 ⁺ | | | 5.18 |
| Total produce: tons per acre | | | | | | |
| None | | 3.01 ⁽¹⁾ | 5.44 | 7.55 | 6.43 | 4.22* |
| Dung | 10 | 6.05 ⁽²⁾ | 7.78 | | | 6.92 |
| | 20 | 8.44 | 10.09 | | | 9.27 |
| Sludge compost | 10 | 7.22 | 8.15 | | | 7.68 |
| | 20 | 8.26 | 10.50 | | | 9.38 |
| Sludge | 10 | 8.05 | 8.83 | | | 8.44 |
| | 20 | 10.50 | 12.07 | | | 11.29 |
| Vegetable compost | 10 | 5.95 | 8.14 | | | 7.04 |
| | 20 | 7.48 | 10.73 | | | 9.10 |
| Mean | | 7.74 ⁺ | 9.54 ⁺ | | | 8.03 |
| Percentage headed, (by number) | | | | | | |
| None | | 0.2 ⁽¹⁾ | 40.4 | 58.8 | 53.0 | 20.3* |
| Dung | 10 | 27.0 ⁽²⁾ | 64.4 | | | 45.7 |
| | 20 | 58.4 | 75.4 | | | 66.9 |
| Sludge compost | 10 | 36.5 | 69.3 | | | 52.9 |
| | 20 | 50.9 | 78.2 | | | 64.6 |
| Sludge | 10 | 52.4 | 57.0 | | | 54.7 |
| | 20 | 52.3 | 84.9 | | | 68.6 |
| Vegetable compost | 10 | 34.5 | 60.6 | | | 47.6 |
| | 20 | 49.8 | 84.5 | | | 67.1 |
| Mean | | 45.2 ⁺ | 71.8 ⁺ | | | 54.4 |

(1) Both plots receiving no organics or N, were badly damaged by birds.

(2) Includes one estimated value.

* Mean over None and 0.3 cwt N per acre only.

+ Excluding 'No organics'.

55/Bf/1.3

Globe beet 1955

| Organic manures | Level of manuring: tons per acre | N: cwt per acre | | | | Mean |
|----------------------------------|-------------------------------------|-------------------|--------------------|------|------|----------|
| | | None | 0.3 | 0.6 | 0.9 | |
| Saleable bulbs: tons per acre | | | | | | |
| | | (±0.917) | | | | (±0.648) |
| None | | 1.37 | 1.58 | 2.72 | 3.13 | 1.47* |
| Dung | 10 | 6.34 | 4.38 | | | 5.36 |
| | 20 | 7.68 | 8.07 | | | 7.87 |
| Sludge compost | 10 | 4.82 | 6.54 | | | 5.68 |
| | 20 | 8.19 | 8.25 | | | 8.22 |
| Sludge | 10 | 4.31 | 5.55 | | | 4.93 |
| | 20 | 6.79 | 7.01 | | | 6.90 |
| Vegetable compost | 10 | 5.61 | 6.64 | | | 6.12 |
| | 20 | 6.22 | 5.81 | | | 6.01 |
| Mean (±0.324) | | 6.24 ⁺ | 6.53 ⁺ | | | 5.55 |
| Total produce: tons per acre | | | | | | |
| None | | 2.78 | 2.76 | 4.74 | 5.54 | 2.77* |
| Dung | 10 | 9.37 | 7.38 | | | 8.37 |
| | 20 | 11.21 | 11.68 | | | 11.45 |
| Sludge compost | 10 | 7.72 | 10.09 | | | 8.91 |
| | 20 | 11.80 | 12.01 | | | 11.91 |
| Sludge | 10 | 7.07 | 9.08 | | | 8.07 |
| | 20 | 10.41 | 11.44 | | | 10.92 |
| Vegetable compost | 10 | 8.44 | 9.90 | | | 9.17 |
| | 20 | 9.47 | 9.50 | | | 9.48 |
| Mean | | 9.44 ⁺ | 10.14 ⁺ | | | 8.62 |
| Plant number: thousands per acre | | | | | | |
| None | | 50.0 | 40.7 | 59.1 | 59.7 | 45.4* |
| Dung | 10 | 79.1 | 76.9 | | | 78.0 |
| | 20 | 83.0 | 75.3 | | | 79.2 |
| Sludge compost | 10 | 75.5 | 76.7 | | | 76.1 |
| | 20 | 82.9 | 80.9 | | | 81.9 |
| Sludge | 10 | 75.5 | 75.3 | | | 75.4 |
| | 20 | 82.1 | 85.3 | | | 83.7 |
| Vegetable compost | 10 | 81.8 | 77.5 | | | 79.6 |
| | 20 | 80.5 | 84.2 | | | 82.4 |
| Mean | | 80.0 ⁺ | 79.0 ⁺ | | | 74.1 |

*Mean over None and 0.3 cwt N per acre only.

⁺Excluding 'No organics'.

IRRIGATION EXPERIMENT

The 5th year

The effects of irrigation and nitrogen - Woburn Butt Close 1955.

For details of cropping, treatments etc. see "Results of the Field Experiments" 1954, Section 54/Bg/1. The 4 irrigation treatments (O A B C) are not necessarily applied to a particular crop in any one year. For particulars of the irrigations and their designations the annual reports should be consulted.

Area of each sub-plot: Cut grass, 0.0264, remainder, 0.0278 acre.

Area harvested: Cut grass, 0.0165, potatoes, 0.0155, sugar beet, 0.0176, barley, 0.0168 acre.

Rainfall and Irrigation: inches

| Week ending | Rain-fall | Irrigation | | | | | | | | | |
|-------------|-----------|------------|------|------|------------|------|------|---------|-----------|------|------|
| | | Potatoes | | | Sugar beet | | | Bar-ley | Cut Grass | | |
| | | A | B | C | A | B | C | B & C | A | B | C |
| May 2 | 0.36 | | | | | | | | | .75 | .75 |
| 9 | 0.28 | | | | | | | .50 | | .06 | .33 |
| 16 | 0.81 | | | | | | | | | | |
| 23 | 1.94 | | | | | | | | | | |
| 30 | 1.34 | | | | | | | | | | |
| June 6 | 0.30 | | | | | | | | | | |
| 13 | 1.50 | | | | | | | | | | |
| 20 | 0.25 | | | | | | | | | | |
| 27 | 0.07 | | | | | | | | | | |
| July 4 | 0.21 | .50 | | .50 | .38 | | .50 | .59 | | | .56 |
| 11 | - | .60 | | .60 | .62 | | .50 | .30 | | | .80 |
| 18 | - | 1.34 | | 1.34 | 1.34 | | 1.34 | | 1.00 | | 1.08 |
| 25 | - | | .57 | .57 | | .50 | .50 | | .50 | | .82 |
| Aug 1 | - | | .50 | .50 | | .79 | .79 | | .50 | 1.00 | .50 |
| 8 | 0.02 | | .75 | .75 | | .62 | .62 | | .50 | .57 | .84 |
| 15 | 0.33 | | .50 | .50 | | .19 | .19 | | .20 | | .67 |
| 22 | 0.30 | | | | | .35 | .35 | | .33 | .50 | .56 |
| 29 | 0.03 | | .50 | .50 | | .50 | .50 | | .75 | | |
| Sept 5 | - | | .54 | .54 | | .63 | .63 | | .75 | .50 | .50 |
| 12 | 0.49 | | .50 | .50 | | .50 | .50 | | .33 | .33 | .33 |
| 19 | 0.11 | | | | | | | | | | |
| 26 | 0.95 | | | | | | | | | | |
| Oct 2 | 0.01 | | | | | | | | | | |
| 9 | 0.41 | | | | | | | | | | |
| 16 | - | | | | | | | | | | |
| 23 | 1.14 | | | | | | | | | | |
| 30 | 0.23 | | | | | | | | | | |
| Nov 6 | 0.69 | | | | | | | | | | |
| Total | 11.77 | 2.44 | 3.86 | 6.30 | 2.34 | 4.08 | 6.42 | 1.39 | 4.86 | 3.71 | 7.74 |

Note: On barley O = A
B = C.

55/Bg/1.2

Cultivations, etc.:

Potatoes. Ploughed: Sept 9, 1954. FYM applied: Jan 17, 1955.
 Ploughed: Jan 24. Fertilizers applied: Apr 18. Potatoes
 planted by machine: Apr 19. Earthed up: June 22. Sprayed
 with arsenious compound, 1 gallon in 40 gallons per acre:
 Sept 27. Lifted: Oct 7. Variety: Majestic.

Sugar beet. Ploughed: Oct 29, 1954. Ground chalk applied at
 20 cwt per acre: Mar 15, 1955. Salt applied: Apr 14.
 Fertilizers applied: Apr 15. Seed drilled at 5.7 lb per acre
 (rubbed and graded): Apr 18. Sprayed with Parathion, $\frac{1}{2}$ pint
 in 40 gallons per acre: June 10. Singled: June 20. Lifted:
 Nov 1. Variety: Klein E.

Barley. Ploughed: Dec 22, 1954. Fertilizers applied, seed
 drilled at $2\frac{1}{2}$ bushels per acre: Mar 18. Sprayed with MCPA,
 low volume, at $2\frac{1}{2}$ pints per acre: May 22. Harvested: Aug 4.
 Variety: Herta.

Cut grass. Nitrochalk applied: Apr 13. Basal fertilizer applied:
 Apr 20. Cut: May 12, June 2, June 24, July 28 (A and C plots
 only), Aug 26, Sept 28 (A, B and C plots only), Nov 6.
 Nitrochalk applied after each cut except the last. Variety:
 Cocksfoot S37.

Standard errors per plot:

| | | | |
|-------------|---------------------------|------------------------------|-----------|
| Potatoes, | Total tubers, whole plot: | 1.959 tons per acre or 12.0% | (6 d.f.) |
| | sub plot: | 0.703 tons per acre or 4.3% | (8 d.f.) |
| Sugar beet, | Total sugar, whole plot: | 4.80 cwt per acre or 11.6% | (6 d.f.) |
| | sub plot: | 3.82 cwt per acre or 9.3% | (8 d.f.) |
| | Tops, whole plot: | 1.229 tons per acre or 11.6% | (6 d.f.) |
| | sub plot: | 0.782 tons per acre or 7.4% | (8 d.f.) |
| Barley, | Grain, whole plot: | 1.99 cwt per acre or 5.5% | (8 d.f.) |
| | sub plot: | 1.89 cwt per acre or 5.2% | (10 d.f.) |
| Cut grass, | Dry matter, whole plot: | 4.05 cwt per acre or 6.8% | (6 d.f.) |
| | sub plot: | 2.76 cwt per acre or 4.6% | (8 d.f.) |

55/Bg/1.3

Summary of Results

| N: cwt per acre including basal | Irrigation | | | | Mean |
|---|------------|-------|-------|-------|--------------------------|
| | 0 | A | B | C | |
| Potatoes, total tubers: tons per acre (± 1.167) [‡] | | | | | |
| 0.5 | 10.83 | 15.57 | 16.73 | 19.03 | 15.54 |
| 1.0 | 11.40 | 16.60 | 18.36 | 21.66 | 17.00 |
| Mean (± 1.131) | 11.11 | 16.08 | 17.55 | 20.35 | 16.27 |
| Difference (± 0.574) | +0.57 | +1.03 | +1.63 | +2.63 | +1.46 (± 0.287) |
| Potatoes, percentage ware ($1\frac{3}{8}$ " riddle) | | | | | |
| 0.5 | 80.2 | 87.9 | 92.1 | 91.2 | 87.8 |
| 1.0 | 80.9 | 89.4 | 90.4 | 92.8 | 88.4 |
| Mean | 80.6 | 88.6 | 91.2 | 92.0 | 88.1 |
| Difference | +0.7 | +1.5 | -1.7 | +1.6 | +0.6 |
| Sugar beet, roots (washed): tons per acre | | | | | |
| 0.4 | 9.83 | 10.06 | 12.40 | 13.33 | 11.40 |
| 0.8 | 10.53 | 12.91 | 14.08 | 15.23 | 13.19 |
| Mean | 10.18 | 11.48 | 13.24 | 14.28 | 12.30 |
| Difference | 0.70 | 2.85 | 1.68 | 1.90 | 1.79 |
| Sugar beet, sugar percentage | | | | | |
| 0.4 | 16.7 | 16.9 | 16.5 | 16.5 | 16.6 |
| 0.8 | 16.9 | 17.4 | 16.5 | 16.8 | 16.9 |
| Mean | 16.8 | 17.1 | 16.5 | 16.6 | 16.8 |
| Difference | 0.2 | 0.5 | 0.0 | 0.3 | 0.3 |
| Sugar beet, total sugar: cwt per acre (± 3.18) [‡] | | | | | |
| 0.4 | 32.8 | 34.1 | 40.8 | 43.9 | 37.9 |
| 0.8 | 35.6 | 45.2 | 46.5 | 51.6 | 44.7 |
| Mean (± 2.77) | 34.2 | 39.6 | 43.6 | 47.7 | 41.3 |
| Difference (± 3.12) | 2.8 | 11.1 | 5.7 | 7.7 | 6.8 (± 1.56) |
| Sugar beet, tops: tons per acre (± 0.778) [‡] | | | | | |
| 0.4 | 7.55 | 8.38 | 8.55 | 11.00 | 8.87 |
| 0.8 | 9.63 | 11.95 | 12.56 | 15.33 | 12.37 |
| Mean (± 0.710) | 8.59 | 10.17 | 10.56 | 13.17 | 10.62 |
| Difference (± 0.638) | 2.08 | 3.57 | 4.01 | 4.33 | 3.50 (± 0.319) |

[‡]for use in comparisons other than vertical.

55/Bg/1.4

| N: cwt per acre including basal | Irrigation | | | Mean |
|------------------------------------|-----------------------------|-------|--|-----------------------|
| | 0 & A | B & C | | |
| Barley, grain: cwt per acre | | | | |
| | (± 0.98) [‡] | | | |
| 0.2 | 31.7 | 32.5 | | 32.1 |
| 0.4 | 40.5 | 39.4 | | 39.9 |
| Mean (± 0.81) | 36.1 | 36.0 | | 36.0 |
| Difference (± 1.09) | 8.8 | 6.9 | | 7.8 (± 0.77) |

| | | | | |
|-----------------------------|------|------|--|------|
| Barley, straw: cwt per acre | | | | |
| 0.2 | 30.6 | 32.3 | | 31.4 |
| 0.4 | 44.2 | 46.9 | | 45.6 |
| Mean | 37.4 | 39.6 | | 38.5 |
| Difference | 13.6 | 14.6 | | 14.2 |

| No. of cuts | Irrigation | | | | Mean |
|-------------------------------------|------------|------|------|------|------------------------|
| | 0 | A | B | C | |
| 5 | | 7 | 6 | 7 | |
| Cut grass, Dry Matter: cwt per acre | | | | | |
| (± 2.59) [‡] | | | | | |
| Level of N | | | | | |
| N ₁ | 37.1 | 52.1 | 44.9 | 63.3 | 49.4 |
| N ₂ | 54.5 | 78.4 | 66.8 | 81.5 | 70.3 |
| Mean (± 2.34) | 45.8 | 65.3 | 55.8 | 72.4 | 59.8 |
| Difference (± 2.25) | 17.4 | 26.3 | 21.9 | 18.2 | 20.9 (± 1.13) |

[‡]for use in comparisons other than vertical.

Cut Grass

N₁ = 0.15 cwt N per acre after each cut except the last
 N₂ = 0.30 cwt N per acre after each cut except the last.

55/Ca/1.1

WINTER WHEAT

The effects of Crop sequences, Varieties, Seed rates and Nitrogen on the incidence of Eyespot (Cercospora herpotrichoides) - Long Hoos I, II and III, 1955. The 2nd preliminary year.

For details of treatments and crop sequences etc., see "Results of the Field Experiments 1954", Section 54/Ca/2.1 with the exception that in 1955 the seed rates were changed to

Holdfast: $1\frac{1}{2}$, 3 bushels per acre

Cappelle: 2, 4 bushels per acre,

in order to give approximately the same number of seeds of each variety per acre.

Area of each plot: 0.0212 acre. Area harvested: 0.0140 acre.

Cultivations, etc.: Ploughed: Oct 8, 1954. Seed combine drilled: Oct 15. 1st application of nitrochalk: Mar 15, 1955. 2nd application: May 9. Combine harvested: Aug 19.

Standard error per plot:

Grain (at 85% D.M.): 1.92 cwt per acre or 5.8% (12 d.f.)

Records of incidence of disease (Eyespot and Take-All), estimates of % area lodged, and counts of plant, shoot and straw numbers were made.

55/Ca/1.2

Summary of Results

| Response to | Responses to treatments | | | | | | |
|-------------|-------------------------|---------------|---------------|-----------|--------|-----------|------|
| | Mean | Variety | | Seed rate | | Nitrogen: | |
| | | Hold- fast | Capp- elle | Single | Double | 0.46 | 0.93 |

Grain (at 85% Dry Matter)

| <u>Previous crop wheat:</u> | Mean yield 17.1 cwt per acre | | | | | | |
|----------------------------------|------------------------------|------|------|------|------|------|---------|
| | (±0.96) | | | | | | (±1.36) |
| Variety (Cappelle - Holdfast) | +6.5 | - | - | +7.1 | +5.9 | +4.8 | +8.2 |
| Seed rate (Double - Single) | -0.8 | -0.2 | -1.4 | - | - | -1.6 | 0.0 |
| Nitrogen (0.93 - 0.46) | +3.6 | +1.9 | +5.3 | +2.8 | +4.4 | - | - |

| <u>Previous crop potatoes:</u> | Mean yield 49.3 cwt per acre | | | | | | |
|----------------------------------|------------------------------|------|------|-------|-------|------|---------|
| | (±0.96) | | | | | | (±1.36) |
| Variety (Cappelle - Holdfast) | +12.9 | - | - | +12.8 | +13.0 | +9.9 | +15.9 |
| Seed rate (Double - Single) | +3.3 | +3.2 | +3.4 | - | - | +2.8 | +3.8 |
| Nitrogen (0.93 - 0.46) | +5.6 | +2.6 | +8.6 | +5.1 | +6.1 | - | - |

Straw

| <u>Previous crop wheat:</u> | Mean yield 24.3 cwt per acre | | | | | | |
|----------------------------------|------------------------------|-------|------|------|------|------|------|
| Variety (Cappelle - Holdfast) | +4.3 | - | - | +4.1 | +4.5 | +8.3 | +0.3 |
| Seed rate (Double - Single) | +0.5 | +0.3 | +0.7 | - | - | +0.7 | +0.3 |
| Nitrogen (0.93 - 0.46) | +8.1 | +12.1 | +4.1 | +8.3 | +7.9 | - | - |

| <u>Previous crop potatoes:</u> | Mean yield 42.0 cwt per acre | | | | | | |
|----------------------------------|------------------------------|------|------|------|------|------|------|
| Variety (Cappelle - Holdfast) | -2.5 | - | - | -4.3 | -0.7 | -3.2 | -1.8 |
| Seed rate (Double - Single) | +5.3 | +3.5 | +7.1 | - | - | +2.8 | +7.8 |
| Nitrogen (0.93 - 0.46) | +4.2 | +3.5 | +4.9 | +1.7 | +6.7 | - | - |

General means. Grain: 33.2 cwt per acre
 Straw: 33.2 cwt per acre
 Mean dry matter % as harvested, Grain: 81.8

WINTER WHEAT

Control of wheat bulb fly by insecticides - Pennell's Piece 1955.

Design: 4 randomized blocks of 6 plots each.

Area of each plot: 0.00643 acre.

Treatments: Insecticides:-

- None (2 plots per block). (0)
- Seed dressed with Technical Dieldrin (97%) at 2.25% of seed w/w using cellulose ether sticker. (1)
- 4% Dieldrin dust at 1 cwt per acre combine drilled with seed. (2)
- Sprayed early with Parathion 0.05% v/v at 100 gallons per acre. (3)
- Sprayed late with Parathion 0.05% v/v at 100 gallons per acre. (4)

Basal dressing per acre: 3 cwt nitrochalk, 20 cwt hydrated lime.

Note: All seed treated with organo-mercurial fungicide.

Cultivations, etc.: Ploughed (for bare fallow): Nov 2, 1953. Lime applied: Mar 3, 1954. Ploughed: May 17. Seed drilled at 2 bushels per acre: Dec 21. "3" plots sprayed with Parathion: Feb 11, 1955. "4" plots sprayed with Parathion: Apr 7. Nitrochalk applied: May 9. Combine harvested: Aug 24. Variety: Cappelle. Previous crop: Bare fallow.

Standard error per plot:

Grain (at 85% dry matter): 3.75 cwt per acre or 8.4% (16 d.f.)

Counts of numbers of plants, tillers, damaged tillers and of wheat bulb fly larvae were made.

Summary of Results

Grain (at 85% dry matter): cwt per acre

| | Insecticide | | | | | Mean |
|-------------------------|---------------------|------|-------|------|------|------|
| | 0 | 1 | 2 | 3 | 4 | |
| Mean (± 1.88) | 41.0 ⁽¹⁾ | 47.2 | 53.7 | 47.9 | 38.7 | 44.9 |
| Increase (± 2.30) | | +6.2 | +12.7 | +6.9 | -2.3 | |

(1) ± 1.33

Mean dry matter % as harvested: 84.4

55/Ca/3

WHEAT

Seed rates in relation to control of wheat bulb fly - Long Hoos 7.
Preliminary years 1954 (spring wheat) and 1955 (winter wheat).

Design: 4 4 x 4 squares with treatments on rows in 1954 and on columns in 1955.

Area of each plot: 0.063 acre. Area harvested: 1954 - 0.0226 acre.
1955 - 0.0262 acre.

Treatments: 1954 and 1955.

Bare fallow.

Seed rates: $\frac{1}{3}$; 1; 3 bushels per acre.

The 1954 treatments were ignored in 1955, so that there were, in effect, 4 randomized blocks of 3 plots in each year.

Basal dressings per acre.

1954. $2\frac{1}{2}$ cwt sulphate of ammonia.

1955. $1\frac{1}{2}$ cwt compound granular fertilizer (12% N, 12% P₂O₅, 15% K₂O)
combine drilled; 3 cwt sulphate of ammonia.

Cultivations, etc.:

1954. Ploughed: Dec 29, 1953. Sulphate of ammonia applied:
Mar 15, 1954. Seed sown: Mar 18. Combine harvested: Sept 22.
Variety: Koga II. Previous crop: Sugar beet.

1955. Ploughed: Oct 14, 1954. Seed (dressed with organo-mercurial compound only), combine drilled: Dec 22. Sulphate of ammonia applied: May 13, 1955. Combine harvested: Aug 25. Variety: Cappelle.

Standard errors per plot: Grain (at 85% D.M.).

1954: 2.12 cwt per acre or 6.3% (6 d.f.)

1955: 5.66 cwt per acre or 16.3% (6 d.f.)

Records were made of the following:

1954. Plant number.

1955. Number of wheat bulb fly larvae, weight per ear and no. of grains per ear.

Summary of Results

Grain (at 85% dry matter): cwt per acre

| | Seed rate: bushels per acre | | | Mean |
|-----------------------------------|-----------------------------|------|------|------|
| | $\frac{1}{3}$ | 1 | 3 | |
| 1954. Spring wheat (± 1.06) | 29.9 | 33.7 | 37.5 | 33.7 |
| 1955. Winter wheat (± 2.83) | 26.1 | 40.3 | 37.4 | 34.6 |

Mean dry matter % as harvested, 1954: 79.4
1955: 84.8

55/Ca/4.1

WINTER WHEAT

Varieties, seed rates, levels and times of application of N: - Woburn, Roadpiece 1955, the 2nd year.

Design: 4 randomized blocks of 8 plots each, certain high order interactions being confounded with block differences. In addition each block contained 2 plots with no nitrogen, the variety x seed rate interaction being confounded.

Area of each plot: 0.0159 acre. Area harvested: 0.0140 acre.

Treatments: All combinations of:-

Varieties: Holdfast; Cappelle.

Seed rates: Holdfast, $1\frac{1}{2}$; 3 bushels per acre.

Cappelle, 2; 4 bushels per acre.

Nitrogen: 0.5; 1.0 cwt N per acre as nitrochalk.

Time of application of N: half dressing in March and again in May; whole dressing mid March; mid April; mid May.

Basal dressing: 1 cwt per acre compound granular fertilizer (12% N, 12% P_2O_5 , 15% K_2O) combine drilled with seed.

Cultivations, etc.: Ploughed: Sept 28, 1954. Combine drilled: Oct 25. March top dressing applied: Mar 15, 1955. April top dressing applied: Apr 20. All plots sprayed with D.N.O.C. at $1\frac{1}{2}$ gallons in 80 gallons: May 19. May top dressing applied: May 24. Combine harvested: Aug 22. Varieties: Holdfast and Cappelle. Previous crop: Wheat.

Note (1) The experiment is a repetition on the same plots of the one carried out in 1954 (see "Results of the Field Experiments 1954", Section 54/Ca/7). There were minor changes in the treatments but the same randomization was used.

(2) Records of incidence of disease (Take-all and Eyespot) and weeds, and counts of plant, shoot and ear numbers were made.

Standard error per plot.

Grain: 3.95 cwt per acre or 26.9% (12 d.f.)

Summary of Results

Grain: cwt per acre

| | T ₁ | T ₂ | T ₃ | T ₄ | Mean |
|---------------------------|----------------|----------------|----------------|----------------|---------------------|
| Mean (± 1.40) | 16.5 | 16.9 | 18.9 | 12.2 | 16.1 |
| (±1.97) | | | | | (±0.99) |
| V ₁ | 17.0 | 15.6 | 18.3 | 12.4 | 15.8 |
| V ₂ | 16.0 | 18.3 | 19.6 | 12.0 | 16.5 |
| Difference (± 2.79) | -1.0 | +2.7 | +1.3 | -0.4 | +0.7 (± 1.40) |
| R ₁ | 14.1 | 16.3 | 16.8 | 9.6 | 14.2 |
| R ₂ | 18.9 | 17.6 | 21.1 | 14.8 | 18.1 |
| Difference (± 2.79) | +4.8 | +1.3 | +4.3 | +5.2 | +3.9 (± 1.40) |
| N ₁ | 14.9 | 11.9 | 15.2 | 12.2 | 13.6 |
| N ₂ | 18.1 | 22.0 | 22.6 | 12.1 | 18.7 |
| Difference (± 2.79) | +3.2 | +10.1 | +7.4 | -0.1 | +5.1 (± 1.40) |

| | R ₁ | R ₂ | N ₀ | N ₁ | N ₂ | Mean |
|---------------------|----------------|----------------|----------------|-----------------|----------------|------|
| Mean (± 0.99) | | | (±1.40) 8.8 | (±0.99) 13.6 | 18.7 | 14.7 |
| (±1.40) | | | (±1.97) | (±1.40) | (±0.88) | |
| V ₁ | 13.7 | 18.0 | 8.8 | 13.6 | 18.0 | 14.4 |
| V ₂ | 14.7 | 18.2 | 8.8 | 13.5 | 19.4 | 14.9 |
| R ₁ | | | 8.9 | 11.9 | 16.5 | 13.1 |
| R ₂ | | | 8.6 | 15.3 | 20.9 | 16.2 |

Mean dry matter % as harvested: 85.5

Treatments

V₁ Holdfast R₁, R₂ 1½, 3 bushels per acre N₀ No N
 V₂ Cappelle R₁, R₂ 2, 4 bushels per acre N₁ 0.46 cwt N per acre
 N₂ 0.93 cwt N per acre

T₁ Nitrochalk half in March half in May T₃ Nitrochalk all in mid April
 T₂ Nitrochalk all in mid March T₄ Nitrochalk all in mid May

The V × R table does not include the plots receiving no nitrogen.

55/Ca/5

SPRING WHEAT

Residual effects of Dung, Nitrogen, Phosphate and Potash - Sawyers I 1955.

Design: 4 randomized blocks of 8 plots each, the interaction DNEK being confounded with block differences.

Area of each plot: 0.0210 acre. Area harvested: 0.0150 acre.

Treatments, applied to potatoes in 1954: All combinations of:-

- Dung: None; 10 tons per acre.
- Nitrogen: None; 0.6 cwt N per acre applied as sulphate of ammonia.
- Phosphate: None; 0.6 cwt P₂O₅ per acre applied as superphosphate.
- Potash: None; 1.0 cwt K₂O per acre applied as muriate of potash.

Basal dressing to wheat: 4 cwt nitrochalk per acre; 21 cwt ground chalk per acre.

Cultivations, etc.: Ploughed: Jan 21, 1955. Chalk applied: Mar 31. Nitrogen applied, seed drilled at 2 bushels per acre: Apr 1. Sprayed with DNOC at 8 lb per acre in 80 gallons: May 2. Combine harvested: Sept 1. Variety: Koga II. Previous crop: Potatoes.

Standard error per plot:

Grain: 2.20 cwt per acre or 7.5% (18 d.f.)

For details of the preceding potato experiment see 54/Cd/1.

Summary of Results

Grain: Mean yield 29.5 cwt per acre

Responses to treatments

| Response to | Mean | Dung: tons per acre | | cwt per acre | | | | | |
|-------------------------------|---------|---------------------|------|--------------|------|-------------------------------|------|------------------|------|
| | | None | 10 | N | | P ₂ O ₅ | | K ₂ O | |
| | | | | None | 0.6 | None | 0.6 | None | 1.0 |
| | (±0.78) | (±1.10) | | | | | | | |
| Dung | +1.2 | - | - | +1.1 | +1.3 | +1.5 | +0.9 | +0.5 | +1.9 |
| N | +1.7 | +1.6 | +1.8 | - | - | +0.8 | +2.6 | +1.4 | +2.0 |
| P ₂ O ₅ | +1.4 | +1.7 | +1.1 | +0.5 | +2.3 | - | - | +2.0 | +0.8 |
| K ₂ O | -0.1 | -0.8 | +0.6 | -0.4 | +0.2 | +0.5 | -0.7 | - | - |

Mean dry matter % as harvested: 85.0

55/Ca/6.1

SPRING WHEAT

Rates and times of application of nitrogen - Rothamsted (R) Great Field I and Woburn (W) Butt Close.

Design (each field): 22 treatments arranged in 4 blocks of 13 plots each, the control and 3 treatments occurring in every block, the other 18 treatments occurring in 2 blocks. The total amounts of N applied per block were equal.

Area of each plot: 0.0212 acre. Area harvested: 0.0140 acre.

Treatments: None, and all combinations of:-

Nitrogen: 0.3; 0.6; 0.9 cwt N per acre applied as 'Nitro-Chalk'.

Times of application: All in seed bed (S); all as early top dressing (E); all as late top dressing (L); $\frac{1}{2}$ S & $\frac{1}{2}$ E; $\frac{1}{2}$ S & $\frac{1}{2}$ L; $\frac{1}{2}$ E & $\frac{1}{2}$ L; $\frac{1}{3}$ S, $\frac{1}{3}$ E & $\frac{1}{3}$ L.

Basal dressing: 1.15 cwt per acre compound fertilizer (13% P₂O₅; 13% K₂O) combine drilled with the seed.

Cultivations, etc.:

Great Field I (R). Ploughed: Nov - Dec 1954. Seed bed 'Nitro-Chalk' applied: Mar 31, 1955. Seed drilled at $2\frac{3}{4}$ bushels per acre with basal fertilizer: Apr 4. Early 'Nitro-Chalk' top dressing applied: Apr 30. Sprayed with D.N.C. 8 lb active material at 80 gallons per acre: May 11. Late 'Nitro-Chalk' top dressing applied: May 19. Combine harvested: Sept 1. Variety: Koga II. Previous crop: Potatoes.

Butt Close (W). Ploughed: Dec 23-28, 1954. 2 tons ground chalk per acre applied: Feb 3, 1955. Seed bed 'Nitro-Chalk' applied, seed drilled at $2\frac{3}{4}$ bushels per acre with basal fertilizer: Mar 22. Early 'Nitro-Chalk' top dressing applied: Apr 20. Late 'Nitro-Chalk' top dressing applied: May 16. Sprayed with MCPA amine at low volume: May 22. Combine harvested: Aug 23. Variety: Koga II. Previous crop: Potatoes.

Standard errors per plot. Grain: cwt per acre.

Great Field I (R): 2.01 cwt per acre or 4.5% (27 d.f.)

Butt Close (W): 2.87 cwt per acre or 11.1% (27 d.f.)

55/Ca/6.2

Summary of Results

Grain: cwt per acre

Rothamsted Great Field I

| | Time of application | | | | | | | Mean |
|-----------------|---------------------|------|------|----------------------------|----------------------------|----------------------------|--|---------------------|
| | S | E | L | $\frac{1}{2}S\frac{1}{2}E$ | $\frac{1}{2}S\frac{1}{2}L$ | $\frac{1}{2}E\frac{1}{2}L$ | $\frac{1}{3}S\frac{1}{3}E\frac{1}{3}L$ | |
| | (±1.52) | | | (±1.01) | | | | (±0.50) |
| N: cwt per acre | | | | | | | | 41.5 ⁽¹⁾ |
| None | | | | | | | | 44.6 |
| 0.3 | 44.1 | 42.9 | 44.5 | 46.0 | 45.9 | 44.9 | 44.1 | 45.1 |
| 0.6 | 45.9 | 45.0 | 43.7 | 43.7 | 45.8 | 44.1 | 46.4 | 45.3 |
| 0.9 | 46.1 | 44.4 | 45.7 | 45.8 | 44.3 | 45.9 | 45.0 | |
| Mean (±0.84) | 45.3 | 44.1 | 44.7 | 45.2 | 45.3 | 44.9 | 45.2 ⁽²⁾ | 44.7 |

(1) ±1.01 (2) ±0.58

Mean dry matter % as harvested: 84.5

Woburn Butt Close

| | Time of application | | | | | | | Mean |
|-----------------|---------------------|------|------|----------------------------|----------------------------|----------------------------|--|---------------------|
| | S | E | L | $\frac{1}{2}S\frac{1}{2}E$ | $\frac{1}{2}S\frac{1}{2}L$ | $\frac{1}{2}E\frac{1}{2}L$ | $\frac{1}{3}S\frac{1}{3}E\frac{1}{3}L$ | |
| | (±2.16) | | | (±1.43) | | | | (±0.72) |
| N: cwt per acre | | | | | | | | 12.2 ⁽¹⁾ |
| None | | | | | | | | 22.2 |
| 0.3 | 26.4 | 21.0 | 24.0 | 21.0 | 21.3 | 22.4 | 20.8 | 28.9 |
| 0.6 | 32.7 | 29.1 | 22.9 | 28.7 | 29.9 | 29.9 | 29.0 | 29.7 |
| 0.9 | 33.4 | 25.9 | 26.4 | 30.8 | 30.6 | 23.7 | 33.5 | |
| Mean (±1.20) | 30.8 | 25.3 | 24.4 | 26.8 | 27.3 | 25.4 | 27.7 ⁽²⁾ | 25.8 |

(1) ±1.43 (2) ±0.83

Mean dry matter % as harvested: 84.1

Time of application

- S In Seedbed.
- E Early top dressing.
- L Late top dressing.

55/Ca/7

SPRING WHEAT

Varieties and levels of nitrogen - Great Field I 1955.

Design: 3 randomized blocks of 5 plots each, plots being split into 2 for the application of nitrogen.

Area of each sub plot: 0.0083 acre. Area harvested: 0.0057 acre.

Treatments: All combinations of:

Whole plots. Varieties: Atle; Atson; Koga II; Peko; Progress.
Sub plots. Nitrogen: 0.3; 0.6 cwt N per acre applied as nitrochalk.

Basal dressing: 1.15 cwt compound granular fertilizer (13% P₂O₅, 13% K₂O) per acre, combine drilled with seed.

Cultivations, etc.: Ploughed: Dec 29, 1954. Nitrogen applied, seed combine drilled at 2½ bushels per acre: Apr 14, 1955. Sprayed with DNOC at 6 lb per acre in 80 gallons: May 12. Combine harvested: Aug 31. Previous crop: Potatoes.

Standard errors per plot, Grain:

Whole plot: 1.75 cwt per acre or 4.7% (8 d.f.)
 Sub plot: 1.35 cwt per acre or 3.6% (10 d.f.)

Summary of Results

Grain: cwt per acre

| | Atle | Atson | Variety Koga II | Peko | Progress | Mean |
|-----------------|----------|-------|--------------------|------|----------|-----------------|
| | (±1.15)* | | | | | |
| N: cwt per acre | | | | | | |
| 0.3 | 35.9 | 35.0 | 40.6 | 38.3 | 38.0 | 37.6 |
| 0.6 | 33.4 | 32.8 | 41.0 | 38.0 | 37.4 | 36.5 |
| Mean (±1.01) | 34.7 | 33.9 | 40.8 | 38.1 | 37.7 | 37.0 |
| Diff. (±1.10) | -2.5 | -2.2 | +0.4 | -0.3 | -0.6 | -1.1 (±0.49) |

*for use in comparisons other than vertical.

Mean dry matter % as harvested: 83.5

55/Cb/1

BARLEY

Seed rates and levels of nitrogen - Great Field I 1955.

Design: 3 randomized blocks of 12 plots each.

Area of each plot: 0.0140 acre.

Treatments: All combinations of:-

Seed rates: 1; 2; 3 bushels per acre.

Nitrogen: None; 0.3; 0.6; 0.9 cwt N per acre applied as sulphate of ammonia.

Basal dressing: 1.15 cwt compound granular fertilizer (13% P₂O₅, 13% K₂O) per acre combine drilled. 1 ton ground chalk per acre.

Cultivations, etc.: Ploughed: Dec 29, 1954. Chalk and sulphate of ammonia applied: April 1, 1955. Seed combine drilled: April 2. Sprayed with D.N.O.C. at 6 lb. per acre in 80 gallons: May 11. Combine harvested: Aug 20. Variety: Proctor. Previous crop: Potatoes.

Standard error per plot:

Grain (at 85% D.M.): 2.43 cwt per acre or 5.7% (22 d.f.).

Summary of Results

| Seed rate: bushels per acre | N: cwt per acre | | | | Mean |
|---|-----------------|---------|------|------|---------|
| | None | 0.3 | 0.6 | 0.9 | |
| Grain (at 85% dry matter): cwt per acre | | | | | |
| | | (±1.40) | | | (±0.70) |
| 1 | 41.4 | 44.7 | 41.7 | 42.9 | 42.7 |
| 2 | 43.3 | 44.6 | 44.7 | 40.7 | 43.3 |
| 3 | 41.4 | 42.4 | 43.2 | 41.6 | 42.1 |
| Mean (±0.81) | 42.1 | 43.9 | 43.2 | 41.7 | 42.7 |
| Straw: cwt per acre | | | | | |
| 1 | 27.7 | 33.0 | 30.9 | 32.8 | 31.1 |
| 2 | 27.7 | 31.9 | 33.0 | 35.3 | 32.0 |
| 3 | 25.5 | 29.6 | 33.2 | 36.6 | 31.2 |
| Mean | 27.0 | 31.5 | 32.4 | 34.9 | 31.4 |

Mean dry matter % as harvested, Grain: 82.6

Records of incidence of disease (Eyespot and Take-All), estimates of % area lodged and counts of ear emergence and plant, straw and ear numbers, were made.

BARLEY

Rates and times of application of nitrogen - Rothamsted (R) Great Field I and Woburn (W) Butt Close.

Design (each field): 22 treatments arranged in 4 blocks of 13 plots each, the control and 3 treatments occurring in every block, the other 18 treatments occurring in 2 blocks. The total amounts of N applied per block were equal.

Area of each plot: 0.0212 acre. Area harvested: 0.0141 acre.

Treatments: None, and all combinations of:-

Nitrogen: N_1 ; N_2 ; N_3 .

Times of application: All in seed bed (S); all as early top dressing (E); all as late top dressing (L); $\frac{1}{2}$ S & $\frac{1}{2}$ E; $\frac{1}{2}$ S & $\frac{1}{2}$ L; $\frac{1}{2}$ E & $\frac{1}{2}$ L; $\frac{1}{3}$ S, $\frac{1}{3}$ E & $\frac{1}{3}$ L.

Where N_1 ; N_2 ; N_3 =

Great Field I (R): 0.23; 0.46; 0.69 cwt N per acre applied as 'Nitro-Chalk'.

Butt Close (W): 0.3; 0.6; 0.9 cwt N per acre applied as 'Nitro-Chalk'.

Basal dressing: 1.15 cwt per acre compound fertilizer (13% P_2O_5 , 13% K_2O) combine drilled with the seed.

Cultivations, etc.:

Great Field I (R). Ploughed: Nov - Dec 1954. Seed drilled at 2 bushels per acre with basal fertilizer, and seed bed 'Nitro-Chalk' applied: Apr 2, 1955. Early 'Nitro-Chalk' top dressing applied: Apr 30. Sprayed with DNC 8 lb active material in 80 gallons per acre: May 11. Late 'Nitro-Chalk' top dressing applied: May 19. Combine harvested: Aug 21. Variety: Herta. Previous crop: Potatoes.

Butt Close (W). Ploughed: Dec 23-28, 1954. Applied ground chalk at 2 tons per acre: Feb 4, 1955. Seed bed 'Nitro-Chalk' applied: Mar 21. Seed drilled at 2 bushels per acre with basal fertilizer: Mar 22. Early top dressing of 'Nitro-Chalk' applied: Apr 20. Late top dressing of 'Nitro-Chalk' applied: May 16. Sprayed with MCPA amine at low volume: May 22. Combine harvested: Aug 22. Variety: Herta. Previous crop: Potatoes.

Standard errors per plot. Grain: cwt per acre.

Great Field I(R): 1.68 cwt per acre or 3.4% (27 d.f.)

Butt Close (W): 3.84 cwt per acre or 9.9% (27 d.f.)*

Note: The Woburn crop was severely and irregularly damaged by birds. Estimates of damage were made before harvesting and the yields have been corrected on the basis of these.

*At 85% dry matter.

Summary of Results

Grain: cwt per acre

Rothamsted Great Field I

| | Time of application | | | | | | | Mean |
|---------------------|---------------------|------|------|----------------------------|----------------------------|----------------------------|--|----------------|
| | S | E | L | $\frac{1}{2}S\frac{1}{2}E$ | $\frac{1}{2}S\frac{1}{2}L$ | $\frac{1}{2}E\frac{1}{2}L$ | $\frac{1}{3}S\frac{1}{3}E\frac{1}{3}L$ | |
| N: cwt per acre | (± 1.26) | | | | | | (± 0.84) | (± 0.42) |
| None | | | | | | | 46.9 ⁽¹⁾ | |
| 0.23 | 48.2 | 47.7 | 49.8 | 46.2 | 47.1 | 47.6 | 48.1 | |
| 0.46 | 50.5 | 51.2 | 47.7 | 49.5 | 50.2 | 48.9 | 49.7 | |
| 0.69 | 48.0 | 46.6 | 51.5 | 48.5 | 47.7 | 48.2 | 48.5 | |
| Mean (± 0.70) | 48.9 | 48.5 | 49.7 | 48.0 | 48.3 | 48.2 | 49.3 ⁽²⁾ | 48.7 |

(1) ± 0.84 (2) ± 0.48

Mean dry matter % as harvested: 85.0

Grain (at 85% dry matter): cwt per acre

Woburn Butt Close

| | Time of application | | | | | | | Mean |
|---------------------|---------------------|------|------|----------------------------|----------------------------|----------------------------|--|----------------|
| | S | E | L | $\frac{1}{2}S\frac{1}{2}E$ | $\frac{1}{2}S\frac{1}{2}L$ | $\frac{1}{2}E\frac{1}{2}L$ | $\frac{1}{3}S\frac{1}{3}E\frac{1}{3}L$ | |
| N: cwt per acre | (± 2.89) | | | | | | (± 1.92) | (± 0.96) |
| None | | | | | | | 21.2 ⁽¹⁾ | |
| 0.3 | 29.7 | 34.7 | 34.0 | 34.8 | 32.6 | 32.2 | 32.6 | |
| 0.6 | 41.4 | 44.4 | 42.0 | 37.3 | 36.2 | 39.3 | 41.2 | |
| 0.9 | 46.5 | 50.5 | 41.9 | 48.7 | 44.7 | 45.4 | 46.8 | |
| Mean (± 1.61) | 39.2 | 43.2 | 39.3 | 40.3 | 37.8 | 38.9 | 41.4 ⁽²⁾ | 38.7 |

(1) ± 1.92 (2) ± 1.11

Mean dry matter % as harvested: 84.5

Time of application

- S In Seedbed.
- E Early top dressing.
- L Late top dressing.

55/Cb/3

BARLEY

Varieties and levels of nitrogen (Growth Study) - Great Field I 1955.

Design: 6 x 6 Latin square.

Area of each plot: 0.0223 acre. Area harvested: 0.0118 acre.

Treatments: All combinations of:

Varieties: Herta; Plumage Archer; Proctor.

Nitrogen: None; 0.46 cwt N per acre applied as nitrochalk.

Basal dressing: 1.15 cwt compound granular fertilizer (13% P₂O₅, 13% K₂O) per acre, combine drilled.

Cultivations, etc.: Ploughed: Dec 29, 1954. Nitrochalk applied, seed combine drilled with placement machine at 2½ bushels per acre: Apr 2, 1955. Sprayed with DNOC at 6 lb per acre in 80 gallons: May 11. Combine harvested: Aug 22. Previous crop: Potatoes.

Standard error per plot:

Grain (at 85% D.M.): 2.45 cwt per acre or 5.0% (20 d.f.).

Records were made of the following:-

At harvest:- Shoot number, 1000 corn weight
Straw height, N analyses of grain, straw and chaff.

At intermediate samplings (at fortnightly intervals from May-August):-

Dry weight of straw and ears
Shoot number
Leaf area
Straw height.

Summary of Results

Grain (at 85% dry matter): cwt per acre

| N: cwt per acre | Herta | Variety Plumage Archer | Proctor | Mean |
|--------------------|-------|------------------------------|---------|---------|
| | | (±1.00) | | |
| None | 48.5 | 43.9 | 51.0 | 47.8 |
| 0.46 | 55.6 | 40.6 | 52.7 | 49.6 |
| Mean (±0.71) | 52.0 | 42.3 | 51.8 | 48.7 |
| Difference (±1.42) | +7.1 | -3.3 | +1.7 | +1.8 |
| | | | | (±0.82) |

Mean dry matter % as harvested: 87.6

55/Cb/4.1

BARLEY

Residual effects of phosphate and potash - Highfield V 1955.

Design: 8 randomized blocks of 8 plots each, the interaction PKS × placement being confounded with block differences.

Area of each plot: 0.00923 acre.

Treatments, applied to lucerne in 1952: All combinations of:-
Phosphate: None; 1.0 cwt P_2O_5 per acre applied as superphosphate.
Potash: None; 1.0 cwt K_2O per acre applied as muriate of potash.
Method of placement: Broadcast on seedbed; ploughed in 10".
Starter: None; 2 cwt granular superphosphate per acre placed beneath seed.

Note: the division of each plot into sub plots in 1954 was ignored in 1955.

Basal dressing: 1 cwt nitrochalk per acre.

Cultivations, etc.: Ploughed: Mar 21, 1955. Seed drilled at $2\frac{3}{4}$ bushels per acre: Apr 6. Nitrochalk applied: Apr 7. Sprayed with 2, 4D, medium volume, $2\frac{1}{2}$ pints per acre: May 20. Combine harvested: Aug 17. Variety: Herta. Previous crop: Lucerne.

Standard error per plot.

Grain (at 85% D.M.): 1.81 cwt per acre or 3.9% (42 d.f.)

For previous years' results see 52/Cf/1, 53/Cg/1 and 54/Ce/1.

55/Cb/4.2

Summary of Results

| Starter | Treatments applied 1952 | | | | | | | | Mean |
|---|-------------------------|----------------------------------|----------------|--|----------------|--|----------------|---------------------|------|
| | No ferti- lizer | Superphosphate Broad- cast | Ploughed in | Muriate of Potash Broad- cast | Ploughed in | Superphosphate and Muriate of Potash Broad- cast | Ploughed in | | |
| Grain (at 85% Dry Matter): cwt per acre | | | | | | | | | |
| | (±0.64) | | | (±0.90) | | | | | |
| None | 44.4 | 45.3 | 46.8 | 45.7 | 46.5 | 47.5 | 45.2 | 45.7 | |
| Super | 44.9 | 45.8 | 44.2 | 47.2 | 46.9 | 47.8 | 45.7 | 45.9 | |
| Mean | 44.7 ⁽¹⁾ | 45.5 | 45.5 | 46.5 | 46.7 | 47.6 | 45.4 | 45.8 | |
| Difference (±1.28) | +0.5 ⁽²⁾ | +0.5 | -2.6 | +1.5 | +0.4 | +0.3 | +0.5 | +0.2 ⁽¹⁾ | |

(1) ±0.45

(2) ±0.90

Straw (at 85% Dry Matter): cwt per acre

| | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|
| None | 34.5 | 36.7 | 38.8 | 35.9 | 37.7 | 42.2 | 38.9 | 37.4 |
| Super | 36.4 | 35.6 | 36.8 | 40.1 | 37.4 | 41.8 | 36.5 | 37.6 |
| Mean | 35.4 | 36.1 | 37.8 | 38.0 | 37.6 | 42.0 | 37.7 | 37.5 |
| Difference | +1.9 | -1.1 | -2.0 | +4.2 | -0.3 | -0.4 | -2.4 | +0.2 |

Mean dry matter % as harvested, Grain: 84.0
Straw: 82.6

55/Cc/1

SPRING BEANS

The control of Black Aphids (*Aphis Fabae*) by spraying and time of sowing - Fosters 1955.

Design: 4 x 4 Latin square, plots split into 2 for the application of spray.

Area of each sub plot: 0.0189 acre.

Treatments: All combinations of:-

Whole plots. Times of sowing: Mar 19; April 6; April 25; May 13.

Sub plots. Spray: None; "Metasystox" (0.05% active ingredient) at 100 gallons per acre.

Note: The beans sown on May 13 were sprayed twice.

Basal dressing: 6 cwt compound granular fertilizer (10% P₂O₅, 20% K₂O) per acre.

Cultivations, etc.: Ploughed: Oct 22, 1954. Basal fertilizer applied, seed drilled at 195 lb per acre: Mar 19, 1955, April 6, April 25, May 13 respectively. Sprayed with "Metasystox": June 23. Last sown crop sprayed with "Metasystox" for second time: July 22. Combine harvested first three sowings: Aug 27. Combine harvested last sowing: Sept 12. Variety: Garton's Tick. Previous crop: Barley.

Standard errors per plot. Grain (at 85% dry matter):

Whole plot: 1.49 cwt per acre or 12.0% (6 d.f.)

Sub plot: 1.54 cwt per acre or 12.4% (12 d.f.)

Counts of Black Aphids and of aphid predators were made at weekly intervals from June to August.

Summary of Results

Grain (at 85% dry matter): cwt per acre

| Spray | Time of sowing | | | | Mean |
|--------------------|----------------|-------|--------|--------|---------|
| | Mar 19 | Apr 6 | Apr 25 | May 13 | |
| | (±0.92)* | | | | |
| None | 16.9 | 13.1 | 6.7 | 2.4 | 9.8 |
| Metasystox | 19.2 | 18.3 | 12.0 | 10.5 | 15.0 |
| Mean (±0.74) | 18.0 | 15.7 | 9.3 | 6.4 | 12.4 |
| Difference (±1.09) | 2.3 | 5.2 | 5.3 | 8.1 | 5.2 |
| | | | | | (±0.54) |

* for use in comparisons other than vertical.

Mean dry matter % as harvested: 82.4.

55/Cc/2.1

SPRING BEANS

Control of weeds by spraying and cultivation - Fosters 1955.

Design: 4 randomized blocks of 5 plots each, 2 blocks being sprayed and 2 unsprayed.

Area of each plot: 0.0202 acre. Area harvested: 0.0126 acre.

Treatments: All combinations of:-

Spraying (on blocks): None; DNBP high volume

Additional cultivations: None (0)

Harrowed once (1)

Harrowed three times (2)

Mechanical weeder once (3)

Mechanical weeder three times (4)

Basal dressing: 6 cwt compound granular fertilizer (10% P₂O₅, 20% K₂O) per acre.

Basal inter-row cultivation: One early horse hoeing.

Cultivations, etc.: Ploughed: Oct 22, 1954. Basal fertilizer applied, seed drilled at 195 lb per acre: Mar 19, 1955. Horse hoed all plots: May 11. Combine harvested: Aug 27. Variety: Spring Tick. Previous crop: Barley.

Treatment cultivations:

(1) Harrowed: April 30.

(2) Harrowed: April 30, May 6, May 31.

(3) Mechanical weeder: April 30.

(4) Mechanical weeder: April 30, May 6, May 31.

DNBP at 4 pts in 100 gallons per acre sprayed: May 26.

Standard error per plot:

Grain: 1.07 cwt per acre or 6.3% (8 d.f.)

55/Cc/2.2

Summary of Results

Grain: cwt per acre

| Spray | Treatment cultivations | | | | | Mean |
|--|------------------------|------|-----------------------------|------|------|------|
| | 0 | 1 | 2 | 3 | 4 | |
| | | | | | | |
| | | | (± 0.76) [Ⓜ] | | | |
| None | 18.5 | 17.5 | 14.3 | 18.5 | 16.4 | 17.0 |
| DNBP | 17.1 | 17.6 | 15.1 | 17.4 | 17.7 | 17.0 |
| Mean (± 0.54) | 17.8 | 17.5 | 14.7 | 18.0 | 17.1 | 17.0 |
| Difference (± 1.07) ⁺ | -1.4 | +0.1 | +0.8 | -1.1 | +1.3 | 0.0 |

[Ⓜ]for use in horizontal comparisons only

⁺for use in the comparison of two differences only

Mean dry matter % as harvested: 86.2

Treatment cultivations

- (0) No extra cultivation.
- (1) Harrowed once.
- (2) Harrowed three times.
- (3) Mechanical weeder once.
- (4) Mechanical weeder three times.

SPRING BEANS

Flower drop - Hormone sprays - Fosters 1955.

Design: 4 randomized blocks of 7 plots each.

Area of each plot: 0.00550 acre. Area harvested: 0.00458 acre.

Treatments: Hormone sprays.

- None (3 plots per block). (0)
- 2; 3 applications of 4 chlorophenoxyacetic acid. (1) & (2)
- 2; 3 applications of α (2:4:5 trichlorophenoxy) propionic acid. (3) & (4)

The sprays, at a concentration of 5 p.p.m., were applied during the flowering period in successive doses at the following rates per acre:-

Treatments (1), (2), (3) and (4): 120 gallons and later an additional 200 gallons.

Treatments (2) and (4): an additional 320 gallons as a final application.

Basal dressing: 9 cwt compound granular fertilizer (10% P₂O₅, 20% K₂O) per acre.

Cultivations, etc.: Ploughed: Oct 19, 1954. Basal fertilizer applied, seed sown at 190 lb per acre: Mar 30, 1955. Sprayed with hormone sprays: June 24, July 1, July 9. Combine harvested: Aug 27. Variety: Gartons' Tick. Previous crop: Barley.

Standard error per plot:

Grain: 2.41 cwt per acre or 20.4% (20 d.f.)

N.B. Counts of numbers of pods were made. Damage by Aphids was severe and irregular.

Summary of Results

Grain: cwt per acre

| | 0 | 1 | 2 | 3 | 4 | Mean |
|-------------------------|---------------------|------|------|------|------|------|
| Mean (± 1.21) | 12.0 ⁽¹⁾ | 13.9 | 12.0 | 9.7 | 11.0 | 11.8 |
| Increase (± 1.39) | | +1.9 | 0.0 | -2.3 | -1.0 | |

(1) ± 0.70 .

Mean dry matter % as harvested: 84.8

- Treatments: 0 No spray
- 1 2 spray applications } 4 chlorophenoxyacetic acid.
- 2 3 spray applications }
- 3 2 spray applications } α (2:4:5 trichlorophenoxy)
- 4 3 spray applications } propionic acid.

55/cd/1

POTATOES

Placement of Nitrogen and Potash - Little Hoos 1955.

Design: 6 randomized blocks of 12 plots each, with levels of N by levels of K partially confounded with block differences.

Area of each plot: 0.0140 acre. Area harvested: 0.0057 acre.

Treatments: All combinations of:-

Nitrogen: None; 0.5; 1.0 cwt N per acre as sulphate of ammonia.

Potash: None; 0.75; 1.5 cwt K₂O per acre as sulphate of potash.

Methods of placement: Broadcast on flat before planting; Side band placement at planting.

Basal dressing: 1.0 cwt P₂O₅ per acre as superphosphate, placement drilled as above.

Cultivations, etc.: Ploughed: Oct 7, 1954. Broadcast fertilizers applied, machine planted with placed fertilizers: Apr 23, 1955. Earthed up: July 6. Sprayed with sulphuric acid, 20% B.O.V.: Sept 28. Hand dug: Oct 7. Variety: Majestic. Previous crop: Beans.

Standard error per plot:

Total tubers: 0.589 tons per acre or 11.4% (31 d.f.)

Summary of Results

Total tubers: tons per acre

| K ₂ O: cwt per acre | N: cwt per acre | | | | | Mean |
|-----------------------------------|-----------------|-----------|------|--------|------|----------|
| | None | Broadcast | | Placed | | |
| | | 0.5 | 1.0 | 0.5 | 1.0 | |
| | (±0.227) | (±0.321) | | | | (±0.131) |
| None | 4.85 | 4.53 | 4.41 | 4.79 | 4.47 | 4.65 |
| | (±0.321) | (±0.455) | | | | (±0.186) |
| Broadcast | | | | | | |
| 0.75 | 4.37 | 4.92 | 4.59 | 3.95 | 6.37 | 4.76 |
| 1.5 | 4.95 | 5.64 | 6.70 | 5.27 | 6.08 | 5.60 |
| Placed | | | | | | |
| 0.75 | 4.67 | 5.73 | 7.59 | 6.07 | 5.65 | 5.73 |
| 1.5 | 4.08 | 5.67 | 6.95 | 6.55 | 6.70 | 5.67 |
| Mean | 4.63 | 5.17 | 5.77 | 5.24 | 5.62 | 5.18 |
| | (±0.131) | (±0.186) | | | | |

55/cd/2.1

POTATOES

Control of virus spread by insecticides - Little Hoos 1955.

Design: 5 × 5 Latin square.

Area of each plot: 0.0602 acre. Area harvested: 0.0139 acre.

Treatments:

No. of insecticide sprayings: None; 2; 4; 6; 8 times sprayed with D.D.T. emulsion, 2 lb. active ingredient, in 100 gallons per acre.

Infector plants: 6 leaf roll and 6 virus Y infected plants planted in each plot.

Note: The tractor used for spraying was driven over all plots on each occasion always passing over the same rows. Yields were taken from the undamaged rows and an estimate of the loss in yield due to tractor damage was made from an area of 0.0820 acre.

Basal dressing: 13 cwt per acre compound granular fertilizer (7% N, 7% P₂O₅, 10¹/₂% K₂O).

Cultivations, etc.: Ploughed: Oct 7, 1954. Machine planted with fertilizer placement: Apr 23, 1955. Earthed up: July 7. Sprayed with sulphuric acid, 15% B.O.V.: Sept 29. Lifted: Oct 7. Variety: Majestic. Previous crop: Beans.

| Sprayings: | 2 | 4 | 6 | 8 |
|------------|---------|---------|---------|---------|
| | June 13 | June 13 | June 13 | June 3 |
| | July 14 | July 5 | June 23 | June 13 |
| | | July 27 | July 5 | June 23 |
| | | Aug 10 | July 14 | July 5 |
| | | | July 27 | July 14 |
| | | | Aug 10 | July 27 |
| | | | | Aug 10 |
| | | | | Aug 26 |

Standard error per plot:

Total tubers: 0.583 tons per acre or 12.3% (12 d.f.)

Note: Aphid counts were made and tuber samples taken to assess virus spread.

55/ca/2.2

Summary of Results

| | Number of sprayings with D.D.T. | | | | | Mean |
|-----------------------------|---------------------------------|-------|-------|-------|-------|------|
| | None | 2 | 4 | 6 | 8 | |
| Total tubers: tons per acre | | | | | | |
| Mean (± 0.261) | 4.59 | 4.68 | 4.72 | 4.94 | 4.86 | 4.76 |
| Increase (± 0.369) | | +0.09 | +0.13 | +0.35 | +0.27 | |

Estimated loss in yield in damaged rows due to:-

8 passages of the tractor: 21.5%

Estimated loss in yield in whole crop due to:-

8 passages of the tractor along 4 rows out of 10: 8.6%

55/Ca/3.1

POTATOES

The control of Blight by Copper Spray - Stackyard 1955.

Design: 5 randomized blocks of 2 plots each, plots being split into 2 for determination of the effect of tractor damage.

Note: There were originally 6 blocks, but the produce from 2 whole plots was accidentally mixed at lifting.

Area of each sub plot: 0.0140 acre.

Treatments:

Whole plots. No spray; Copper fungicide 5 lb in 45 gallons per acre sprayed twice. The tractor used for spraying was driven over all the plots on each occasion.

Sub plots. 4 rows damaged by two passages of the tractor were compared with 4 undamaged rows.

Basal dressing: 20 cwt compound granular fertilizer (7% N, 7% P₂O₅, 10½% K₂O) per acre.

Cultivations, etc.: Ploughed: Oct 14, 1954. Sprayed with T.C.A., 20 lb in 40 gallons: Dec 6 and again Feb 2, 1955. Potatoes planted by machine with placed fertilizer: Apr 23. Earthed up: July 18. Fungicide treatment applied: Aug 4 and again Aug 29. Sprayed with sulphuric acid, 20% B.O.V.: Oct 4. Lifted: Oct 26. Variety: Majestic. Previous crop: Barley.

Standard errors per plot. Total tubers:

Whole plot: 1.09 tons per acre or 16.9% (4 d.f.)

Sub plot: 0.766 tons per acre or 11.8% (8 d.f.)

Note: Blight counts were made.

55/ca/3.2

Summary of Results

| | None | Spray Copper fungicide | Mean |
|-----------------------------|-------------|------------------------------|----------|
| Total tubers: tons per acre | | | |
| | (1) and (2) | | (±0.242) |
| Undamaged rows | 6.09 | 6.77 | 6.43 |
| Damaged rows | 6.51 | 6.52 | 6.51 |
| Mean (±0.488) | 6.30 | 6.65 | 6.47 |

(1) ±0.342 for use in vertical comparisons only.

(2) ±0.545 for use in all others.

Percentage ware (1½" riddle)

| | | | |
|----------------|------|------|------|
| Undamaged rows | 76.3 | 77.1 | 76.7 |
| Damaged rows | 77.8 | 79.2 | 78.5 |
| Mean | 77.0 | 78.2 | 77.6 |

55/Cd/4.1

POTATOES

The control of Blight by Copper and Sulphuric Sprays - Little Knott 1955.

Design: 4 × 4 Latin square, plots being split into 2 for determination of the effect of tractor damage.

Area of each sub plot: 0.0140 acre.

Treatments:

Whole plots: No spray; Copper fungicide 5 lb in 45 gallons per acre sprayed twice; 100 gallons sulphuric acid, 20% B.O.V. sprayed to destroy haulm; Copper fungicide and sulphuric acid sprayed as above. The tractor used for spraying was driven over all the plots on each occasion.

Sub plots: 4 rows damaged by two passages of the tractor were compared with 4 undamaged rows.

Basal dressing per acre: 10 tons dung; 20 cwt compound granular fertilizer (7% N, 7% P₂O₅, 10 $\frac{1}{2}$ % K₂O).

Cultivations, etc.: Ploughed: Sept 29, 1954. Dung applied and cultivated in: Mar 18, 1955. Potatoes planted by machine with placed fertilizer: Apr 27. Earthed up: July 4. Fungicide treatment applied: Aug 18 and Sept 7. Sulphuric acid treatment applied: Sept 13. Lifted: Oct 26. Variety: King Edward. Previous crop: Beans - N. end; Wheat - S. end.

Standard errors per plot: Total tubers.

Whole plot: 0.875 tons per acre or 12.8% (6 d.f.)

Sub plot: 0.291 tons per acre or 4.2% (12 d.f.)

Note: Blight Counts were made.

55/Cd/4.2

Summary of Results

| | Spray | | | | Mean |
|-----------------------------|------------------|------------------|----------------|-------------------------------------|--------------------------|
| | None | Copper fungicide | Sulphuric acid | Copper fungicide and Sulphuric acid | |
| Total tubers: tons per acre | | | | | |
| | (± 0.449)* | | | | |
| Undamaged rows | 7.73 | 6.08 | 7.30 | 6.81 | 6.98 |
| Damaged rows | 7.53 | 5.81 | 6.97 | 6.66 | 6.74 |
| Mean (± 0.438) | 7.63 | 5.95 | 7.13 | 6.74 | 6.86 |
| Difference (± 0.206) | -0.20 | -0.27 | -0.33 | -0.15 | -0.24 (± 0.103) |

Percentage ware ($1\frac{1}{2}$ " riddle)

| | | | | | |
|----------------|------|------|------|------|------|
| Undamaged rows | 78.5 | 72.5 | 79.2 | 73.2 | 75.9 |
| Damaged rows | 78.5 | 64.4 | 73.9 | 73.9 | 72.7 |
| Mean | 78.5 | 68.5 | 76.6 | 73.6 | 74.3 |
| Difference | 0.0 | -8.1 | -5.3 | +0.7 | -3.2 |

*For use in comparisons other than vertical.

55/Ce/1

LUCERNE

Single and repeated applications of potash - Great Harpenden II 1955 - the first year.

Design: 6 randomized blocks of 8 plots each.

Area of each plot: 0.0147 acre.

Treatments 1955:

Potash seedbed dressings: None (2 plots per block); 0.33; 0.66; 1.0 (2 plots per block); 2.0; 3.0 cwt K_2O per acre applied as muriate of potash.

Note: It is intended to repeat the following dressings annually for two further years: None; 0.33; 0.66; 1.0 (1 plot per block) cwt K_2O per acre.

Basal dressing: 14 cwt per acre ground chalk, 2.3 cwt per acre superphosphate placed beneath seed.

Cultivations, etc.: Ploughed: Oct 12, 1954. Chalk applied: Apr 6, 1955. Potash applied: May 6. Seed drilled at 174 lb. per acre with placed superphosphate: May 9. Cut: Sept 14. Variety: Du Puits. Previous crop: Barley.

Standard error per plot:

Dry Matter: 1.01 cwt per acre or 17.7% (36 d.f.)

Note: The yields have been corrected for a linear fertility trend along the blocks.

Summary of Results

| | Dry matter: cwt per acre (1 cut) | | | | | | Mean |
|-------------------------|----------------------------------|------|------|--------------------|-----|-----|------|
| | K ₂ O: cwt per acre | | | | | | |
| | None | 0.33 | 0.66 | 1.0 | 2.0 | 3.0 | |
| Mean (± 0.41) | 5.2 ⁽¹⁾ | 5.7 | 5.8 | 5.6 ⁽¹⁾ | 6.3 | 6.1 | 5.7 |
| Increase (± 0.50) | | 0.5 | 0.6 | 0.4 ⁽²⁾ | 1.1 | 0.9 | |

(1) ± 0.29

(2) ± 0.41

Mean dry matter % as harvested: 36.9

BROCCOLI

Effect of dung and nitrogen on Virus spread - Great Knott I, 1955.

Design: 6 x 6 Latin square.

Area of each plot: 0.0207 acre. Area harvested: 0.0143 acre.

Treatments: All combinations of:-

Dung: None; 20 tons per acre.

Nitrogen: None; 0.6; 1.2 cwt N per acre applied as nitrochalk, half before planting, half in spring.

Basal dressing per acre: 4 tons ground chalk; 4 cwt superphosphate and 2 cwt muriate of potash.

Cultivations, etc.: Ploughed: Sept 30, 1953 and again Dec 7. Chalk applied: Feb 24, 1954. Ploughed: Feb 28. Dung applied, ploughed: June 4. Basal fertilizers applied: June 19. 1st application of nitrochalk: July 12. Planted: July 16. Gaps replanted: July 27, Aug 6, Aug 13, Aug 27, Aug 31. 2nd application of nitrochalk: Mar 17, 1955. Harvested: Apr 26 to May 5. Variety: Continuity. Previous crop: Wheat.

Note: The replanted broccoli failed.

Standard error per plot,

No. of saleable curds: 0.543 thousands per acre or 20.9% (20 d.f.)

Summary of Results

| Dung: tons per acre | N: cwt per acre | | | Mean |
|--|-----------------|-------|-------|----------------|
| | None | 0.6 | 1.2 | |
| Number of saleable curds: thousands per acre (±0.222) | | | | |
| None | 2.66 | 2.69 | 2.44 | 2.60 |
| 20 | 2.11 | 3.01 | 2.69 | 2.61 |
| Mean (±0.157) | 2.39 | 2.85 | 2.57 | 2.60 |
| Difference (±0.314) | -0.55 | +0.32 | +0.25 | +0.01 (±0.181) |
| Weight per saleable curd: lb | | | | |
| None | 1.28 | 1.27 | 1.45 | 1.33 |
| 20 | 1.44 | 1.42 | 1.56 | 1.48 |
| Mean | 1.36 | 1.35 | 1.50 | 1.40 |
| Difference | +0.16 | +0.15 | +0.11 | +0.15 |

General means: Total no. of curds: 4.57 thousands per acre
 Percentage saleable curds out of total no. of curds: 56.6

Records of incidence of cauliflower mosaic were made.

55/Cg/1

KALE

Placement of nitrogen, phosphate and potash - Great Harpenden II 1955.

Design: 4 randomized blocks of 10 plots each.

Area of each plot: 0.00909 acre. Area harvested: 0.00727 acre.

Treatments: None (2 plots per block) and all combinations of
Fertilizer: P; K; PK; NPK.

Method of application: Broadcast in seed bed; Drilled in band 2"
to side of seed and 3" below soil surface

where N = 0.4 cwt N per acre as sulphate of ammonia.
P = 0.6 cwt P₂O₅ per acre as superphosphate.
K = 1.0 cwt K₂O per acre as sulphate of potash.

In addition top dressings were applied:-

To "NPK" plots: 0.4 cwt N per acre as sulphate of ammonia.
To all other plots: 0.8 cwt N per acre as sulphate of ammonia.

Basal dressing: 7 cwt ground chalk per acre.

Cultivations, etc.: Ploughed: Oct 12, 1954. Ground chalk applied:
Apr 6, 1955. Broadcast fertilizers applied: Apr 20. Seed
drilled at 4 lb per acre with sideband fertilizer: Apr 21. Top
dressing applied: July 5. Cut: 2 blocks, Nov 29; remainder, Dec 6.
Variety: Marrow-stem. Previous crop: Barley.

Standard error per plot:

Yield: 1.55 tons per acre or 13.2% (28 d.f.)

Summary of Results

Yield: tons per acre

| N Top dressing: cwt per acre | 0.8 | | | | | 0.4 | Mean |
|---------------------------------|-------|----------|-------|-------|-------|-------------------|------|
| | None | P | K | PK | NPK | | |
| Treatments at sowing | | | | | | | |
| <u>Method of application</u> | | (±0.774) | | | | | |
| Broadcast | | 12.15 | 12.14 | 10.93 | 11.36 | 11.64 | |
| Drilled | | 12.11 | 11.49 | 12.77 | 11.82 | 12.05 | |
| Mean (±0.547) | 11.13 | 12.13 | 11.82 | 11.85 | 11.59 | 11.70 | |
| Difference (±1.094) | | -0.04 | -0.65 | +1.84 | +0.46 | +0.41 (±0.547) | |

55/E/1

CHEMICAL ANALYSES OF ORGANIC MATERIALS, USED IN THE THREE
COURSE ROTATION AND MARKET GARDEN EXPERIMENTS 1955

| Material | Percentage dry matter in sample | Percentage in dry matter | | | |
|-----------------------------------|---------------------------------------|--------------------------|------|-------------------------------|------------------|
| | | Organic matter | N | P ₂ O ₅ | K ₂ O |
| Three Course Rotation, Rothamsted | | | | | |
| Wheat Straw | 84.6 | 92.8 | 0.48 | 0.18 | 1.24 |
| Market Garden Experiment, Woburn | | | | | |
| Sewage Sludge* | | | | | |
| 1 | 45.4 | 41.2 | 2.59 | 3.74 | 0.21 |
| 2 | 45.8 | 41.9 | 2.53 | 3.42 | 0.16 |
| Sludge-Straw Compost* | | | | | |
| 1 | 39.8 | 43.2 | 2.43 | 2.94 | 0.47 |
| 2 | 35.1 | 45.3 | 2.36 | 2.88 | 0.37 |
| Vegetable Compost* | | | | | |
| 1 | 24.8 | 51.9 | 2.46 | 3.93 | 2.92 |
| 2 | 22.8 | 56.4 | 2.40 | 4.52 | 2.71 |
| Farmyard Manure* | | | | | |
| 1 | 23.1 | 60.0 | 3.16 | 4.34 | 3.70 |
| 2 | 21.2 | 61.1 | 1.96 | 1.00 | 1.29 |

*Sample 1 applied Sept, 1954; Sample 2 applied April, 1955.

METEOROLOGICAL RECORDS ROTHAMSTED 1955

(Departure from long period means in brackets)

| Month | Total sunshine: hours | Mean temperature: °F | | | Ground (2) frosts | Total rainfall: in. 1/1000 acre gauge | Rain (3) days | Drain- age through 20 in. soil: in. m.p.h. | Wind: (4) |
|-------|-----------------------|----------------------|-------|-----------------|-------------------|---------------------------------------|---------------|--|-----------|
| | | Air (1) | 1 ft. | In ground 4 ft. | | | | | |
| Jan. | 34 (-18) | 35.2 (-2.1) | 36.8 | 43.6 | 23 | 2.75 (+0.25) | 19 | 2.16 | 5.5 |
| Feb. | 78 (+9) | 33.4 (-4.9) | 36.7 | 42.1 | 20 | 2.31 (+0.38) | 19 | 1.08 | 8.7 |
| Mar. | 149 (+32) | 36.9 (-4.4) | 36.6 | 40.0 | 24 | 1.45 (-0.46) | 10 | 1.12 | 6.5 |
| Apr. | 173 (+16) | 48.4 (+2.6) | 46.7 | 43.1 | 10 | 0.49 (-1.46) | 9 | - | 5.4 |
| May | 208 (+12) | 49.7 (-2.2) | 50.1 | 47.0 | 1 | 4.62 (+2.47) | 16 | 1.96 | 6.2 |
| June | 153 (-50) | 57.0 (-0.3) | 57.6 | 50.1 | 0 | 2.11 (-0.06) | 9 | 0.55 | 4.8 |
| July | 249 (+54) | 62.8 (+2.1) | 64.7 | 55.4 | 0 | 0.25 (-2.32) | 4 | - | 4.3 |
| Aug. | 191 (+6) | 63.9 (+3.7) | 64.8 | 57.6 | 0 | 0.81 (-1.78) | 8 | - | 3.2 |
| Sept. | 155 (+9) | 57.7 (+1.7) | 58.6 | 57.6 | 0 | 1.88 (-0.50) | 14 | - | 3.3 |
| Oct. | 122 (+18) | 48.1 (-0.8) | 50.7 | 54.3 | 6 | 3.29 (+0.31) | 17 | 1.66 | 4.1 |
| Nov. | 43 (-20) | 43.9 (+1.5) | 44.5 | 49.8 | 10 | 2.07 (-0.76) | 18 | 1.69 | 3.8 |
| Dec. | 51 (+6) | 41.2 (+2.6) | 41.3 | 46.4 | 14 | 2.60 (+0.04) | 23 | 1.82 | 5.8 |
| Year | 1606 (+74) | 48.2 - | 49.0 | 48.9 | 108 | 24.63 (-3.89) | 166 | 12.04 | 5.1 |

(1) Mean of maximum and minimum.

(2) Number of nights grass minimum was 30°F or less.

(3) Mean or total.

(3) Number of days rainfall was 0.01 in. or more.
 (4) At 2 metres above ground level.

55/E/2.2

METEOROLOGICAL RECORDS WOBURN 1955

| Month | Total sun- shine: hours | Mean Temperature: °F | | Grass minimum: °F | Total rainfall: in. 8" gauge | Rain ⁽²⁾ days |
|-------------------|----------------------------------|-------------------------|--------------------|-------------------------|---------------------------------------|-----------------------------|
| | | Air ⁽¹⁾ | In ground 1 ft. | | | |
| January | 38 | 35.9 | 36.9 | 29.5 | 2.44 | 17 |
| February | 81 | 33.7 | 36.7 | 26.8 | 1.63 | 17 |
| March | 142 | 36.7 | 37.1 | 24.6 | 1.13 | 12 |
| April | 155 | 48.6 | 48.4 | 35.3 | 0.47 | 10 |
| May | 196 | 49.4 | 51.2 | 37.5 | 4.57 | 17 |
| June | 151 | 56.7 | 59.1 | 47.7 | 2.15 | 12 |
| July | 242 | 62.8 | 65.7 | 47.8 | 0.19 | 2 |
| August | 171 | 64.2 | 66.0 | 47.8 | 0.70 | 7 |
| September | 158 | 57.3 | 58.3 | 44.2 | 1.65 | 10 |
| October | 115 | 48.0 | 49.4 | 36.0 | 1.79 | 13 |
| November | 52 | 43.6 | 43.9 | 32.8 | 1.26 | 14 |
| December | 61 | 41.7 | 41.0 | 31.9 | 2.29 | 19 |
| Year [*] | 1562 | 48.2 | 49.5 | 36.8 | 20.27 | 150 |

(1) Mean of maximum and minimum.

(2) Number of days rainfall was 0.01 in. or more.

^{*}Mean or total.

ROTHAMSTED REPORT FOR 1977, PART 1

CONVERSION FACTORS

Factors for the Conversion of Imperial to Metric Units

| | |
|--------------------------------------|--|
| 1 inch (in.) | = 2.540 centimetres (cm) |
| 1 foot (ft) (=12 in.) | = 30.48 cm |
| 1 yard (yd) (=3 ft) | = 0.9144 metre (m) |
| 1 square yard (yd ²) | = 0.8361 m ² |
| 1 acre (ac) (=4840 yd ²) | = 0.4047 hectare (ha) |
| 1 ounce (oz) | = 28.35 grams (g) |
| 1 pound (lb) | = 0.4536 kilogram (kg) |
| 1 hundredweight (cwt) (=112 lb) | = 50.80 kg |
| 1 ton (=2240 lb) | = 1016 kg = 1.016 metric tons (tonnes) (t) |
| 1 pint | = 0.5682 litre (l) |
| 1 gallon (gal) (=8 pints) | = 4.546 litres |
| 1 fluid ounce = 1/20 pint | = 0.02841 litre = 28.41 ml |
| 1 cubic foot | = 28.32 litres |

| <i>To convert</i> | <i>Multiply by</i> |
|---|--------------------|
| oz ac ⁻¹ to g ha ⁻¹ | 70.06 |
| lb ac ⁻¹ to kg ha ⁻¹ | 1.121 |
| cwt ac ⁻¹ to kg ha ⁻¹ | 125.5 |
| cwt ac ⁻¹ to t ha ⁻¹ | 0.1255 |
| ton ac ⁻¹ to kg ha ⁻¹ | 2511 |
| ton ac ⁻¹ to t ha ⁻¹ | 2.511 |
| gal ac ⁻¹ to l ha ⁻¹ | 11.233 |

The following factors are accurate to about 2 parts in 100:

$$1 \text{ lb ac}^{-1} = 1.1 \text{ kg ha}^{-1}$$

$$1 \text{ gal ac}^{-1} = 11 \text{ litres ha}^{-1}$$

$$1 \text{ ton ac}^{-1} = 2.5 \text{ t ha}^{-1}$$

In general reading of the text there will be no great inaccuracy in regarding:

$$1 \text{ lb} = 0.5 \text{ kg}$$

$$1 \text{ lb ac}^{-1} = 1 \text{ kg ha}^{-1}$$

Temperatures

To convert °F into °C subtract 32 and multiply by $\frac{5}{9}$ (0.556)
 To convert °C into °F multiply by $\frac{9}{5}$ (1.8) and add 32

CONVERSION FACTORS

Factors for the Conversion of Metric to Imperial Units

| | |
|----------------------------------|--|
| 1 centimetre (cm) | = 0.3937 inch (in.) = 0.03281 ft |
| 1 metre (m) | = 1.094 yards (yd) |
| 1 square metre (m ²) | = 1.196 square yards (yd ²) |
| 1 hectare (ha) | = 2.471 acres (ac) |
| 1 gram (g) | = 0.03527 ounce (oz) |
| 1 kilogram (kg) | = 2.205 pounds (lb) |
| 1 kg | = 0.01968 hundredweight (cwt) = 0.0009842 ton |
| 1 metric ton (tonne) (t) | = 0.9842 ton |
| 1 litre | = 1.760 pints = 0.2200 gallon (gal) |
| 1 litre = 1000 millilitres (ml) | = 35.20 fluid ounces = 0.03531 cubic foot (ft ³) |

| <i>To convert</i> | <i>Multiply by</i> |
|--|--------------------|
| g ha ⁻¹ to oz ac ⁻¹ | 0.01427 |
| kg ha ⁻¹ to lb ac ⁻¹ | 0.8921 |
| kg ha ⁻¹ to cwt ac ⁻¹ | 0.007966 |
| t ha ⁻¹ to cwt ac ⁻¹ | 7.966 |
| kg ha ⁻¹ to tons ac ⁻¹ | 0.0003983 |
| t ha ⁻¹ to tons ac ⁻¹ | 0.3983 |
| l ha ⁻¹ to gal ac ⁻¹ | 0.08902 |

Plant nutrients

Plant nutrients are best stated in terms of amounts of the elements (P, K, Na, Ca, Mg, S); the old 'oxide' terminology (P₂O₅, K₂O, Na₂O, CaO, MgO, SO₃) is still used in work involving fertilisers and liming since Regulations require statements of P₂O₅, K₂O, etc.

For quick conversions

(accurate to within 2%) the following factors may be used:

| | |
|----------------------------------|---------------------------------|
| $2\frac{1}{2} \times P = P_2O_5$ | $\frac{3}{7} \times P_2O_5 = P$ |
| $1\frac{1}{2} \times K = K_2O$ | $\frac{5}{6} \times K_2O = K$ |
| $1\frac{3}{8} \times Ca = CaO$ | $\frac{7}{10} \times CaO = Ca$ |
| $1\frac{3}{4} \times Mg = MgO$ | $\frac{3}{5} \times MgO = Mg$ |

For accurate conversions:

| <i>To convert</i> | <i>Multiply by</i> | <i>To convert</i> | <i>Multiply by</i> |
|------------------------------------|--------------------|------------------------------------|--------------------|
| P ₂ O ₅ to P | 0.4364 | P to P ₂ O ₅ | 2.2915 |
| K ₂ O to K | 0.8301 | K to K ₂ O | 1.2047 |
| CaO to Ca | 0.7146 | Ca to CaO | 1.3994 |
| MgO to Mg | 0.6031 | Mg to MgO | 1.6581 |