

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1955

[Full Table of Content](#)



55/R/A/4 Barnfield - Mangolds and Sugar Beet

Rothamsted Research

Rothamsted Research (1956) *55/R/A/4 Barnfield - Mangolds and Sugar Beet* ; Yields Of The Field Experiments 1955, pp 10 - 12 - DOI: <https://doi.org/10.23637/ERADOC-1-175>

55/A/4.1

MANGOLDS AND SUGAR BEET - BARNFIELD 1955

The 80th and 10th years

For history, details of treatments etc., see "Results of the Field Experiments 1939-47" Vol. 1, Section A/5.

Because of a serious infestation of couch grass (Agropyron repens) it was decided to apply trichloroacetic acid to the following plots: all the O series, 1N. In 1955 half of each plot was sprayed at half the standard rate. No damage to the crops was noted, and the halves were not harvested separately.

Cultivations, etc.: Dung applied: Dec 30, 1954. Ploughed: Jan 3, 1955. Certain plots sprayed with trichloroacetic acid, sodium salt: at 20 lb. in 40 gallons per acre: Apr 9. Fertilizers applied: Apr 22. Seed drilled, Mangolds - 9 lb; Sugar beet - 16 lb per acre. Calcium chloride applied to 4NB only: June 2. Sprayed with miscible D.D.T. 3 pints per acre: June 11. Singling started: July 19. Top dressings applied; nitrate of soda - July 20, sulphate of ammonia - July 22. Lifting started: Mangolds, - Nov 13, Sugar beet - Nov 16. Varieties Mangolds - Yellow Globe, Sugar beet - Klein E.

Erratum to "Results of the Field Experiments" 1951, p. 51/A/4.2.
Sugar Beet: Tops. Cross Dressing - 0. Yields for strips 4-8 should read "1.65, 1.49, 1.57, 1.63, 1.79", NOT "6.61, ..., 7.14"

Summary of Results

Strip	Cross Dressing				
	O	N	A	AC	C
Mangolds, Roots: tons per acre					
1	3.55	11.93	10.20	7.44	10.34
2	5.04	9.95	7.98	7.95	10.26
4	1.75	(a) 7.34 (b) 7.29*	5.38	7.95	8.04
5	1.03	6.01	2.50	3.99	6.07
6	1.01	4.30	4.67	7.98	8.72
7	0.89	7.22	6.97	8.18	9.38
8	0.44	2.22	2.47	4.32	4.05
9	5.66				
Mangolds, Leaves: tons per acre					
1	1.47	3.64	3.27	3.00	2.69
2	1.71	3.00	3.10	3.42	2.20
4	0.94	(a) 3.40 (b) 3.13*	2.05	2.86	2.00
5	0.74	3.44	1.81	2.03	2.25
6	0.81	2.81	1.88	2.54	1.81
7	0.72	3.96	2.49	3.69	2.54
8	0.60	1.66	2.34	2.91	2.64
9	3.42				
Mangolds, Plant Number: thousands per acre					
1	19.8	23.2	21.4	16.2	22.0
2	22.2	21.5	19.8	17.6	21.4
4	22.8	(a) 24.7 (b) 23.5*	19.0	17.8	21.3
5	22.6	24.1	20.3	16.8	22.0
6	22.1	22.6	21.5	19.2	22.5
7	21.3	23.1	21.5	18.9	21.9
8	21.0	21.2	20.2	20.9	23.0
9	22.5				

*No nitrate of soda. Nitrogen applied as calcium and potassium nitrates.

55/A/4.3

Strip	Cross Dressing				
	O	N	A	AC	C
Sugar Beet, Roots (washed): tons per acre					
1	2.67	7.83	7.96	6.78	8.46
2	3.03	4.78	3.02	2.86	6.92
4	1.32	(b) 4.67*	3.33	6.56	6.59
5	1.31	3.55	2.82	4.37	6.07
6	1.27	3.10	3.29	6.14	6.66
7	1.04	4.46	4.24	6.61	5.14
8	0.90	3.04	2.58	4.76	4.04
9	2.94				
Sugar Beet, Tops: tons per acre					
1	2.39	9.28	8.94	5.96	4.93
2	3.32	7.03	4.54	3.91	5.42
4	1.15	(b) 5.67*	4.40	5.52	4.69
5	1.06	6.94	3.61	6.30	5.52
6	1.27	4.93	4.25	6.55	5.42
7	1.05	6.74	4.20	8.40	4.25
8	0.93	4.54	3.27	6.74	5.52
9	3.52				
Sugar Beet, Plant Number: thousands per acre					
1	19.6	22.3	21.5	19.6	21.9
2	22.4	22.8	23.0	17.5	22.0
4	23.1	(b) 21.4*	22.1	20.8	20.5
5	23.9	24.3	22.4	22.0	23.4
6	23.1	23.1	21.4	20.8	23.5
7	24.2	23.5	23.4	21.5	22.8
8	22.8	24.2	24.2	22.8	24.0
9	23.2				
Sugar Beet, Sugar Percentage					
1	16.1	14.9	15.3	16.0	16.2
2	15.5	13.9	14.0	14.1	15.8
4	15.8	(b) 14.3*	14.8	15.7	16.2
5	16.0	14.2	15.5	15.6	16.4
6	15.4	14.6	15.3	15.7	16.1
7	15.9	15.6	16.0	15.6	16.1
8	15.3	14.9	15.9	14.9	15.5
9	15.4				

*No nitrate of soda. Nitrogen applied as calcium and potassium nitrates.