

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1955

[Full Table of Content](#)



55/W/A/8 Stackyard Woburn - Barley

Rothamsted Research

Rothamsted Research (1956) *55/W/A/8 Stackyard Woburn - Barley* ; Yields Of The Field Experiments 1955, pp 16 - 16 - DOI: <https://doi.org/10.23637/ERADOC-1-175>

55/A/8

BARLEY - WOBURN STACKYARD 1955

For history, details of treatments etc., see "Results of the Field Experiments" 1939-47, Vol. I, Section A/7 and 1953 Section A/8. In 1955 dressings of calcium carbonate were applied ranging from 20 to 50 cwt per acre according to the acidity of the plots.

Cultivations, etc.: Ploughed: Nov 3, 1954. First dressing of chalk applied: Feb 15, 1955. Second dressing of chalk applied: Mar 8. Ploughed: Mar 18. Third dressing of chalk applied: Apr 6. Seed drilled at 3 bushels per acre: Apr 7. 'Nitrochalk' applied: Apr 12. Sprayed with MCPA amine at $2\frac{1}{2}$ pints per acre in low volume: May 22. Harvested: Aug 24. Variety: Plumage Archer.

Summary of Results

Plot	'Nitrochalk' dressing: cwt per acre	Grain: cwt per acre	Straw: cwt per acre
1	2	5.6	7.1
3	4	8.0	10.7
7	6	8.4	12.1
6	2	12.2	11.5
9	4	18.7	19.2
4	6	11.9	15.0
11b (3)	2	11.7	12.7
11b (1)	4	19.8	22.4
11b (2)	6	22.4	25.9
10b	2	6.1	7.1
11a	4	9.8	11.9
10a	6	4.4	7.8
5	2	6.9	8.1
8	4	9.6	11.3
2	6	5.6	7.9