

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1955

[Full Table of Content](#)



55/R/CA/1 Winter Wheat - Eyespot Rotation, Varieties, Seed Rates and N - 2nd Year

Rothamsted Research

Rothamsted Research (1956) *55/R/CA/1 Winter Wheat - Eyespot Rotation, Varieties, Seed Rates and N - 2nd Year* ; Yields Of The Field Experiments 1955, pp 71 - 72 - DOI:

<https://doi.org/10.23637/ERADOC-1-175>

55/Ca/1.1

WINTER WHEAT

The effects of Crop sequences, Varieties, Seed rates and Nitrogen on the incidence of Eyespot (Cercospora herpotrichoides) - Long Hoos I, II and III, 1955. The 2nd preliminary year.

For details of treatments and crop sequences etc., see "Results of the Field Experiments 1954", Section 54/Ca/2.1 with the exception that in 1955 the seed rates were changed to

Holdfast: $1\frac{1}{2}$, 3 bushels per acre

Cappelle: 2, 4 bushels per acre,

in order to give approximately the same number of seeds of each variety per acre.

Area of each plot: 0.0212 acre. Area harvested: 0.0140 acre.

Cultivations, etc.: Ploughed: Oct 8, 1954. Seed combine drilled: Oct 15. 1st application of nitrochalk: Mar 15, 1955. 2nd application: May 9. Combine harvested: Aug 19.

Standard error per plot:

Grain (at 85% D.M.): 1.92 cwt per acre or 5.8% (12 d.f.)

Records of incidence of disease (Eyespot and Take-All), estimates of % area lodged, and counts of plant, shoot and straw numbers were made.

55/Ca/1.2

Summary of Results

Response to	Responses to treatments						
	Mean	Variety		Seed rate		Nitrogen:	
		Hold- fast	Capp- elle	Single	Double	0.46	0.93

Grain (at 85% Dry Matter)

<u>Previous crop wheat:</u>	Mean yield 17.1 cwt per acre						
	(±0.96)						(±1.36)
Variety (Cappelle - Holdfast)	+6.5	-	-	+7.1	+5.9	+4.8	+8.2
Seed rate (Double - Single)	-0.8	-0.2	-1.4	-	-	-1.6	0.0
Nitrogen (0.93 - 0.46)	+3.6	+1.9	+5.3	+2.8	+4.4	-	-

<u>Previous crop potatoes:</u>	Mean yield 49.3 cwt per acre						
	(±0.96)						(±1.36)
Variety (Cappelle - Holdfast)	+12.9	-	-	+12.8	+13.0	+9.9	+15.9
Seed rate (Double - Single)	+3.3	+3.2	+3.4	-	-	+2.8	+3.8
Nitrogen (0.93 - 0.46)	+5.6	+2.6	+8.6	+5.1	+6.1	-	-

Straw

<u>Previous crop wheat:</u>	Mean yield 24.3 cwt per acre						
Variety (Cappelle - Holdfast)	+4.3	-	-	+4.1	+4.5	+8.3	+0.3
Seed rate (Double - Single)	+0.5	+0.3	+0.7	-	-	+0.7	+0.3
Nitrogen (0.93 - 0.46)	+8.1	+12.1	+4.1	+8.3	+7.9	-	-

<u>Previous crop potatoes:</u>	Mean yield 42.0 cwt per acre						
Variety (Cappelle - Holdfast)	-2.5	-	-	-4.3	-0.7	-3.2	-1.8
Seed rate (Double - Single)	+5.3	+3.5	+7.1	-	-	+2.8	+7.8
Nitrogen (0.93 - 0.46)	+4.2	+3.5	+4.9	+1.7	+6.7	-	-

General means. Grain: 33.2 cwt per acre
 Straw: 33.2 cwt per acre
 Mean dry matter % as harvested, Grain: 81.8