Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1955



Full Table of Content

55/R/CA/2 Winter Wheat - Control of Bulb Fly (Insecticides)

Rothamsted Research

Rothamsted Research (1956) 55/R/CA/2 Winter Wheat - Control of Bulb Fly (Insecticides); Yields Of The Field Experiments 1955, pp 73 - 73 - DOI: https://doi.org/10.23637/ERADOC-1-175

55/Ca/2

WINTER WHEAT

Control of wheat bulb fly by insecticides - Pennell's Piece 1955.

Design: 4 randomized blocks of 6 plots each.

Area of each plot: 0.00643 acre.

Treatments: Insecticides:
None (2 plots per block).

Seed dressed with Technical Dieldrin (97%) at 2.25% of seed

w/w using cellulose ether sticker.

4% Dieldrin dust at 1 cwt per acre combine drilled with seed.

Sprayed early with Parathion 0.05% v/v at 100 gallons per

acre.

Sprayed late with Parathion 0.05% v/v at 100 gallons per

acre.

(4)

Basal dressing per acre: 3 cwt nitrochalk, 20 cwt hydrated lime.

Note: All seed treated with organo-mercurial fungicide.

Cultivations, etc.: Ploughed (for bare fallow): Nov 2, 1953. Lime applied: Mar 3, 1954. Ploughed: May 17. Seed drilled at 2 bushels per acre: Dec 21. "3" plots sprayed with Parathion: Feb 11, 1955. "4" plots sprayed with Parathion: Apr 7. Nitrochalk applied: May 9. Combine harvested: Aug 24. Variety: Cappelle. Previous crop: Bare fallow.

Standard error per plot:
Grain (at 85% dry matter): 3.75 cwt per acre or 8.4% (16 d.f.)

Counts of numbers of plants, tillers, damaged tillers and of wheat bulb fly larvae were made.

Summary of Results

Grain (at 85% dry matter): cwt per acre

	Insecticide					
	0	1	2	3	4	Mean
Mean (±1.88)	41.0(1)	47.2	53.7	47.9	38.7	44.9
Increase (±2.30)		+6.2	+12.7	+6.9	-2.3	
(1) ±1.33						

Mean dry matter % as harvested: 84.4