


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1955

[Full Table of Content](#)



55/W/CA/4 Winter Wheat - Varieties, Seed Rates, Levels and Times of N

Rothamsted Research

Rothamsted Research (1956) *55/W/CA/4 Winter Wheat - Varieties, Seed Rates, Levels and Times of N*; Yields Of The Field Experiments 1955, pp 75 - 76 - DOI:

<https://doi.org/10.23637/ERADOC-1-175>

55/Ca/4.1

WINTER WHEAT

Varieties, seed rates, levels and times of application of N: - Woburn, Roadpiece 1955, the 2nd year.

Design: 4 randomized blocks of 8 plots each, certain high order interactions being confounded with block differences. In addition each block contained 2 plots with no nitrogen, the variety x seed rate interaction being confounded.

Area of each plot: 0.0159 acre. Area harvested: 0.0140 acre.

Treatments: All combinations of:-

Varieties: Holdfast; Cappelle.

Seed rates: Holdfast, $1\frac{1}{2}$; 3 bushels per acre.

Cappelle, 2; 4 bushels per acre.

Nitrogen: 0.5; 1.0 cwt N per acre as nitrochalk.

Time of application of N: half dressing in March and again in May; whole dressing mid March; mid April; mid May.

Basal dressing: 1 cwt per acre compound granular fertilizer (12% N, 12% P_2O_5 , 15% K_2O) combine drilled with seed.

Cultivations, etc.: Ploughed: Sept 28, 1954. Combine drilled: Oct 25. March top dressing applied: Mar 15, 1955. April top dressing applied: Apr 20. All plots sprayed with D.N.O.C. at $1\frac{1}{2}$ gallons in 80 gallons: May 19. May top dressing applied: May 24. Combine harvested: Aug 22. Varieties: Holdfast and Cappelle. Previous crop: Wheat.

Note (1) The experiment is a repetition on the same plots of the one carried out in 1954 (see "Results of the Field Experiments 1954", Section 54/Ca/7). There were minor changes in the treatments but the same randomization was used.

(2) Records of incidence of disease (Take-all and Eyespot) and weeds, and counts of plant, shoot and ear numbers were made.

Standard error per plot.

Grain: 3.95 cwt per acre or 26.9% (12 d.f.)

Summary of Results

Grain: cwt per acre

	T ₁	T ₂	T ₃	T ₄	Mean
Mean (± 1.40)	16.5	16.9	18.9	12.2	16.1
(±1.97)					(±0.99)
V ₁	17.0	15.6	18.3	12.4	15.8
V ₂	16.0	18.3	19.6	12.0	16.5
Difference (± 2.79)	-1.0	+2.7	+1.3	-0.4	+0.7 (± 1.40)
R ₁	14.1	16.3	16.8	9.6	14.2
R ₂	18.9	17.6	21.1	14.8	18.1
Difference (± 2.79)	+4.8	+1.3	+4.3	+5.2	+3.9 (± 1.40)
N ₁	14.9	11.9	15.2	12.2	13.6
N ₂	18.1	22.0	22.6	12.1	18.7
Difference (± 2.79)	+3.2	+10.1	+7.4	-0.1	+5.1 (± 1.40)

	R ₁	R ₂	N ₀	N ₁	N ₂	Mean
Mean (± 0.99)			(±1.40) 8.8	(±0.99) 13.6	18.7	14.7
(±1.40)			(±1.97)	(±1.40)	(±0.88)	
V ₁	13.7	18.0	8.8	13.6	18.0	14.4
V ₂	14.7	18.2	8.8	13.5	19.4	14.9
R ₁			8.9	11.9	16.5	13.1
R ₂			8.6	15.3	20.9	16.2

Mean dry matter % as harvested: 85.5

Treatments

V₁ Holdfast R₁, R₂ 1½, 3 bushels per acre N₀ No N
 V₂ Cappelle R₁, R₂ 2, 4 bushels per acre N₁ 0.46 cwt N per acre
 N₂ 0.93 cwt N per acre

T₁ Nitrochalk half in March half in May T₃ Nitrochalk all in mid April
 T₂ Nitrochalk all in mid March T₄ Nitrochalk all in mid May

The V × R table does not include the plots receiving no nitrogen.