


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1955

[Full Table of Content](#)



55/R/CC/2 Spring Beans - Control of Weeds (Spraying and Cultivations)

Rothamsted Research

Rothamsted Research (1956) *55/R/CC/2 Spring Beans - Control of Weeds (Spraying and Cultivations)* ; Yields Of The Field Experiments 1955, pp 89 - 90 - DOI:
<https://doi.org/10.23637/ERADOC-1-175>

55/Cc/2.1

SPRING BEANS

Control of weeds by spraying and cultivation - Fosters 1955.

Design: 4 randomized blocks of 5 plots each, 2 blocks being sprayed and 2 unsprayed.

Area of each plot: 0.0202 acre. Area harvested: 0.0126 acre.

Treatments: All combinations of:-

Spraying (on blocks): None; DNBP high volume

Additional cultivations: None (0)

Harrowed once (1)

Harrowed three times (2)

Mechanical weeder once (3)

Mechanical weeder three times (4)

Basal dressing: 6 cwt compound granular fertilizer (10% P₂O₅, 20% K₂O) per acre.

Basal inter-row cultivation: One early horse hoeing.

Cultivations, etc.: Ploughed: Oct 22, 1954. Basal fertilizer applied, seed drilled at 195 lb per acre: Mar 19, 1955. Horse hoed all plots: May 11. Combine harvested: Aug 27. Variety: Spring Tick. Previous crop: Barley.

Treatment cultivations:

(1) Harrowed: April 30.

(2) Harrowed: April 30, May 6, May 31.

(3) Mechanical weeder: April 30.

(4) Mechanical weeder: April 30, May 6, May 31.

DNBP at 4 pts in 100 gallons per acre sprayed: May 26.

Standard error per plot:

Grain: 1.07 cwt per acre or 6.3% (8 d.f.)

55/Cc/2.2

Summary of Results

Grain: cwt per acre

Spray	Treatment cultivations					Mean
	0	1	2	3	4	
			(± 0.76) [Ⓜ]			
None	18.5	17.5	14.3	18.5	16.4	17.0
DNBP	17.1	17.6	15.1	17.4	17.7	17.0
Mean (± 0.54)	17.8	17.5	14.7	18.0	17.1	17.0
Difference (± 1.07) ⁺	-1.4	+0.1	+0.8	-1.1	+1.3	0.0

[Ⓜ]for use in horizontal comparisons only

⁺for use in the comparison of two differences only

Mean dry matter % as harvested: 86.2

Treatment cultivations

- (0) No extra cultivation.
- (1) Harrowed once.
- (2) Harrowed three times.
- (3) Mechanical weeder once.
- (4) Mechanical weeder three times.