


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1955

[Full Table of Content](#)



55/R/A/3 Agdell - Rotation

Rothamsted Research

Rothamsted Research (1956) *55/R/A/3 Agdell - Rotation* ; Yields Of The Field Experiments 1955, pp 9 - 9 - DOI: <https://doi.org/10.23637/ERADOC-1-175>

55/A/3

WHEAT AFTER FALLOW - HOOSFIELD 1955

Without manure 1851 and since

For history, details of treatments etc. see "Results of the Field Experiments" 1939-47 Vol.I, Section A/3.

Cultivations, etc.:

Cropped plots. Ploughed: Sept 25, 1954. Seed drilled at 3 bushels per acre: Dec 7. Harvested: Aug 20, 1955. Variety: Squareheads Master 13/4.

Fallowed plots. Ploughed: Sept 25, 1954 and Sept 28, 1955.

Summary of Results

Mean yields: cwt per acre

Plot	B3	B4	B1	
No. of years of fallow	1	1	3	Mean
Grain	9.4	10.2	11.2	10.3
Straw	12.9	12.3	15.3	13.5

Mean dry matter % as threshed, Grain: 85.3
Straw: 86.2

SPRING WHEAT - AGDELL 1955

For history, details of treatments etc., see "Results of the Field Experiments" 1939-47 Vol.I, Section A/4, and 1954, page 54/A/3. In 1955 spring wheat was grown with uniform treatment, manured with nitrogen only. Yields were not recorded. Detailed sampling both of crops and soils was carried out.

Cultivations, etc.:

Ploughed: Nov 4, 1954. Seed drilled at 2 bushels per acre: Apr 4, 1955. Basal dressing of nitrochalk, 4 cwt per acre applied: Apr 5. Sprayed with MCPA at 2½ pts. in 18 gallons per acre: May 23. Combine harvested: Sept 2. Variety: Koga II.